

Acta Biologica Hungarica 66(3), pp. 339–347 (2015)
DOI: 10.1556/018.66.2015.3.9

IDENTIFICATION OF *ASPERGILLUS* SPECIES IN CENTRAL EUROPE ABLE TO PRODUCE G-TYPE AFLATOXINS

NIKOLETT BARANYI,¹ DANIELA JAKŠIĆ DESPOT,² ANDREA PALÁGYI,¹ NOÉMI KISS,¹
SÁNDOR KOCSUBÉ,¹ ANDRÁS SZEKERES,¹ ANITA KECSKEMÉTI,¹ OTTÓ BENCSEK,¹
CSABA VÁGVÖLGYI,¹ MAJA ŠEGVIĆ KLARIĆ² and JÁNOS VARGA^{1*}

¹Department of Microbiology, Faculty of Science and Informatics, University of Szeged,
Közép fasor 52, H-6726 Szeged, Hungary

²Department of Microbiology, Faculty of Pharmacy and Biochemistry, University of Zagreb,
Schrottova 39, 10000 Zagreb, Croatia

(Received: January 8, 2015; accepted: February 9, 2015)

The occurrence of potential aflatoxin producing fungi was examined in various agricultural products and indoor air in Central European countries including Hungary, Serbia and Croatia. For species identification, both morphological and sequence based methods were applied. *Aspergillus flavus* was detected in several samples including maize, cheese, nuts, spices and indoor air, and several isolates were able to produce aflatoxins. Besides, three other species of *Aspergillus* section *Flavi*, *A. nomius*, *A. pseudonomius* and *A. parasiticus* were also isolated from cheese, maize and indoor air, respectively. This is the first report on the occurrence of *A. nomius* and *A. pseudonomius* in Central Europe. All *A. nomius*, *A. pseudonomius* and *A. parasiticus* isolates were able to produce aflatoxins B₁, B₂, G₁ and G₂. The *A. nomius* isolate came from cheese produced very high amounts of aflatoxins (above 1 mg ml⁻¹). All *A. nomius*, *A. pseudonomius* and *A. parasiticus* isolates produced much higher amounts of aflatoxin G₁ than aflatoxin B₁. Further studies are in progress to examine the occurrence of producers of these highly carcinogenic mycotoxins in agricultural products and indoor air in Central Europe.

Keywords: Aflatoxins – *Aspergillus* – cheese – indoor air – sequence-based identification

INTRODUCTION

Aflatoxins are decaketide-derived secondary metabolites which are produced through a complex biosynthetic pathway. These compounds exhibit hepatocarcinogenic and hepatotoxic properties, and referred to as the most harmful naturally occurring carcinogens [3, 40]. From this reason, aflatoxins are among the economically most important mycotoxins produced by various *Aspergillus* species mainly belonging to *Aspergillus* section *Flavi* [40]. The most important producers of aflatoxins are *A. flavus*, *A. parasiticus* and *A. nomius* [40]. While *A. flavus* produces B-type aflatoxins, *A. nomius* and *A. parasiticus* are able to produce also G-type aflatoxins [40, 41]. *Aspergillus flavus* and *A. parasiticus* are identified worldwide from various substrates [40]. *Aspergillus nomius* is an aflatoxin producing filamentous fungus which was described only in 1987 [21]. Isolates of this species were originally isolated from

*Corresponding author; e-mail address: jvarga@bio.u-szeged.hu

insects and wheat in the USA, and *Cycas* sp. from Polinesia [21]. Later it was also identified in various agricultural soils in the USA, and the presence of the sexual cycle was also confirmed in this species [16]. Recently, a species closely related to *A. nomius* named *A. pseudonomius* has also been described [41], which was found to be an important source of aflatoxin contamination on Brazil nuts [23]. However, to our knowledge, these species have never been reported from Central Europe.

In this study, we examined the occurrence of aflatoxin-producing species in various food products and indoor air in Central Europe, including Hungary, Serbia and Croatia, using sequence-based and analytical methods.

MATERIALS AND METHODS

Chemicals

Methanol, acetonitrile, dichloromethane, ethyl-acetate and n-hexane for sample preparation and eluents were purchased from VWR (Hungary). The formic acid, trifluoroacetic acid and aflatoxins B₁, B₂, G₁, G₂ standards were obtained from Sigma-Aldrich (Budapest, Hungary). Deionized water both for sample preparation and high-performance liquid chromatography (HPLC) runs was produced by Millipore Milli-Q Gradient A10 water purification equipment (Millipore, Hungary).

Sample collection

The cheese samples (n = 15), surface sterilized maize samples (n = 80), onions (n = 16), nuts (n = 7) and spice samples (n = 16) were plated onto dichloran rose bengal chlortetracycline (DRBC) media [19]. Indoor air in Croatian apartments (n = 50) was sampled using a Mas-100 Eco microbiological air sampler (Merck, Darmstadt, Germany) with 400 holes (hole to agar impactor) and dichloran 18% glycerol agar (DG-18) plates [27]. In Hungary (n = 26), the plate sedimentation method was used [34]. Plates were incubated at 25 °C in darkness and monitored periodically for characteristic mycelium growing from the kernels. Outgrowing mycelia were purified and transferred to malt extract agar (MEA). The isolates were subcultured on malt extract agar (MEA) and Czapek yeast autolysate (CYA) media for morphological identification [34]. Assignment to *Aspergillus* section *Flavi* was confirmed after subculturing on CYA, MEA and Potato Dextrose agar (PDA), according to standard procedures [9, 28, 31].

Genotypic studies

Isolation of genomic DNA from mycelia grown in liquid YPD medium (1% Bacto yeast extract, 1% Bacto peptone, 1% D-glucose) for five days were performed by the

Masterpure™ yeast DNA purification kit (Epicentre Biotechnologies, Madison, WI, USA) according to the manufacturer's instructions. A fragment of the calmodulin gene was amplified with primers cmd5 and cmd6 as described by Hong et al. [15]. DNA sequences were determined at Agowa GmbH (Berlin, Germany). Sequence analysis was performed by nucleotide-nucleotide BLAST similarity search at the website of the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/BLAST>) [1], and sequences were also compared with our own sequence database. Species identification was determined from the lowest expect value of the BLAST output. Partial calmodulin sequences were deposited in the GenBank sequence database under accession numbers KP310011-KP310014.

Examination of aflatoxin producing abilities

During the examination of aflatoxin production the *A. parasiticus* isolates were cultivated on Czapek yeast autolysate (CYA) agar in darkness at 25 °C for 15 days. Aflatoxin extraction was performed from 4 agar plugs (in diameter of 6 mm) with 1 ml of mixture of methanol/dichloromethane/ethyl-acetate (1/2/3, v/v/v) supplemented with 1% of formic acid [37]. The extracts were centrifuged at 10,000 g for 10 min and the organic phases were removed and evaporated to dryness under slight stream of nitrogen. For the measurement of aflatoxins, the *A. nomius* and *A. pseudonomius* isolates were cultivated in darkness for 7 days on 2 ml of yeast extract – sucrose (YES) media at 25 °C [34, 38]. Aflatoxin extraction was performed with 2 ml of dichloromethane, the extracts were centrifuged at 10,000 g for 10 min and the organic phases were dried.

After the evaporation the samples were derivatized with trifluoroacetic acid as described previously [12]. For the analysis the modular HPLC system (Shimadzu, Japan) equipped with a DGU-14A vacuum degasser, two LC-20AD pumps, SIL-20A autosampler, CBM-20A column thermostat RF-20A fluorescence detector and CBM-20A system controller, which was controlled by ClassVP 6.2 software. Separations were achieved on a Lichocart 250×4 mm, 5 μm (Merck, Hungary) column with coupled with a Lichospher 100 RP-18 (Merck, Hungary) guard column and the injection volume was 5 μl. The composition of the isocratic mobile phase were water/methanol/acetonitrile 65/17.5/17.5 (V/V/V), and the flow rate was maintained at 1 ml/min, while column temperature was 40 °C. The fluorescence detector was set to an excitation wavelength of 365 nm and an emission wavelength of 430 nm. For the quantification of aflatoxins, linear calibration curves were used in the concentration range of 50–1000 ng/ml in the case of aflatoxins B₁ and G₁ and 14–722 ng/ml for aflatoxins B₂ and G₂, and the equations of resulted calibration curves were $y = 0.0049x + 14.8$, $y = 0.0012x + 15.6$, $y = 0.0034x + 4.2$ and $y = 0.0008x + 4.5$ for aflatoxins G₁, B₁, G₂ and B₂, respectively.




Fig. 1. Colony morphologies on PDA plates (A, D), light-microscopic pictures (B, E), and HPLC chromatograms of aflatoxins (C, F) of *A. nomius* SZMC 22631 and *A. pseudonomius* SZMC 22273, respectively

RESULTS AND DISCUSSION

Large numbers of samples were analyzed from maize, dried fruits, cheese, nuts, wheat and spices (Hungary), from maize (Serbia), and from indoor air (Croatia, Hungary) [42]. Species identifications were carried out using sequence analysis of part of the calmodulin gene. This gene is suitable for sequence-based identification of isolates belonging to *Aspergillus* section *Flavi* [41]. Potential aflatoxin producing species were not identified on dried fruits including raisins, figs and dates (data not shown). However, aflatoxigenic species have been found in the remaining samples (Table 1). *Aspergillus flavus* was identified in many of the samples (Table 1), while *A. nomius* was detected in only one sample of cheese from Hungary (Fig. 1). Both the morphological features and calmodulin sequence data indicated that this isolate belongs to *A. nomius* (Fig. 1). To our knowledge, this is the first report on the occurrence of *A. nomius* in Central Europe. This species was originally isolated from mouldy wheat in the USA, and later from various substrates in India, Japan and Thailand. Recently, Olsen et al. [26] and several other authors have observed that *A. nomius* is an important producer of aflatoxins in Brazil nuts [2, 5, 14, 33], on cocoa [7], almond [17], Indonesian nutmegs [25], maize, rice, black beans and cassava from Thailand [29, 30], and from various soil samples in Thailand, Iran and Japan [11, 20, 32]. It was also identified in house dust samples from Mexico [43]. This species has also been found to be able to cause human infections including onychomycosis in Italy [44], keratitis in India [22], invasive aspergillosis, cavitary and fibrosing pulmonary and pleural aspergillosis in Hong Kong [39] and breakthrough pneumonia in Italy [4]. Besides, it was also identified as the causative agent of stonebrood disease in honeybees in Great Britain [13].

Another species identified in the samples was *A. pseudonomius*, which was described recently in insects and soil from the USA [41], and was found to be partly responsible for aflatoxin contamination of Brazil nuts in Brazil [23]. Recently this species was also identified from house dust samples in Thailand and Micronesia [43].

Table 1
Occurrence of *Aspergillus* isolates from various substrates

Source	Number of <i>Aspergillus</i> isolates	Isolates of <i>Aspergillus</i> section <i>Flavi</i>	<i>A. flavus</i>	<i>A. nomius</i>	<i>A. pseudonomius</i>	<i>A. parasiticus</i>
Maize	258	111	110	0	1	0
Cheese	5	1	0	1	0	0
Nuts	18	12	12	0	0	0
Onion	105	0	0	0	0	0
Spices	28	5	5	0	0	0
Indoor air	171 (Croatia), 42 (Hungary)	67 (Croatia), 6 (Hungary)	65 (Croatia), 6 (Hungary)	0	0	2 (Croatia), 0 (Hungary)

Table 2
Aflatoxin producing abilities of the identified *A. parasiticus*, *A. nomius* and *A. pseudonomius* isolates

	Aflatoxin B ₁ (µg ml ⁻¹)	Aflatoxin B ₂ (µg ml ⁻¹)	Aflatoxin G ₁ (µg ml ⁻¹)	Aflatoxin G ₂ (µg ml ⁻¹)	Total aflatoxins (µg ml ⁻¹)
<i>A. parasiticus</i> MFBF10887 (SZMC 22727) ^a	18.16	0.28	194.61	3.71	216.76
<i>A. parasiticus</i> MFBF 11062B (SZMC 22728)	9.95	0.17	129.42	2.01	141.54
<i>A. nomius</i> III/4/2 (SZMC 22631)	370.88	9.89	814.44	19.63	1214.84
<i>A. pseudonomius</i> NS 640 B/22 (SZMC 22273)	283.95	5.89	446.93	7.76	744.53

^aMFBF: Microbial Collection, Department of Microbiology, Faculty of Pharmacy and Biochemistry, University of Zagreb, Croatia; SZMC: Szeged Microbiological Collection, Szeged, Hungary

In this study, *A. pseudonomius* was found in maize samples collected in Serbia. This is the first report on the occurrence of this species in Europe.

Besides, another species able to produce both B and G-type aflatoxins, *A. parasiticus* was also detected in indoor samples from Croatia, which species was also detected previously in similar habitats [8, 27] (Table 1). However, the isolates were previously identified using only morphological methods. More recently, *A. parasiticus* was also detected in wheat in Slovakia [10], and in maize in Hungary [36].

Regarding the mycotoxin producing abilities of the isolates, the *A. nomius*, *A. pseudonomius* and *A. parasiticus* isolates were able to produce aflatoxins B₁, B₂, G₁ and G₂ in different amounts (Table 1, Fig. 1). The amounts observed were extremely high in the case of the *A. nomius* isolate came from cheese (Table 1, Fig. 1). Usually aflatoxin B₁ concentrations exceed that of aflatoxin G₁ in these species [18]. However, in our studies, all the *A. parasiticus*, *A. nomius* and *A. pseudonomius* isolates produced higher amounts of aflatoxin G₁ than aflatoxin B₁ at 25 °C (Table 2). These data are in agreement with some previous studies [6, 24, 35]. These studies indicated that the ratio of aflatoxins B and G is greatly influenced by temperature. Further studies are needed to clarify the significance of these findings.

CONCLUSIONS

Large numbers of agricultural products and indoor air samples were analyzed in this study to clarify the species distribution of potential aflatoxin producing fungi in our region. Besides *A. flavus*, *A. parasiticus* was also identified in indoor samples from

Croatia, *A. nomius* from cheese from Hungary, and *A. pseudonomius* from maize from Serbia. All four isolates were able to produce aflatoxins B₁, B₂, G₁ and G₂. *Aspergillus pseudonomius*, *A. nomius* and *A. parasiticus* isolates produced aflatoxin G₁ in larger quantities than aflatoxin B₁ under the conditions used, while the *A. nomius* isolate produced these mycotoxins in very high amounts. Further studies are in progress to examine the occurrence of producers of these highly carcinogenic mycotoxins in agricultural products and indoor air in Central Europe.

ACKNOWLEDGEMENTS

Part of the work presented was supported by OTKA grant No. K84077, and by the European Union through the Hungary–Serbia IPA Cross-border Cooperation Program (ToxFreeFeed, HU-SRB/1002/122/062). This research was realized in the frames of TÁMOP 4.2.4. A/2-11-1-2012-0001 “National Excellence Program – Elaborating and operating an inland student and researcher personal support system convergence program”. The project was subsidized by the European Union and co-financed by the European Social Fund.

REFERENCES

1. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., Lipman, D. J. (1990) Basic local alignment search tool. *J. Mol. Biol.* 215, 403–410.
2. Baquião, A. C., de Oliveira, M. M., Reis, T. A., Zorzete, P., Diniz Atayde, D., Correa, B. (2013) Polyphasic approach to the identification of *Aspergillus* section *Flavi* isolated from Brazil nuts. *Food Chem.* 139, 1127–1132.
3. Baranyi, N., Kocsubé, S., Vágvölgyi, C., Varga, J. (2013) Current trends in aflatoxin research. *Acta Biol. Szeged.* 57, 95–107.
4. Caira, M., Posteraro, B., Sanguinetti, M., de Carolis, E., Leone, G., Pagano, L. (2012) First case of breakthrough pneumonia due to *Aspergillus nomius* in a patient with acute myeloid leukemia. *Med. Mycol.* 50, 746–750.
5. Calderari, T. O., Iamanaka, B. T., Frisvad, J. C., Pitt, J. I., Sartori, D., Pereira, J. L., Fungaro, M. H., Taniwaki, M. H. (2013) The biodiversity of *Aspergillus* section *Flavi* in brazil nuts: from rainforest to consumer. *Int. J. Food Microbiol.* 160, 267–272.
6. Carbone, I., Jakobek, J. L., Ramirez-Prado, J. H., Horn, B. W. (2007) Recombination, balancing selection and adaptive evolution in the aflatoxin gene cluster of *Aspergillus parasiticus*. *Mol. Ecol.* 16, 4401–4417.
7. Copetti, M. V., Iamanaka, B. T., Pereira, J. L., Fungaro, M. H., Taniwaki, M. H. (2011) Aflatoxigenic fungi and aflatoxin in cocoa. *Int. J. Food Microbiol.* 148, 141–144.
8. Cvetnić, Z., Pepeljnjak, S. (1995) Aflatoxin producing potential of *Aspergillus flavus* and *A. parasiticus* isolated from samples of smoked dried meat. *Die Nahrung* 39, 302–307.
9. de Hoog, G. S., Guarro, J., Gene, J., Figueras, M. J. (2000) *Atlas of Clinical Fungi*. 2nd ed. Centraalbureau voor Schimmelcultures, Utrecht/Universitat Rovira i Virgili, Reus.
10. Dovičičová, M., Tančinová, D., Labuda, R., Sulyok, M. (2012) *Aspergillus parasiticus* from wheat grain of Slovak origin and its toxigenic potency. *Czech J. Food Sci.* 30, 483–487.
11. Ehrlich, K. C., Kobbeman, K., Montalbano, B. G., Cotty, P. J. (2007) Aflatoxin-producing *Aspergillus* species from Thailand. *Int. J. Food Microbiol.* 114, 153–159.
12. Fazekas, B., Tar, A., Kovacs, M. (2005) Aflatoxin and ochratoxin A content of spices in Hungary. *Food Addit. Contam.* 22, 856–863.

13. Foley, K., Fazio, G., Jensen, A. B., Hughes, W. O. (2014) The distribution of *Aspergillus* spp. opportunistic parasites in hives and their pathogenicity to honey bees. *Vet. Microbiol.* 169, 203–210.
14. Gonçalves, J. S., Ferracin, L. M., Carneiro Vieira, M. L., Iamanaka, B. T., Taniwaki, M. H., Pelegrinelli Fungaro, M. H. (2012) Molecular analysis of *Aspergillus* section *Flavi* isolated from Brazil nuts. *World J. Microbiol. Biotechnol.* 28, 1817–1825.
15. Hong, S. B., Cho, H. S., Shin, H. D., Frisvad, J. C., Samson, R. A. (2006) Novel *Neosartorya* species isolated from soil in Korea. *Int. J. Syst. Evol. Microbiol.* 56, 477–486.
16. Horn, B. W., Moore, G. G., Carbone, I. (2011) Sexual reproduction in aflatoxin-producing *Aspergillus nomius*. *Mycologia* 103, 174–183.
17. Kenjo, T., Ishide, Y., Aoyama, K., Ichinoe, M. (2007) Fungal population and distribution of aflatoxinogenic fungi in commercial almond powder products. *Shokuhin Eiseigaku Zasshi* 48, 90–96. (in Japanese)
18. Kensler, T. W., Roebuck, B. D., Wogan, G. N., Groopman, J. D. (2011) Aflatoxin: a 50-year odyssey of mechanistic and translational toxicology. *J. Toxicol. Sci.* 120, S28–S48.
19. King, A. D. Jr., Hocking, A. D., Pitt, J. I. (1979) Dichloran-rose bengal medium for enumeration and isolation of molds from foods. *Appl. Environ. Microbiol.* 37, 959–964.
20. Kumeda, Y., Asao, T., Takahashi, H., Ichinoe, M. (2003) High prevalence of B and G aflatoxin producing fungi in sugarcane fields in Japan: heteroduplex panel analysis identifies a new genotype within *Aspergillus* section *Flavi* and *Aspergillus nomius*. *FEMS Microbiol. Ecol.* 45, 229–238.
21. Kurtzman, C. P., Horn, B. W., Hesseltine, C. W. (1987) *Aspergillus nomius*, a new aflatoxin-producing species related to *Aspergillus flavus* and *Aspergillus tamarii*. *Antonie Van Leeuwenhoek* 53, 147–158.
22. Manikandan, P., Varga, J., Kocsubé, S., Samson, R. A., Anita, R., Revathi, R., Dóczy, I., Németh, T. M., Narendran, V., Vágvölgyi, C., Manoharan, C., Kredics, L. (2009) Mycotic keratitis due to *Aspergillus nomius*. *J. Clin. Microbiol.* 47, 3382–3385.
23. Massi, F. P., Vieira, M. L., Sartori, D., Penha, R. E., de Freitas Munhoz, C., Ferreira, J. M., Iamanaka, B. T., Taniwaki, M. H., Frisvad, J. C., Fungaro, M. H. (2014) Brazil nuts are subject to infection with B and G aflatoxin-producing fungus, *Aspergillus pseudonomius*. *Int. J. Food Microbiol.* 186, 14–21.
24. Matumba, L., Sulyok, M., Njoroge, S. M., Njumbe Ediage, E., Van Poucke, C., De Saeger, S., Krska, R. (2014) Uncommon occurrence ratios of aflatoxin B₁, B₂, G₁, and G₂ in maize and groundnuts from Malawi. *Mycotoxin Res.* 31, 57–62.
25. Okano, K., Tomita, T., Ohzu, Y., Takai, M., Ose, A., Kotsuka, A., Ikeda, N., Sakata, J., Kumeda, Y., Nakamura, N., Ichinoe, M. (2012) Aflatoxins B and G contamination and aflatoxinogenic fungi in nutmeg. *Shokuhin Eiseigaku Zasshi* 53, 211–216.
26. Olsen, M., Johnsson, P., Möller, T., Paladino, R., Lindblad, M. (2008) *Aspergillus nomius*, an important aflatoxin producer in Brazil nuts? *World Mycotoxin J.* 1, 123–126.
27. Pepeljnjak, S., Šegvić, M. (2004) An overview of mycotoxins and toxigenic fungi in Croatia. In: Logrieco, A., Visconti, A. (eds) *An Overview on Toxigenic Fungi and Mycotoxins in Europe*. Kluwer Academic Publishers, Dordrecht, pp. 33–50.
28. Pitt, J. I., Hocking, A. D. (2009) *Fungi and Food Spoilage*. Springer Science+Business Media LLC, New York (NY).
29. Pitt, J. I., Hocking, A. D., Bhudhasamai, K., Miscamble, B. F., Wheeler, K. A., Tanboon-Ek, P. (1993) The normal mycoflora of commodities from Thailand. 1. Nuts and oilseeds. *Int. J. Food Microbiol.* 20, 211–226.
30. Pitt, J. I., Hocking, A. D., Bhudhasamai, K., Miscamble, B. F., Wheeler, K. A., Tanboon-Ek, P. (1994) The normal mycoflora of commodities from Thailand. 2. Beans, rice, small grains and other commodities. *Int. J. Food Microbiol.* 23, 35–53.
31. Raper, K. B., Fennell, D. I. (1965) *The Genus Aspergillus*. Williams & Wilkins, Baltimore, US.
32. Razzaghi-Abyaneh, M., Shams-Ghahfarokhi, M., Allameh, A., Kazeroon-Shiri, A., Ranjbar-Bahadori, S., Mirzahoseini, H., Rezaee, M. B. (2006) A survey on distribution of *Aspergillus* section *Flavi* in corn field soils in Iran: population patterns based on aflatoxins, cyclopiazonic acid and sclerotia production. *Mycopathologia* 161, 183–192.

33. Reis, T. A., Baquião, A. C., Atayde, D. D., Grabarz, F., Corrêa, B. (2014) Characterization of *Aspergillus* section *Flavi* isolated from organic Brazil nuts using a polyphasic approach. *Food Microbiol.* 42, 34–39.
34. Samson, R. A., Hoekstra, E. S., Lund, F., Filtenborg, O., Frisvad, J. C. (2004) *Introduction to Food- and Airborne Fungi*. CBS Fungal Biodiversity Center, Utrecht, Netherlands.
35. Schmidt-Heydt, M., Rüfer, C. E., Abdel-Hadi, A., Magan, N., Geisen, R. (2010) The production of aflatoxin B₁ or G₁ by *Aspergillus parasiticus* at various combinations of temperature and water activity is related to the ratio of *aflS* to *aflR* expression. *Mycotoxin Res.* 26, 241–246.
36. Sebök, F., Dobolyi, C., Hartman, M., Risa, A., Kritafon, C., Szoboszlai, S., Kriszt, B. (2014) Occurrence of potentially aflatoxin producing *Aspergillus* species in maize fields of Hungary. In: *Abstract Book of the Meeting of the Hungarian Microbiological Society and the EU FP7 PROMISE Regional Meeting*, pp. 62–63.
37. Smedsgaard, J. (1997) Micro-scale extraction procedure for standardized screening of fungal metabolite production in cultures. *J. Chromatogr. A* 760, 264–270.
38. Stubblefield, R. D., Shotwell, O. L. (1977) Reverse phase analytical and preparative high pressure liquid chromatography of aflatoxins. *J. Assoc. Off. Anal. Chem.* 60, 784–790.
39. Tam, E. W., Chen, J. H., Lau, E. C., Ngan, A. H., Fung, K. S., Lee, K. C., Lam, C. W., Yuen, K. Y., Lau, S. K., Woo, P. C. (2014) Misidentification of *Aspergillus nomius* and *Aspergillus tamarii* as *Aspergillus flavus*: characterization by internal transcribed spacer, β -tubulin, and calmodulin gene sequencing, metabolic fingerprinting, and matrix-assisted laser desorption ionization-time of flight mass spectrometry. *J. Clin. Microbiol.* 52, 1153–1160.
40. Varga, J., Frisvad, J. C., Samson, R. A. (2009) A reappraisal of fungi producing aflatoxins. *World Mycotoxin J.* 2, 263–277.
41. Varga, J., Frisvad, J. C., Samson, R. A. (2011) Two new aflatoxin producing species, and an overview of *Aspergillus* section *Flavi*. *Stud. Mycol.* 69, 57–80.
42. Varga, J., Kerepes, L., Kocsubé, S., Szigeti, G., Baranyi, N., Vágvölgyi, C. (2013) Examination of the diversity of indoor molds in a Hungarian student hostel. *Acta Biol. Szeged.* 57, 21–24.
43. Visagie, C. M., Hirooka, Y., Tanney, J. B., Whitfield, E., Mwange, K., Meijer, M., Amend, A. S., Seifert, K. A., Samson, R. A. (2014) *Aspergillus*, *Penicillium* and *Talaromyces* isolated from house dust samples collected around the world. *Stud. Mycol.* 78, 63–139.
44. Zotti, M., Machetti, M., Persi, A., Barabino, G., Parodi, A. (2011) Onychomycosis: first case due to *Aspergillus nomius*. *Acta Derm. Venereol.* 91, 591–592.