

# Extremal Fuzzy Space

**Khairia Mohamed Mira**

Department of Mathematics, Faculty of Science, Tripoli University, Libya.

Email: [khairiamera@yahoo.co.uk](mailto:khairiamera@yahoo.co.uk)

**Khadiga Salem Abdeen**

Department of Mathematics, Faculty of Science, Tripoli University, Libya.

Email: [khadiga@yahoo.co.uk](mailto:khadiga@yahoo.co.uk)

Received: April 21, 2021; Accepted: May 10, 2021; Published: May 17, 2021

Copyright © 2021 by author(s) and Scitech Research Organisation(SRO).

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>

---

## Abstract

The extremal topology on an arbitrary set  $X$  was defined as a maximal non-discrete topology [1] and [2]. In this paper we introduce an extremal topology on a fuzzy set  $X$ , which is also a maximal non-discrete topology, and it has to be in a specific form. This form depends on some ultrafilters  $\mathfrak{S}$ . We consider some properties for this kind of topologies when  $\mathfrak{S}$  is free. The subspaces and the base of this topology is also considered.

## Keywords

An extremal topology; Fuzzy space; ultrafilter; door space.

---

## 1. Introduction

If  $X$  is a set, in 1991, the extremal topology was defined by Papazyan, see [1], which is a maximal non-discrete topology  $\tau$  on  $X$ , so every topology which is strictly finer than  $\tau$  is discrete.

In 1965 Zadeh [6], introduced the fundamental concept of fuzzy sets, which formed the backbone of fuzzy mathematics. After that some studies presented by Chang and Wong investigated the basic concepts and general properties of fuzzy topologies.

On the other side, some studies are presented by, for instance, Sola and me about the concept of the extremal topologies.

In this paper we will consider an extremal topology on a fuzzy set  $X$  and prove that this topology has to be in a specific form  $\tau_{x_0}$ , for each  $x_0 \in X$ . After that we will consider some topological properties on it.

## 2 Preliminaries

The material of this section mostly comes from [6], [7], [10], [8] and [11].

**Definition 2.1.** An extremal topology on a nonempty set  $X$  is a maximal non-discrete topology  $\tau$ .

Let  $X$  be non-empty set and  $I = [0, 1]$  unit interval, we denoted all functions from  $X$  to  $I$  by  $I^X$ .

**Definition 2.2.** A Fuzzy set  $A$  in  $X$  is every element of  $I^X$ , so a fuzzy set  $A$  in  $X$  is characterized by a membership function  $\mu_A : X \rightarrow I$  which associates with each point  $x \in X$  and it is denoted by  $A = \{(x, \mu_A(x)) : x \in X\} \subseteq X \times I$ .

For fuzzy sets  $A$  and  $B$  in  $X$  and  $x \in X$ , we have

- $A = B \iff \mu_A(x) = \mu_B(x)$  .
- $C = A \cup B \iff \mu_C(x) = \max\{\mu_A(x), \mu_B(x)\}$ .
- $C = A \cap B \iff \mu_C(x) = \min\{\mu_A(x), \mu_B(x)\}$ .
- $\mu_{A^c}(x) = 1 - \mu_A(x)$ .

More generally, for a family of fuzzy sets,  $\wp = \{A_i : i \in I\}$ , the union  $C = \cup_i A_i$  , and the intersection,  $D = \cap_i A_i$  , are defined by

$$\mu_C(x) = \sup\{\mu_{A_i}(x) : i \in I, x \in X\}, \text{ and } \mu_D(x) = \inf\{\mu_{A_i}(x) : i \in I, x \in X\}$$

The symbol  $\tilde{0}$  will be used to denote the empty fuzzy set, so  $\mu_{\tilde{0}}(x) = 0$  for all  $x \in X$ , whereas  $\tilde{1}$  will be denoted for the set  $X$  where  $\mu_{\tilde{1}}(x) = 1$  for all  $x \in X$ .

For  $\lambda \in (0, 1]$ , a fuzzy point  $P_{x_0}^\lambda$  in  $X$  is a fuzzy set on  $X$  with support  $x_0$  and value  $\lambda$ , i.e:

$$P_{x_0}^\lambda = \begin{cases} \lambda, & \text{if } x = x_0, \text{ and} \\ 0, & \text{otherwise.} \end{cases}$$

so it can be written as  $P_{x_0}^\lambda = \{(x_0, \lambda)\}$ .

**Definition 2.3.** . A fuzzy topology on a set  $X$  is a family  $\tau$  of fuzzy sets in  $X$  which satisfies following conditions:

- $\tilde{1}, \tilde{0} \in \tau$  .
- If  $A, B \in \tau$  then  $A \cap B \in \tau$ .
- If  $A_i \in \tau$  then  $\cup_i A_i \in \tau$

the pair  $(X, \tau)$  is called a fuzzy topological space where its members are called open fuzzy sets. A fuzzy set is closed iff its complement  $A^c$  is open. As in general topology, the indiscrete fuzzy topology contains only  $\tilde{0}$ , and  $\tilde{1}$ , while the discrete fuzzy topology contains all fuzzy sets.

A fuzzy Filter on a fuzzy set  $X$  is a nonempty collection  $\mathfrak{F} \subset I^X$  of nonempty fuzzy sets on  $X$  which satisfies that  $F_1 \cap F_2 \in \mathfrak{F}$  for all  $F_1, F_2 \in \mathfrak{F}$  and if  $F_1 \subseteq F_2$  and  $F_1 \in \mathfrak{F}$  then  $F_2 \in \mathfrak{F}$ . The maximal, with respect to set inclusion, fuzzy filter on  $X$  is called a fuzzy ultrafilter on  $X$ . A collection  $\mathfrak{B}$  of subsets of  $\mathfrak{F}$  is a base for  $\mathfrak{F}$  iff for each  $F \in \mathfrak{F}$  there is some  $B \in \mathfrak{B}$  such that  $B \leq F$ .

A fuzzy topological space  $X$  is said to be a fuzzy door space if every fuzzy subset of  $X$  is either fuzzy open or fuzzy closed. Whereas, a topological space  $X$  is called a  $T_{1/2}$ -space if every one-point set is either open or closed. Clearly a door space is a  $T_{1/2}$ -space.

A point  $P_y^r$  is said to be an accumulation point of  $A$  iff for every open set  $U$  contains  $P_y^r$  satisfies  $U \cap A_p \neq \emptyset$ , where  $\mu_{A_p}(y) = 0$  and  $\mu_{A_p}(x) = \mu_A(x)$  otherwise.

### 3 Extremal Fuzzy Topology

**Definition 3.1.** Let  $r \in I$ , we define

- $P(X |_{x_0}^r)$  be the collection of all fuzzy subsets  $A \subseteq X$  where  $\mu_A(x_0) = r$ , so  $\mu_{X|_{x_0}^r}(x_0) = r$  and otherwise  $\mu_{X|_{x_0}^r}(x) = 1$ .
- $\mathfrak{S} \subseteq P(X |_{x_0}^r)$  be a filter on  $X$ . So for any  $F \subseteq \mathfrak{S}$ ,  $\mu_F(x_0) = r$ .

**Theorem 3.2.** Let  $X$  be a fuzzy space,  $P_{x_0}^r$  be a fuzzy point in  $X$  and  $\mathfrak{S}$  be a filter as above then:

$$\tau_{x_0} = P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$$

is a topology on  $X$ .

*Proof.* It is clear to prove  $\tau_{x_0}$  is a topology, where  $\tilde{0} \in P(X |_{x_0}^0)$  and  $\tilde{1} = X |_{x_0}^0 \cup (P_{x_0}^1 \cup F)$  for some  $F \in \mathfrak{S}$ .  $\square$

**Corollary 3.3.** The topology  $\tau_{x_0} = P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$  is a door space and hence a  $T_{1/2}$  space.

*Proof.* If  $A \subseteq X$  such that  $\mu_A(x_0) = 0$ , then  $A \in P(X |_{x_0}^0)$  is open. If  $\mu_A(x_0)$  and  $\mu_A(y)$  are not vanish for some  $y \neq x_0$ , then  $A$  has the form  $P_{x_0}^r \cup F$  for some  $F \in \mathfrak{S}$  and  $\mu_F(y) \neq 0$ , so  $A$  is also open. But in case  $\mu_A(x_0) \neq 0$  and  $\mu_A(y) = 0$  for all  $y \neq x_0$ , in this case  $A$  is closed. therefore  $\tau_{x_0}$  is a door space and hence a  $T_{1/2}$  space.  $\square$

**Lemma 3.4.** Let  $\mathfrak{B}$  be a filter base for a filter  $\mathfrak{S}$  and let  $x_0 \in X$ . Then  $\mathfrak{B}_{x_0} = \{P_x^i\} \cup \{P_{x_0}^r \cup B : B \in \mathfrak{B}\}$  be a base for the previous topology  $\tau_{x_0}$  for each  $x \in X$  and  $i, r \in (0, 1]$ .

*Proof.* If  $u$  be an open set in  $\tau_{x_0}$  which contains  $P_x^r$ . In case  $u \in P(X |_{x_0}^0)$ , the one point set  $P_x^r$  is contained in  $u$ . But if  $u$  has the form  $P_{x_0}^r \cup F$  for some  $F \in \mathfrak{S}$ , so there is  $B \in \mathfrak{B}$  such that  $B \subseteq F$ , and this yields to  $P_x^r \in P_{x_0}^r \cup B \subseteq P_{x_0}^r \cup F$ .  $\square$

**Corollary 3.5.** Let  $X$  be a non-empty fuzzy set and  $\tau_{x_0} = P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$  for some  $x_0 \in X$ . If  $\mathfrak{S}$  is a free filter on  $X |_{x_0}^0$ , then  $\tau_{x_0}$  is a Hausdorff space, furthermore, it has at most one accumulation point.

*Proof.* Let  $P_x^r \neq P_y^i$  for each  $x \neq y$  in  $X$  and  $r, i \in (0, 1]$ . If  $x \neq x_0$  and  $y \neq x_0$ , then  $P_x^r, P_y^i \in P(X |_{x_0}^0)$  are disjoint open set. But if  $x = x_0$  and  $y \neq x_0$ . Since  $\mathfrak{S}$  is a free filter, so  $\bigcap_{F \in \mathfrak{S}} F = \emptyset$ , then there is  $F \in \mathfrak{S}$  such that  $\mu_F(y) = 0$ . Let  $u = P_{x_0}^r \cup F$ . Therefore  $u$  and  $P_y^i$  are disjoint open sets contains, respectively,  $P_{x_0}^r$  and  $P_y^i$ . Thus  $\tau$  is a Hausdorff space.

By 3.3,  $\tau$  is a door space, and by [4], A Hausdorff door space has at most one accumulation point.  $\square$

**Corollary 3.6.** If  $\mathfrak{S} \subseteq P(X |_{x_0}^0)$  be an ultrafilter on a fuzzy set  $X$ , then the collection of the nonempty sets  $\mathfrak{S}_A = \{A \cap F : F \in \mathfrak{S}\} \subseteq P(A |_{x_0}^0)$  be an ultrafilter on  $A \subseteq X$ .

*Proof.* It is easy to prove that.  $\square$

In case  $\mathfrak{S}$  is an ultrafilter, we can deduce the following result which is the main aim of this paper:

**Theorem 3.7.** Let  $X$  be a fuzzy space and  $\mathfrak{S}$  be an ultrafilter, then any extremal topology on  $X$  has the following form:

$$\tau_{x_0} = P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$$

*Proof.* suppose that  $\tau_{x_0}$  be an extremal topology on a fuzzy set  $X$ , so there exists a fuzzy point, lets call it  $P_{x_0}^r$ , which is not in  $\tau_{x_0}$ , for  $r \in (0, 1]$ . Let  $\mathfrak{B} = \{B \in P(X |_{x_0}^0) : P_{x_0}^r \cup B \in \tau_{x_0}\}$  be a filter base for some a filter  $\wp$  on  $X |_{x_0}^0$ . Let  $\mathfrak{S}$  be an ultrafilter contains  $\wp$ .

Assume that  $\eta$  be a topology on  $X$  which is generated by  $\tau_{x_0} \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$ . Since  $P_{x_0}^r$  is not belong to  $\tau_{x_0}$  and  $F \neq \emptyset$  for all  $F \in \mathfrak{S}$ , so  $P_{x_0}^r$  also is not belong to  $\eta$ . This shows that  $\eta$  is non-discrete, and since  $\tau_{x_0}$  is extremal, this leads to  $\{P_{x_0}^r \cup F : F \in \mathfrak{S}\} \subset \tau_{x_0}$ . Also we have  $P_y^r \in \tau_{x_0}$  for all  $y \neq x_0$  because  $\tau_{x_0}$  is extremal, therefore  $P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\} \subseteq \tau_{x_0}$ .

On the other hand, let  $u \in \tau_{x_0}$  such that  $u$  is not in  $P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$ , then  $u = P_{x_0}^r \cup A$  where  $r \in (0, 1]$  such that  $A \cap F = \emptyset$  for all  $F \in \mathfrak{S}$ .

Since  $P_{x_0}^r \cup F \in \tau_{x_0}$ , then  $P_{x_0}^r = (P_{x_0}^r \cup A) \cap (P_{x_0}^r \cup F) \in \tau_{x_0}$ , this contradicts the extremality of  $\tau_{x_0}$ , therefore  $u \in P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$ . Thus  $\tau_{x_0} = P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$ .

Conversely, assume that  $\tau_{x_0} = P(X |_{x_0}^0) \cup \{P_{x_0}^r \cup F : F \in \mathfrak{S}\}$  and assume that  $\tau_{x_0} \subsetneq \tau$  for some a non-discrete topology  $\tau$  on  $X$ . So there exists  $u \in \tau$  which is not in  $\tau_{x_0}$ , then  $u$  has the form  $P_{x_0}^r \cup A$  where  $r \in (0, 1]$  and  $A \cap F = \emptyset$  for all  $F \in \mathfrak{S}$ . Similar to the previous discussing we get a contradiction. therefore  $\tau = \tau_{x_0}$ , and this shows that  $\tau_{x_0}$  is an extremal topology. □

**Corollary 3.8.** *Let  $\emptyset \neq A \subseteq X$ . The subspace  $\tau_A$  of an extremal topology  $\tau_{x_0}$  on  $X$  is an extremal topology on  $A$ .*

*Proof.* For a nonempty fuzzy subset  $A$  of  $X$ , we have  $\tau_A = \{A \cap u : u \in \tau_{x_0}\} = P(A |_{x_0}^0) \cup \{A \cap (P_{x_0}^r \cup F) : F \in \mathfrak{S}\}$  Since  $A \cap (P_{x_0}^r \cup F) = P_{x_0}^i \cup (A \cap F)$ , where  $i = \min\{r, \mu_A(x_0)\} \in (0, 1]$ . By 3.6,  $A \cap F = F_A \in \mathfrak{S}_A$ . Then  $\tau_A$  has the form  $P(A |_{x_0}^0) \cup \{P_{x_0}^i \cup F_A : F_A \in \mathfrak{S}_A\}$  which is an extremal topology on  $A$ . □

## References

- [1] Papazyan, T. Extremal topologies on a semigroup, Topology and its applications 39 (1991) 229-243
- [2] Mera, K. M. and Sola, M. A. (2005) Extremal Topology, Damascus University Journal for basic science, vol.21, No1.
- [3] Sola, M. A. A note on extremal topologies, unpublished paper.
- [4] Kelley, John L. General Topology. Springer. ISBN 3540901256. (1991).
- [5] Willard, S. (1970). General Topology, Addison-Wesley Publishing Company, INC
- [6] Zadeh, L. A. (1965), Fuzzy Sets, Information and Control, 8, pp. 338-353.
- [7] Chang, C. L. (1968), Fuzzy topological spaces, J. Math. Anal. Appl., 24, pp. 182-190.
- [8] M. A. Vicente AND M. S. Aranguren, "Fuzzy Filters", journal of Mathematical analysis and applications 129, 56G-568 (1988)
- [9] Zadeh, L. A., A Fuzzy-Set-Theoretic Interpretation of Linguistic Hedges, ERL Memorandum M335, University of California, Berkeley, April (1972). (To appear in Information Sciences.)
- [10] Wong, C. K., Fuzzy topology: Product and quotient theorems, J. Math. Anal. Appl. 45 (1974), 512-521.
- [11] C. De Mitri and E. Pascali, On Sequential Compactness and Semicompactness in Fuzzy Topology, journal of Mathematical analysis and applications 93. 324-327 (1983)