

A Study on Sum Formulas of Generalized Pentanacci Sequence: Closed Forms of the Sum Formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$

Yüksel Soykan

Department of Mathematics, Art and Science Faculty,
Zonguldak Bülent Ecevit University, 67100, Zonguldak, Turkey.
Email: yuksel_soykan@hotmail.com

Received: April 30, 2021; Accepted: May 20, 2021; Published: May 31, 2021

Copyright © 2021 by author(s) and Scitech Research Organisation(SRO).
This work is licensed under the Creative Commons Attribution International License (CC BY).
<http://creativecommons.org/licenses/by/4.0/>

Abstract

In this paper, closed forms of the sum formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$ for generalized Pentanacci numbers are presented. As special cases, we give summation formulas of Pentanacci, Pentanacci-Lucas, and other fifth-order recurrence sequences.

2020 Mathematics Subject Classification.

11B37, 11B39, 11B83.

Keywords

Pentanacci numbers, Pentanacci-Lucas numbers, sum formulas, summing formulas.

1. Introduction

The generalized Pentanacci sequence $\{W_n(W_0, W_1, W_2, W_3, W_4; r, s, t, u, v)\}_{n \geq 0}$ (or shortly $\{W_n\}_{n \geq 0}$) is defined as follows:

$$\begin{aligned} W_n &= rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4} + vW_{n-5}, \\ W_0 &= c_0, W_1 = c_1, W_2 = c_2, W_3 = c_3, W_4 = c_4, n \geq 5 \end{aligned} \quad (1.1)$$

where W_0, W_1, W_2, W_3, W_4 are arbitrary real or complex numbers and r, s, t, u, v are real numbers. The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = -\frac{u}{v}W_{-n+1} - \frac{t}{v}W_{-n+2} - \frac{s}{v}W_{-n+3} - \frac{r}{v}W_{-n+4} + \frac{1}{v}W_{-n+5}$$

for $n = 1, 2, 3, \dots$ when $v \neq 0$. Therefore, recurrence (1.1) holds for all integer n . Pentanacci sequence has been studied by many authors, see for example [8], [9], [11], [26].

Cite this paper: Soykan, Y. (2021). A Study on Sum Formulas of Generalized Pentanacci Sequence: Closed Forms of the Sum Formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$. *Journal of Progressive Research in Mathematics*, 18(2), 20-38. Retrieved from <http://scitecresearch.com/journals/index.php/jprm/article/view/2050>

Table 1 A few special case of generalized Pentanacci sequences.

No	Sequences (Numbers)	Notation	References
1	Generalized Pentanacci	$\{V_n\} = \{W_n(W_0, W_1, W_2, W_3, W_4; 1, 1, 1, 1, 1)\}$	[26]
2	Generalized Fifth order Pell	$\{V_n\} = \{W_n(W_0, W_1, W_2, W_3, W_4; 2, 1, 1, 1, 1)\}$	[27]
3	Generalized Fifth order Jacobsthal	$\{V_n\} = \{W_n(W_0, W_1, W_2, W_3, W_4; 1, 1, 1, 1, 2)\}$	[28]
4	Generalized 5-primes	$\{V_n\} = \{W_n(W_0, W_1, W_2, W_3, W_4; 2, 3, 5, 7, 11)\}$	[29]

For some specific values of W_0, W_1, W_2, W_3, W_4 and r, s, t, u, v it is worth presenting these special Pentanacci numbers in a table as a specific name. In literature, for example, the following names and notations (see Table 2) are used for the special cases of r, s, t, u, v and initial values.

Table 2 A few members of generalized Pentanacci sequences.

Sequences (Numbers)	Notation	OEIS [12]	Ref
Pentanacci	$\{P_n\} = \{W_n(0, 1, 1, 2, 4; 1, 1, 1, 1, 1)\}$	A001591	[26]
Pentanacci-Lucas	$\{Q_n\} = \{W_n(5, 1, 3, 7, 15; 1, 1, 1, 1, 1)\}$	A074048	[26]
fifth order Pell	$\{P_n^{(5)}\} = \{W_n(0, 1, 2, 5, 13; 2, 1, 1, 1, 1)\}$	A141448	[27]
fifth order Pell-Lucas	$\{Q_n^{(5)}\} = \{W_n(5, 2, 6, 17, 46; 2, 1, 1, 1, 1)\}$		[27]
modified fifth-order Pell	$\{E_n^{(5)}\} = \{W_n(0, 1, 1, 3, 8; 2, 1, 1, 1, 1)\}$		[27]
fifth order Jacobsthal	$\{J_n^{(5)}\} = \{W_n(0, 1, 1, 1, 1; 1, 1, 1, 1, 2)\}$	A226310	[28,2]
fifth order Jacobsthal-Lucas	$\{j_n^{(5)}\} = \{W_n(2, 1, 5, 10, 20; 1, 1, 1, 1, 2)\}$	A226311	[28,2]
modified fifth order Jacobsthal	$\{K_n^{(5)}\} = \{W_n(3, 1, 3, 10, 20; 1, 1, 1, 1, 2)\}$		[28]
fifth-order Jacobsthal Perrin	$\{Q_n^{(5)}\} = \{W_n(3, 0, 2, 8, 16; 1, 1, 1, 1, 2)\}$		[28]
adjusted fifth-order Jacobsthal	$\{S_n^{(5)}\} = \{W_n(0, 1, 1, 2, 4; 1, 1, 1, 1, 2)\}$		[28]
modified fifth-order Jacobsthal-Lucas	$\{R_n^{(5)}\} = \{W_n(5, 1, 3, 7, 15; 1, 1, 1, 1, 2)\}$		[28]
5-primes	$\{G_n\} = \{W_n(0, 0, 0, 1, 2; 2, 3, 5, 7, 11)\}$		[29]
Lucas 5-primes	$\{H_n\} = \{W_n(5, 2, 10, 41, 150; 2, 3, 5, 7, 11)\}$		[29]
modified 5-primes	$\{E_n\} = \{W_n(0, 0, 0, 1, 1; 2, 3, 5, 7, 11)\}$		[29]

For easy writing, from now on, we drop the superscripts from the sequences, for example we write P_n for $P_n^{(5)}$.

We present some works on summing formulas of the numbers in the following Table 3.

Table 3. A few special study of sum formulas.

Name of sequence	Papers which deal with summing formulas
Pell and Pell-Lucas	[1,4,30],[6,7]
Generalized Fibonacci	[5,13,14,15,16,17,19]
Generalized Tribonacci	[3,10,18]
Generalized Tetranacci	[20,25,31]
Generalized Pentanacci	[21,22]
Generalized Hexanacci	[23,24]

In this work, we investigate linear summation formulas of generalized Pentanacci numbers.

2 Sum Formulas of Generalized Pentanacci Numbers with Positive Subscripts

The following Theorem presents some linear summing formulas of generalized Pentanacci numbers with positive subscripts.

Theorem 2.1. *Let x be a real (or complex) number. For $n \geq 0$ we have the following formulas:*

(a) *If $rx + sx^2 + tx^3 + ux^4 + vx^5 - 1 \neq 0$ then*

$$\sum_{k=0}^n x^k W_k = \frac{\Theta_1(x)}{rx + sx^2 + tx^3 + ux^4 + vx^5 - 1}$$

where

$$\Theta_1(x) = x^{n+4}W_{n+4} - (rx - 1)x^{n+3}W_{n+3} - (sx^2 + rx - 1)x^{n+2}W_{n+2} - (sx^2 + tx^3 + rx - 1)x^{n+1}W_{n+1} + vx^{n+5}W_n - x^4W_4 + x^3(rx - 1)W_3 + x^2(sx^2 + rx - 1)W_2 + x(sx^2 + tx^3 + rx - 1)W_1 + (sx^2 + tx^3 + ux^4 + rx - 1)W_0.$$

(b) *If $r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + v^2x^5 + 2sx + 2rtx^2 + 2rvx^3 - 2sux^3 + 2tvx^4 - 1 \neq 0$ then*

$$\sum_{k=0}^n x^k W_{2k} = \frac{\Theta_2(x)}{r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + v^2x^5 + 2sx + 2rtx^2 + 2rvx^3 - 2sux^3 + 2tvx^4 - 1}$$

where

$$\Theta_2(x) = -(ux^2 + sx - 1)x^{n+1}W_{2n+2} + (t + rs + vx + rux)x^{n+2}W_{2n+1} + (u + t^2x - u^2x^2 + v^2x^3 + rt + 2tvx^2 + rvx - sux)x^{n+2}W_{2n} + (v + ru - svx + tux)x^{n+2}W_{2n-1} + v(r + vx^2 + tx)x^{n+2}W_{2n-2} + x^2(ux^2 + sx - 1)W_4 - x^3(t + rs + vx + rux)W_3 + x(r^2x + ux^2 - s^2x^2 + 2sx + rtx^2 + rvx^3 - sux^3 - 1)W_2 - x^3(v + ru - svx + tux)W_1 + (r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + 2sx + 2rtx^2 + rvx^3 - 2sux^3 + tvx^4 - 1)W_0.$$

(c) *If $r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + v^2x^5 + 2sx + 2rtx^2 + 2rvx^3 - 2sux^3 + 2tvx^4 - 1 = 0$ then*

$$\sum_{k=0}^n x^k W_{2k+1} = \frac{\Theta_3(x)}{r^2x + 2ux^2 - s^2x^2 + t^2x^3 - u^2x^4 + v^2x^5 + 2sx + 2rtx^2 + 2rvx^3 - 2sux^3 + 2tvx^4 - 1}$$

where

$$\Theta_3(x) = (r + vx^2 + tx)x^{n+1}W_{2n+2} + (s - s^2x + t^2x^2 - u^2x^3 + v^2x^4 + ux + rvx^2 - 2sux^2 + 2tvx^3 + rtx)x^{n+1}W_{2n+1} + (t + vx - svx^2 + rux - stx)x^{n+1}W_{2n} + (u - u^2x^2 + v^2x^3 + tvx^2 + rvx - sux)x^{n+1}W_{2n-1} - v(ux^2 + sx - 1)x^{n+1}W_{2n-2} - x^2(r + vx^2 + tx)W_4 + x(r^2x + ux^2 + sx + rtx^2 + rvx^3 - 1)W_3 - x^2(t + vx - svx^2 + rux - stx)W_2 + (r^2x + ux^2 - s^2x^2 + t^2x^3 + 2sx + 2rtx^2 + rvx^3 - sux^3 + tvx^4 - 1)W_1 + vx^2(ux^2 + sx - 1)W_0$$

Proof.

(a) Using the recurrence relation

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4} + vW_{n-5}$$

i.e.

$$vW_{n-5} = W_n - rW_{n-1} - sW_{n-2} - tW_{n-3} - uW_{n-4}$$

we obtain

$$\begin{aligned} vx^0W_0 &= x^0W_5 - rx^0W_4 - sx^0W_3 - tx^0W_2 - ux^0W_1 \\ vx^1W_1 &= x^1W_6 - rx^1W_5 - sx^1W_4 - tx^1W_3 - ux^1W_2 \\ vx^2W_2 &= x^2W_7 - rx^2W_6 - sx^2W_5 - tx^2W_4 - ux^2W_3 \\ vx^3W_3 &= x^3W_8 - rx^3W_7 - sx^3W_6 - tx^3W_5 - ux^3W_4 \\ &\vdots \\ vx^{n-4}W_{n-4} &= x^{n-4}W_{n+1} - rx^{n-4}W_n - sx^{n-4}W_{n-1} - tx^{n-4}W_{n-2} - ux^{n-4}W_{n-3} \\ vx^{n-3}W_{n-3} &= x^{n-3}W_{n+2} - rx^{n-3}W_{n+1} - sx^{n-3}W_n - tx^{n-3}W_{n-1} - ux^{n-3}W_{n-2} \\ vx^{n-2}W_{n-2} &= x^{n-2}W_{n+3} - rx^{n-2}W_{n+2} - sx^{n-2}W_{n+1} - tx^{n-2}W_n - ux^{n-2}W_{n-1} \\ vx^{n-1}W_{n-1} &= x^{n-1}W_{n+4} - rx^{n-1}W_{n+3} - sx^{n-1}W_{n+2} - tx^{n-1}W_{n+1} - ux^{n-1}W_n \\ vx^nW_n &= x^nW_{n+5} - rx^nW_{n+4} - sx^nW_{n+3} - tx^nW_{n+2} - ux^nW_{n+1} \end{aligned}$$

If we add the equations side by side (and using $W_{n+5} = rW_{n+4} + sW_{n+3} + tW_{n+2} + uW_{n+1} + vW_n$), we get (a)

(b) and (c) Using the recurrence relation

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4} + vW_{n-5}$$

i.e.

$$rW_{n-1} = W_n - sW_{n-2} - tW_{n-3} - uW_{n-4} - vW_{n-5}$$

we obtain

$$\begin{aligned} rx^1W_3 &= x^1W_4 - sx^1W_2 - tx^1W_1 - ux^1W_0 - vx^1W_{-1} \\ rx^2W_5 &= x^2W_6 - sx^2W_4 - tx^2W_3 - ux^2W_2 - vx^2W_1 \\ rx^3W_7 &= x^3W_8 - sx^3W_6 - tx^3W_5 - ux^3W_4 - vx^3W_3 \\ rx^4W_9 &= x^4W_{10} - sx^4W_8 - tx^4W_7 - ux^4W_6 - vx^4W_5 \\ &\vdots \\ rx^{n-1}W_{2n-1} &= x^{n-1}W_{2n} - sx^{n-1}W_{2n-2} - tx^{n-1}W_{2n-3} - ux^{n-1}W_{2n-4} - vx^{n-1}W_{2n-5} \\ rx^nW_{2n+1} &= x^nW_{2n+2} - sx^nW_{2n} - tx^nW_{2n-1} - ux^nW_{2n-2} - vx^nW_{2n-3} \\ rx^{n+1}W_{2n+3} &= x^{n+1}W_{2n+4} - sx^{n+1}W_{2n+2} - tx^{n+1}W_{2n+1} - ux^{n+1}W_{2n} - vx^{n+1}W_{2n-1} \\ rx^{n+2}W_{2n+5} &= x^{n+2}W_{2n+6} - sx^{n+2}W_{2n+4} - tx^{n+2}W_{2n+3} - ux^{n+2}W_{2n+2} - vx^{n+2}W_{2n+1} \end{aligned}$$

Now, if we add the above equations side by side, we get

$$\begin{aligned}
r(-x^0W_1 + \sum_{k=0}^n x^k W_{2k+1}) &= (x^n W_{2n+2} - x^0 W_2 - x^{-1} W_0 + \sum_{k=0}^n x^{k-1} W_{2k}) \\
&\quad -s(-x^0 W_0 + \sum_{k=0}^n x^k W_{2k}) - t(-x^{n+1} W_{2n+1} + \sum_{k=0}^n x^{k+1} W_{2k+1}) \\
&\quad -u(-x^{n+1} W_{2n} + \sum_{k=0}^n x^{k+1} W_{2k}) \\
&\quad -v(-x^{n+2} W_{2n+1} - x^{n+1} W_{2n-1} + x^1 W_{-1} + \sum_{k=0}^n x^{k+2} W_{2k+1})
\end{aligned}$$

Since

$$W_{-1} = -\frac{u}{v}W_0 - \frac{t}{v}W_1 - \frac{s}{v}W_2 - \frac{r}{v}W_3 + \frac{1}{v}W_4$$

we obtain

$$\begin{aligned}
r(-x^0W_1 + \sum_{k=0}^n x^k W_{2k+1}) &= (x^n W_{2n+2} - x^0 W_2 - x^{-1} W_0 + x^{-1} \sum_{k=0}^n x^k W_{2k}) \quad (2.1) \\
&\quad -s(-x^0 W_0 + \sum_{k=0}^n x^k W_{2k}) - t(-x^{n+1} W_{2n+1} + x^1 \sum_{k=0}^n x^k W_{2k+1}) \\
&\quad -u(-x^{n+1} W_{2n} + x^1 \sum_{k=0}^n x^k W_{2k}) - v(-x^{n+2} W_{2n+1} - x^{n+1} W_{2n-1} \\
&\quad + x^1(-\frac{u}{v}W_0 - \frac{t}{v}W_1 - \frac{s}{v}W_2 - \frac{r}{v}W_3 + \frac{1}{v}W_4) + x^2 \sum_{k=0}^n x^k W_{2k+1}).
\end{aligned}$$

Similarly, using the recurrence relation

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4} + vW_{n-5}$$

i.e.

$$rW_{n-1} = W_n - sW_{n-2} - tW_{n-3} - uW_{n-4} - vW_{n-5}$$

we write the following obvious equations;

$$\begin{aligned}
rx^1W_2 &= x^1W_3 - sx^1W_1 - tx^1W_0 - ux^1W_{-1} - vx^1W_{-2} \\
rx^2W_4 &= x^2W_5 - sx^2W_3 - tx^2W_2 - ux^2W_1 - vx^2W_0 \\
rx^3W_6 &= x^3W_7 - sx^3W_5 - tx^3W_4 - ux^3W_3 - vx^3W_2 \\
rx^4W_8 &= x^4W_9 - sx^4W_7 - tx^4W_6 - ux^4W_5 - vx^4W_4 \\
&\quad \vdots \\
rx^{n-1}W_{2n-2} &= x^{n-1}W_{2n-1} - sx^{n-1}W_{2n-3} - tx^{n-1}W_{2n-4} - ux^{n-1}W_{2n-5} - vx^{n-1}W_{2n-6} \\
rx^nW_{2n} &= x^nW_{2n+1} - sx^nW_{2n-1} - tx^nW_{2n-2} - ux^nW_{2n-3} - vx^nW_{2n-4} \\
rx^{n+1}W_{2n+2} &= x^{n+1}W_{2n+3} - sx^{n+1}W_{2n+1} - tx^{n+1}W_{2n} - ux^{n+1}W_{2n-1} - vx^{n+1}W_{2n-2} \\
rx^{n+2}W_{2n+4} &= x^{n+2}W_{2n+5} - sx^{n+2}W_{2n+3} - tx^{n+2}W_{2n+2} - ux^{n+2}W_{2n+1} - vx^{n+2}W_{2n}
\end{aligned}$$

Now, if we add the above equations side by side, we obtain

$$\begin{aligned}
r(-x^0W_0 + \sum_{k=0}^n x^k W_{2k}) &= (-x^0W_1 + \sum_{k=0}^n x^k W_{2k+1}) - s(-x^{n+1}W_{2n+1} + \sum_{k=0}^n x^{k+1}W_{2k+1}) \\
&\quad - t(-x^{n+1}W_{2n} + \sum_{k=0}^n x^{k+1}W_{2k}) - u(-x^{n+2}W_{2n+1} - x^{n+1}W_{2n-1} \\
&\quad + x^1W_{-1} + \sum_{k=0}^n x^{k+2}W_{2k+1}) \\
&\quad - v(-x^{n+2}W_{2n} - x^{n+1}W_{2n-2} + x^1W_{-2} + \sum_{k=0}^n x^{k+2}W_{2k})
\end{aligned}$$

Since

$$\begin{aligned}
W_{-1} &= -\frac{u}{v}W_0 - \frac{t}{v}W_1 - \frac{s}{v}W_2 - \frac{r}{v}W_3 + \frac{1}{v}W_4 \\
W_{-2} &= -\frac{u}{v}(-\frac{u}{v}W_0 - \frac{t}{v}W_1 - \frac{s}{v}W_2 - \frac{r}{v}W_3 + \frac{1}{v}W_4) - \frac{t}{v}W_0 - \frac{s}{v}W_1 - \frac{r}{v}W_2 + \frac{1}{v}W_3
\end{aligned}$$

we have

$$\begin{aligned}
r(-x^0W_0 + \sum_{k=0}^n x^k W_{2k}) &= (-x^0W_1 + \sum_{k=0}^n x^k W_{2k+1}) - s(-x^{n+1}W_{2n+1} + x^1 \sum_{k=0}^n x^k W_{2k+1}) \quad (2.2) \\
&\quad - t(-x^{n+1}W_{2n} + x^1 \sum_{k=0}^n x^k W_{2k}) - u(-x^{n+2}W_{2n+1} - x^{n+1}W_{2n-1} \\
&\quad + x^1(-\frac{u}{v}W_0 - \frac{t}{v}W_1 - \frac{s}{v}W_2 - \frac{r}{v}W_3 + \frac{1}{v}W_4) + x^2 \sum_{k=0}^n x^k W_{2k+1}) \\
&\quad - v(-x^{n+2}W_{2n} - x^{n+1}W_{2n-2} + x^1(-\frac{u}{v}(-\frac{u}{v}W_0 - \frac{t}{v}W_1 - \frac{s}{v}W_2 \\
&\quad - \frac{r}{v}W_3 + \frac{1}{v}W_4) - \frac{t}{v}W_0 - \frac{s}{v}W_1 - \frac{r}{v}W_2 + \frac{1}{v}W_3) + x^2 \sum_{k=0}^n x^k W_{2k}).
\end{aligned}$$

Then, solving the system (2.1)-(2.2), the required result of (b) and (c) follow. \square

3 Special Cases

In this section, for the special cases of x , we present the closed form solutions (identities) of the sums $\sum_{k=0}^n x^k W_k$, $\sum_{k=0}^n x^k W_{2k}$ and $\sum_{k=0}^n x^k W_{2k+1}$ for the specific case of sequence $\{W_n\}$.

3.1 The case $x = 1$

In this subsection we consider the special case $x = 1$.

The case $x = 1$ of Theorem 2.1 is given in Soykan [21].

3.2 The case $x = -1$

In this subsection we consider the special case $x = -1$ and we present the closed form solutions (identities) of the sums $\sum_{k=0}^n (-1)^k W_k$, $\sum_{k=0}^n (-1)^k W_{2k}$ and $\sum_{k=0}^n (-1)^k W_{2k+1}$ for the specific case of the sequence $\{W_n\}$.

Taking $r = s = t = u = v = 1$ in Theorem 2.1 (a), (b) and (c), we obtain the following Proposition.

Proposition 3.1. *If $r = s = t = u = v = 1$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{2}((-1)^n (-W_{n+4} + 2W_{n+3} - W_{n+2} + 2W_{n+1} + W_n) + W_4 - 2W_3 + W_2 - 2W_1 + W_0)$.
- (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{2}((-1)^n (W_{2n+2} - W_{2n} - 2W_{2n-1} - W_{2n-2}) + W_4 - 3W_2 - 2W_1 + W_0)$.
- (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{2}((-1)^n (W_{2n+2} - W_{2n} + W_{2n-2}) + W_4 - 2W_3 - W_2 + 2W_1 + W_0)$.

From the above Proposition, we have the following Corollary which gives sum formulas of Pentanacci numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2, P_4 = 4$).

Corollary 3.2. *For $n \geq 0$, Pentanacci numbers have the following properties.*

- (a) $\sum_{k=0}^n (-1)^k P_k = \frac{1}{2}((-1)^n (-P_{n+4} + 2P_{n+3} - P_{n+2} + 2P_{n+1} + P_n) - 1)$.
- (b) $\sum_{k=0}^n (-1)^k P_{2k} = \frac{1}{2}((-1)^n (P_{2n+2} - P_{2n} - 2P_{2n-1} - P_{2n-2}) - 1)$.
- (c) $\sum_{k=0}^n (-1)^k P_{2k+1} = \frac{1}{2}((-1)^n (P_{2n+2} - P_{2n} + P_{2n-2}) + 1)$.

Taking $W_n = Q_n$ with $Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7, Q_4 = 15$ in the above Proposition, we have the following Corollary which presents sum formulas of Pentanacci-Lucas numbers.

Corollary 3.3. *For $n \geq 0$, Pentanacci-Lucas numbers have the following properties.*

- (a) $\sum_{k=0}^n (-1)^k Q_k = \frac{1}{2}((-1)^n (-Q_{n+4} + 2Q_{n+3} - Q_{n+2} + 2Q_{n+1} + Q_n) + 7)$.
- (b) $\sum_{k=0}^n (-1)^k Q_{2k} = \frac{1}{2}((-1)^n (Q_{2n+2} - Q_{2n} - 2Q_{2n-1} - Q_{2n-2}) + 9)$.
- (c) $\sum_{k=0}^n (-1)^k Q_{2k+1} = \frac{1}{2}((-1)^n (Q_{2n+2} - Q_{2n} + Q_{2n-2}) + 5)$.

Taking $r = 2, s = t = u = v = 1$ in Theorem 2.1 (a), (b) and (c), we obtain the following Proposition.

Proposition 3.4. *If $r = 2, s = t = u = v = 1$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{3}((-1)^n (-W_{n+4} + 3W_{n+3} - 2W_{n+2} + 3W_{n+1} + W_n) + W_4 - 3W_3 + 2W_2 - 3W_1 + 2W_0)$.
- (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{5}((-1)^n (W_{2n+2} - W_{2n} - 3W_{2n-1} - 2W_{2n-2}) + W_4 - 6W_2 - 3W_1 + 3W_0)$.
- (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{5}((-1)^n (2W_{2n+2} - 2W_{2n} - W_{2n-1} + W_{2n-2}) + 2W_4 - 5W_3 - 2W_2 + 4W_1 + W_0)$.

From the last Proposition, we have the following Corollary which gives linear sum formulas of fifth-order Pell numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5, P_4 = 13$).

Corollary 3.5. *For $n \geq 0$, fifth-order Pell numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k P_k = \frac{1}{3}((-1)^n (-P_{n+4} + 3P_{n+3} - 2P_{n+2} + 3P_{n+1} + P_n) + P_4 - 3P_3 + 2P_2 - 3P_1 + 2P_0)$.
- (b) $\sum_{k=0}^n (-1)^k P_{2k} = \frac{1}{5}((-1)^n (P_{2n+2} - P_{2n} - 3P_{2n-1} - 2P_{2n-2}) + P_4 - 6P_2 - 3P_1 + 3P_0)$.
- (c) $\sum_{k=0}^n (-1)^k P_{2k+1} = \frac{1}{5}((-1)^n (2P_{2n+2} - 2P_{2n} - P_{2n-1} + P_{2n-2}) + 2P_4 - 5P_3 - 2P_2 + 4P_1 + P_0)$.

Taking $W_n = Q_n$ with $Q_0 = 5, Q_1 = 2, Q_2 = 6, Q_3 = 17, Q_4 = 46$ in the last Proposition, we have the following Corollary which presents linear sum formulas of fifth-order Pell-Lucas numbers.

Corollary 3.6. *For $n \geq 0$, fifth-order Pell-Lucas numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k Q_k = \frac{1}{3}((-1)^n (-Q_{n+4} + 3Q_{n+3} - 2Q_{n+2} + 3Q_{n+1} + Q_n) + Q_4 - 3Q_3 + 2Q_2 - 3Q_1 + 2Q_0)$.
- (b) $\sum_{k=0}^n (-1)^k Q_{2k} = \frac{1}{5}((-1)^n (Q_{2n+2} - Q_{2n} - 3Q_{2n-1} - 2Q_{2n-2}) + Q_4 - 6Q_2 - 3Q_1 + 3Q_0)$.
- (c) $\sum_{k=0}^n (-1)^k Q_{2k+1} = \frac{1}{5}((-1)^n (2Q_{2n+2} - 2Q_{2n} - Q_{2n-1} + Q_{2n-2}) + 2Q_4 - 5Q_3 - 2Q_2 + 4Q_1 + Q_0)$.

Taking $r = 1, s = 1, t = 1, u = 1, v = 2$ in Theorem 2.1 (a), (b) and (c), we obtain the following Proposition.

Proposition 3.7. *If $r = s = t = 1, u = 1, v = 2$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{3}((-1)^n (-W_{n+4} + 2W_{n+3} - W_{n+2} + 2W_{n+1} + 2W_n) + W_4 - 2W_3 + W_2 - 2W_1 + W_0)$.
- (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{5}((-1)^n (W_{2n+2} + W_{2n+1} + W_{2n} - 4W_{2n-1} - 4W_{2n-2}) + W_4 + W_3 - 4W_2 - 4W_1 + W_0)$.
- (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{5}((-1)^n (2W_{2n+2} + 2W_{2n+1} - 3W_{2n} - 3W_{2n-1} + 2W_{2n-2}) + 2W_4 - 3W_3 - 3W_2 + 2W_1 + 2W_0)$.

Taking $W_n = J_n$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1, J_4 = 1$ in the last Proposition, we have the following Corollary which presents linear sum formulas of fifth-order Jacobsthal numbers.

Corollary 3.8. *For $n \geq 0$, fifth order Jacobsthal numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k J_k = \frac{1}{3}((-1)^n (-J_{n+4} + 2J_{n+3} - J_{n+2} + 2J_{n+1} + 2J_n) - 2)$.
- (b) $\sum_{k=0}^n (-1)^k J_{2k} = \frac{1}{5}((-1)^n (J_{2n+2} + J_{2n+1} + J_{2n} - 4J_{2n-1} - 4J_{2n-2}) - 6)$.
- (c) $\sum_{k=0}^n (-1)^k J_{2k+1} = \frac{1}{5}((-1)^n (2J_{2n+2} + 2J_{2n+1} - 3J_{2n} - 3J_{2n-1} + 2J_{2n-2}) - 2)$.

From the last Proposition, we have the following Corollary which gives linear sum formulas of fifth order Jacobsthal-Lucas numbers (take $W_n = j_n$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10, j_4 = 20$).

Corollary 3.9. *For $n \geq 0$, fifth order Jacobsthal-Lucas numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k j_k = \frac{1}{3}((-1)^n (-j_{n+4} + 2j_{n+3} - j_{n+2} + 2j_{n+1} + 2j_n) + 5)$.
- (b) $\sum_{k=0}^n (-1)^k j_{2k} = \frac{1}{5}((-1)^n (j_{2n+2} + j_{2n+1} + j_{2n} - 4j_{2n-1} - 4j_{2n-2}) + 8)$.
- (c) $\sum_{k=0}^n (-1)^k j_{2k+1} = \frac{1}{5}((-1)^n (2j_{2n+2} + 2j_{2n+1} - 3j_{2n} - 3j_{2n-1} + 2j_{2n-2}) + 1)$.

Taking $W_n = K_n$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10, K_4 = 20$ in the last proposition, we have the following corollary which presents linear sum formula of modified fifth order Jacobsthal numbers.

Corollary 3.10. *For $n \geq 0$, modified fifth order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k K_k = \frac{1}{3}((-1)^n (-K_{n+4} + 2K_{n+3} - K_{n+2} + 2K_{n+1} + 2K_n) + 4)$.
- (b) $\sum_{k=0}^n (-1)^k K_{2k} = \frac{1}{5}((-1)^n (K_{2n+2} + K_{2n+1} + K_{2n} - 4K_{2n-1} - 4K_{2n-2}) + 17)$.
- (c) $\sum_{k=0}^n (-1)^k K_{2k+1} = \frac{1}{5}((-1)^n (2K_{2n+2} + 2K_{2n+1} - 3K_{2n} - 3K_{2n-1} + 2K_{2n-2}) + 9)$.

From the last proposition, we have the following corollary which gives linear sum formula of fifth-order Jacobsthal Perrin numbers (take $W_n = Q_n$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8, Q_4 = 16$).

Corollary 3.11. *For $n \geq 0$, fifth-order Jacobsthal Perrin numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k Q_k = \frac{1}{3}((-1)^n (-Q_{n+4} + 2Q_{n+3} - Q_{n+2} + 2Q_{n+1} + 2Q_n) + 5)$.
- (b) $\sum_{k=0}^n (-1)^k Q_{2k} = \frac{1}{5}((-1)^n (Q_{2n+2} + Q_{2n+1} + Q_{2n} - 4Q_{2n-1} - 4Q_{2n-2}) + 19)$.
- (c) $\sum_{k=0}^n (-1)^k Q_{2k+1} = \frac{1}{5}((-1)^n (2Q_{2n+2} + 2Q_{2n+1} - 3Q_{2n} - 3Q_{2n-1} + 2Q_{2n-2}) + 8)$.

Taking $W_n = S_n$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2, S_4 = 4$ in the proposition, we have the following corollary which presents linear sum formula of adjusted fifth-order Jacobsthal numbers.

Corollary 3.12. *For $n \geq 0$, adjusted fifth-order Jacobsthal numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k S_k = \frac{1}{3}((-1)^n (-S_{n+4} + 2S_{n+3} - S_{n+2} + 2S_{n+1} + 2S_n) - 1)$.
 (b) $\sum_{k=0}^n (-1)^k S_{2k} = \frac{1}{5}((-1)^n (S_{2n+2} + S_{2n+1} + S_{2n} - 4S_{2n-1} - 4S_{2n-2}) - 2)$.
 (c) $\sum_{k=0}^n (-1)^k S_{2k+1} = \frac{1}{5}((-1)^n (2S_{2n+2} + 2S_{2n+1} - 3S_{2n} - 3S_{2n-1} + 2S_{2n-2}) + 1)$.

From the last proposition, we have the following corollary which gives linear sum formula of modified fifth-order Jacobsthal-Lucas numbers (take $W_n = R_n$ with $R_0 = 5, R_1 = 1, R_2 = 3, R_3 = 7, R_4 = 15$).

Corollary 3.13. *For $n \geq 0$, modified fifth-order Jacobsthal-Lucas numbers have the following property:*

- (a) $\sum_{k=0}^n (-1)^k R_k = \frac{1}{3}((-1)^n (-R_{n+4} + 2R_{n+3} - R_{n+2} + 2R_{n+1} + 2R_n) + 7)$.
 (b) $\sum_{k=0}^n (-1)^k R_{2k} = \frac{1}{5}((-1)^n (R_{2n+2} + R_{2n+1} + R_{2n} - 4R_{2n-1} - 4R_{2n-2}) + 11)$.
 (c) $\sum_{k=0}^n (-1)^k R_{2k+1} = \frac{1}{5}((-1)^n (2R_{2n+2} + 2R_{2n+1} - 3R_{2n} - 3R_{2n-1} + 2R_{2n-2}) + 12)$.

Taking $r = 2, s = 3, t = 5, u = 7, v = 11$ in Theorem 2.1 (a), (b) and (c), we obtain the following proposition.

Proposition 3.14. *If $r = 2, s = 3, t = 5, u = 7, v = 11$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n (-1)^k W_k = \frac{1}{9}((-1)^n (-W_{n+4} + 3W_{n+3} + 5W_{n+1} + 11W_n) + W_4 - 3W_3 - 5W_1 - 2W_0)$.
 (b) $\sum_{k=0}^n (-1)^k W_{2k} = \frac{1}{73}((-1)^n (-3W_{2n+2} + 14W_{2n+1} + 69W_{2n} - 23W_{2n-1} - 88W_{2n-2}) - 3W_4 + 14W_3 - 4W_2 - 23W_1 - 15W_0)$.
 (c) $\sum_{k=0}^n (-1)^k W_{2k+1} = \frac{1}{73}((-1)^n (8W_{2n+2} + 60W_{2n+1} - 38W_{2n} - 109W_{2n-1} - 33W_{2n-2}) + 8W_4 - 13W_3 - 38W_2 - 36W_1 - 33W_0)$.

From the last proposition, we have the following corollary which gives linear sum formulas of 5-primes numbers (take $W_n = G_n$ with $G_0 = 0, G_1 = 0, G_2 = 0, G_3 = 1, G_4 = 2$).

Corollary 3.15. *For $n \geq 0$, 5-primes numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k G_k = \frac{1}{9}((-1)^n (-G_{n+4} + 3G_{n+3} + 5G_{n+1} + 11G_n) - 1)$.
 (b) $\sum_{k=0}^n (-1)^k G_{2k} = \frac{1}{73}((-1)^n (-3G_{2n+2} + 14G_{2n+1} + 69G_{2n} - 23G_{2n-1} - 88G_{2n-2}) + 8)$.
 (c) $\sum_{k=0}^n (-1)^k G_{2k+1} = \frac{1}{73}((-1)^n (8G_{2n+2} + 60G_{2n+1} - 38G_{2n} - 109G_{2n-1} - 33G_{2n-2}) + 3)$.

Taking $W_n = H_n$ with $H_0 = 5, H_1 = 2, H_2 = 10, H_3 = 41, H_4 = 150$ in the last proposition, we have the following corollary which presents linear sum formulas of Lucas 5-primes numbers.

Corollary 3.16. *For $n \geq 0$, Lucas 5-primes numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k H_k = \frac{1}{9}((-1)^n (-H_{n+4} + 3H_{n+3} + 5H_{n+1} + 11H_n) + 7)$.
 (b) $\sum_{k=0}^n (-1)^k H_{2k} = \frac{1}{73}((-1)^n (-3H_{2n+2} + 14H_{2n+1} + 69H_{2n} - 23H_{2n-1} - 88H_{2n-2}) - 37)$.
 (c) $\sum_{k=0}^n (-1)^k H_{2k+1} = \frac{1}{73}((-1)^n (8H_{2n+2} + 60H_{2n+1} - 38H_{2n} - 109H_{2n-1} - 33H_{2n-2}) + 50)$.

From the last proposition, we have the following corollary which gives linear sum formulas of modified 5-primes numbers (take $W_n = E_n$ with $E_0 = 0, E_1 = 0, E_2 = 0, E_3 = 1, E_4 = 1$).

Corollary 3.17. *For $n \geq 0$, modified 5-primes numbers have the following properties:*

- (a) $\sum_{k=0}^n (-1)^k E_k = \frac{1}{9}((-1)^n (-E_{n+4} + 3E_{n+3} + 5E_{n+1} + 11E_n) - 2)$.
 (b) $\sum_{k=0}^n (-1)^k E_{2k} = \frac{1}{73}((-1)^n (-3E_{2n+2} + 14E_{2n+1} + 69E_{2n} - 23E_{2n-1} - 88E_{2n-2}) + 11)$.
 (c) $\sum_{k=0}^n (-1)^k E_{2k+1} = \frac{1}{73}((-1)^n (8E_{2n+2} + 60E_{2n+1} - 38E_{2n} - 109E_{2n-1} - 33E_{2n-2}) - 5)$.

3.3 The case $x = i$

In this subsection we consider the special case $x = i$.

Taking $x = i, r = s = t = u = v = 1$ in Theorem 2.1 (a), (b) and (c), we obtain the following proposition.

Proposition 3.18. *If $r = s = t = u = v = 1$ then for $n \geq 0$ we have the following formulas:*

- (a) $\sum_{k=0}^n i^k W_k = \frac{1}{-1+i} (i^n (W_{n+4} - (1+i)W_{n+3} - (2-i)W_{n+2} + 2iW_{n+1} + iW_n) - W_4 + (1+i)W_3 + (2-i)W_2 - 2iW_1 - W_0).$
- (b) $\sum_{k=0}^n i^k W_{2k} = \frac{1}{-3+3i} (i^n ((1+2i)W_{2n+2} - (2+2i)W_{2n+1} - W_{2n} - 2W_{2n-1} - iW_{2n-2}) + (2-i)W_4 - (2-2i)W_3 - (3+2i)W_2 + 2iW_1 - (4-3i)W_0).$
- (c) $\sum_{k=0}^n i^k W_{2k+1} = \frac{1}{-3+3i} (i^n (-W_{2n+2} + 2iW_{2n+1} + (-1+2i)W_{2n} + (1+i)W_{2n-1} + (1+2i)W_{2n-2}) + iW_4 - (1+3i)W_3 + (2+i)W_2 - (2-2i)W_1 + (2-i)W_0).$

From the above Proposition, we have the following Corollary which gives linear sum formulas of Pentanacci numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2, P_4 = 4$).

Corollary 3.19. *For $n \geq 0$, Pentanacci numbers have the following properties.*

- (a) $\sum_{k=0}^n i^k P_k = \frac{1}{-1+i} (i^n (P_{n+4} - (1+i)P_{n+3} - (2-i)P_{n+2} + 2iP_{n+1} + iP_n) - i).$
- (b) $\sum_{k=0}^n i^k P_{2k} = \frac{1}{-3+3i} (i^n ((1+2i)P_{2n+2} - (2+2i)P_{2n+1} - P_{2n} - 2P_{2n-1} - iP_{2n-2}) + 1).$
- (c) $\sum_{k=0}^n i^k P_{2k+1} = \frac{1}{-3+3i} (i^n (-P_{2n+2} + 2iP_{2n+1} + (-1+2i)P_{2n} + (1+i)P_{2n-1} + (1+2i)P_{2n-2}) + (-2+i)).$

Taking $W_n = Q_n$ with $Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7, Q_4 = 15$ in the above Proposition, we have the following Corollary which presents linear sum formulas of Pentanacci-Lucas numbers.

Corollary 3.20. *For $n \geq 0$, Pentanacci-Lucas numbers have the following properties.*

- (a) $\sum_{k=0}^n i^k Q_k = \frac{1}{-1+i} (i^n (Q_{n+4} - (1+i)Q_{n+3} - (2-i)Q_{n+2} + 2iQ_{n+1} + iQ_n) + (-7+2i)).$
- (b) $\sum_{k=0}^n i^k Q_{2k} = \frac{1}{-3+3i} (i^n ((1+2i)Q_{2n+2} - (2+2i)Q_{2n+1} - Q_{2n} - 2Q_{2n-1} - iQ_{2n-2}) + (-13+10i)).$
- (c) $\sum_{k=0}^n i^k Q_{2k+1} = \frac{1}{-3+3i} (i^n (-Q_{2n+2} + 2iQ_{2n+1} + (-1+2i)Q_{2n} + (1+i)Q_{2n-1} + (1+2i)Q_{2n-2}) + (7-6i)).$

Corresponding sums of the other fifth order generalized Pentanacci numbers can be calculated similarly.

4 Sum Formulas of Generalized Pentanacci Numbers with Negative Subscripts

The following Theorem presents some linear summing formulas of generalized Pentanacci numbers with negative subscripts.

Theorem 4.1. *Let x be a real (or complex) number. For $n \geq 1$ we have the following formulas:*

- (a) *(Sum of the generalized Pentanacci numbers with negative indices) If $v + rx^4 + sx^3 + tx^2 + ux - x^5 \neq 0$, then*

$$\sum_{k=1}^n x^k W_{-k} = \frac{\Theta_4(x)}{v + rx^4 + sx^3 + tx^2 + ux - x^5}$$

- (b) *If $2sx^4 - u^2x + 2ux^3 + r^2x^4 - s^2x^3 + t^2x^2 + v^2 - x^5 + 2rtx^3 + 2rvx^2 - 2sux^2 + 2tvx \neq 0$ then*

$$\sum_{k=1}^n x^k W_{-2k} = \frac{\Theta_5(x)}{2sx^4 - u^2x + 2ux^3 + r^2x^4 - s^2x^3 + t^2x^2 + v^2 - x^5 + 2rtx^3 + 2rvx^2 - 2sux^2 + 2tvx}$$

(c) If $2sx^4 - u^2x + 2ux^3 + r^2x^4 - s^2x^3 + t^2x^2 + v^2 - x^5 + 2rtx^3 + 2rvx^2 - 2sux^2 + 2tvx \neq 0$ then

$$\sum_{k=1}^n x^k W_{-2k+1} = \frac{\Theta_6(x)}{2sx^4 - u^2x + 2ux^3 + r^2x^4 - s^2x^3 + t^2x^2 + v^2 - x^5 + 2rtx^3 + 2rvx^2 - 2sux^2 + 2tvx}$$

where

$$\Theta_4(x) = -x^{n+1}W_{4-n} + (r-x)x^{n+1}W_{-n+3} + (s+rx-x^2)x^{n+1}W_{-n+2} + (t+rx^2+sx-x^3)x^{n+1}W_{-n+1} + (u+rx^3+sx^2+tx-x^4)x^{n+1}W_{-n} + xW_4 - x(r-x)W_3 + x(-s-rx+x^2)W_2 + x(-t-rx^2-sx+x^3)W_1 + x(-u-rx^3-sx^2-tx+x^4)W_0.$$

$$\Theta_5(x) = -(v+rx^2+tx)x^{n+1}W_{-2n+3} + (sx^2+r^2x^2+rv+ux-x^3+rtx)x^{n+1}W_{-2n+2} - (tx^2-sv+vx+ru-x-stx)x^{n+1}W_{-2n+1} + (2sx^3+t^2x+ux^2+r^2x^3-s^2x^2+tv-x^4+2rtx^2+rvx-sux)x^{n+1}W_{-2n} + v(u+sx-x^2)x^{n+1}W_{-2n-1} + x(-u-sx+x^2)W_4 + x(v+ru+tx+rsx)W_3 + x(-2sx^2+s^2x-r^2x^2-rv+su-ux+x^3-rtx)W_2 + x(-sv+tu+vx+ru)W_1 + x(-2sx^3-t^2x-2ux^2-r^2x^3+s^2x^2-tv+u^2+x^4-2rtx^2-rvx+2sux)W_0.$$

$$\Theta_6(x) = x^{n+2}(u+sx-x^2)W_{-2n+3} - (v+ru+tx+rsx)x^{n+2}W_{-2n+2} + (2sx^2-s^2x+r^2x^2+rv-su+ux-x^3+rtx)x^{n+2}W_{-2n+1} + (sv-tu-vx-ru)W_{-2n} - v(v+rx^2+tx)x^{n+1}W_{-2n-1} + x(v+rx^2+tx)W_4 + x(-sx^2-r^2x^2-rv-ux+x^3-rtx)W_3 + x(tx^2-sv+vx+ru-x-stx)W_2 + x(-2sx^3-t^2x-ux^2-r^2x^3+s^2x^2-tv+x^4-2rtx^2-rvx+2sux)W_1 + vx(-u-sx+x^2)W_0.$$

Proof.

(a) Using the recurrence relation

$$W_{-n} = \frac{1}{v}W_{-n+5} - \frac{u}{v}W_{-n+1} - \frac{t}{v}W_{-n+2} - \frac{s}{v}W_{-n+3} - \frac{r}{v}W_{-n+4}$$

i.e.

$$vW_{-n} = W_{-n+5} - rW_{-n+4} - sW_{-n+3} - tW_{-n+2} - uW_{-n+1}$$

we obtain

$$\begin{aligned} vx^n W_{-n} &= x^n W_{-n+5} - rx^n W_{-n+4} - sx^n W_{-n+3} - tx^n W_{-n+2} - ux^n W_{-n+1} \\ vx^{n-1} W_{-n+1} &= x^{n-1} W_{-n+6} - rx^{n-1} W_{-n+5} - sx^{n-1} W_{-n+4} - tx^{n-1} W_{-n+3} - ux^{n-1} W_{-n+2} \\ vx^{n-2} W_{-n+2} &= x^{n-2} W_{-n+7} - rx^{n-2} W_{-n+6} - sx^{n-2} W_{-n+5} - tx^{n-2} W_{-n+4} - ux^{n-2} W_{-n+3} \\ &\vdots \\ vx^5 W_{-5} &= x^5 W_0 - rx^5 W_{-1} - sx^5 W_{-2} - tx^5 W_{-3} - ux^5 W_{-4} \\ vx^4 W_{-4} &= x^4 W_1 - rx^4 W_0 - sx^4 W_{-1} - tx^4 W_{-2} - ux^4 W_{-3} \\ vx^3 W_{-3} &= x^3 W_2 - rW_1 - sx^3 W_0 - tx^3 W_{-1} - ux^3 W_{-2} \\ vx^2 W_{-2} &= x^2 W_3 - rx^2 W_2 - sx^2 W_1 - tx^2 W_0 - ux^2 W_{-1} \\ vx^1 W_{-1} &= x^1 W_4 - rx^1 W_3 - sx^1 W_2 - tx^1 W_1 - ux^1 W_0. \end{aligned}$$

If we add the equations side by side, we get (a).

(b) and (c) Using the recurrence relation

$$\begin{aligned} W_{-n+5} &= rW_{-n+4} + sW_{-n+3} + tW_{-n+2} + uW_{-n+1} + vW_{-n} \\ \Rightarrow W_{-n} &= \frac{1}{v}W_{-n+5} - \frac{u}{v}W_{-n+1} - \frac{t}{v}W_{-n+2} - \frac{s}{v}W_{-n+3} - \frac{r}{v}W_{-n+4} \end{aligned}$$

i.e.

$$uW_{-n+1} = W_{-n+5} - rW_{-n+4} - sW_{-n+3} - tW_{-n+2} - vW_{-n}$$

we obtain

$$\begin{aligned}
ux^n W_{-2n+1} &= x^n W_{-2n+5} - rx^n W_{-2n+4} - sx^n W_{-2n+3} - tx^n W_{-2n+2} - vx^n W_{-2n} \\
ux^{n-1} W_{-2n+3} &= x^{n-1} W_{-2n+7} - rx^{n-1} W_{-2n+6} - sx^{n-1} W_{-2n+5} - tx^{n-1} W_{-2n+4} - vx^{n-1} W_{-2n+2} \\
ux^{n-2} W_{-2n+5} &= x^{n-2} W_{-2n+9} - rx^{n-2} W_{-2n+8} - sx^{n-2} W_{-2n+7} - tx^{n-2} W_{-2n+6} - vx^{n-2} W_{-2n+4} \\
ux^{n-3} W_{-2n+7} &= x^{n-3} W_{-2n+11} - rx^{n-3} W_{-2n+10} - sx^{n-3} W_{-2n+9} - tx^{n-3} W_{-2n+8} - vx^{n-3} W_{-2n+6} \\
&\vdots \\
ux^3 W_{-5} &= x^3 W_{-1} - rx^3 W_{-2} - sx^3 W_{-3} - tx^3 W_{-4} - vx^3 W_{-6} \\
ux^2 W_{-3} &= x^2 W_1 - rx^2 W_0 - sx^2 W_{-1} - tx^2 W_{-2} - vx^2 W_{-4} \\
ux^1 W_{-1} &= x^1 W_3 - rx^1 W_2 - sx^1 W_1 - tx^1 W_0 - vx^1 W_{-2}
\end{aligned}$$

If we add the equations side by side, we get

$$\begin{aligned}
u \sum_{k=1}^n x^k W_{-2k+1} &= (-x^{n+1} W_{-2n+3} - x^{n+2} W_{-2n+1} + x^1 W_3 + x^2 W_1 + \sum_{k=1}^n x^{k+2} W_{-2k+1}) \\
&\quad -r(-x^{n+1} W_{-2n+2} - x^{n+2} W_{-2n} + x^2 W_0 + x^1 W_2 + \sum_{k=1}^n x^{k+2} W_{-2k}) \\
&\quad -s(-x^{n+1} W_{-2n+1} + x^1 W_1 + \sum_{k=1}^n x^{k+1} W_{-2k+1}) - t(-x^{n+1} W_{-2n} \\
&\quad + x^1 W_0 + \sum_{k=1}^n x^{k+1} W_{-2k}) - v(\sum_{k=1}^n x^k W_{-2k}).
\end{aligned}$$

Then we have

$$\begin{aligned}
u \sum_{k=1}^n x^k W_{-2k+1} &= (-x^{n+1} W_{-2n+3} - x^{n+2} W_{-2n+1} + x^1 W_3 + x^2 W_1 + x^2 \sum_{k=1}^n x^k W_{-2k+1}) \quad (4.1) \\
&\quad -r(-x^{n+1} W_{-2n+2} - x^{n+2} W_{-2n} + x^2 W_0 + x^1 W_2 + x^2 \sum_{k=1}^n x^k W_{-2k}) \\
&\quad -s(-x^{n+1} W_{-2n+1} + x^1 W_1 + x^1 \sum_{k=1}^n x^k W_{-2k+1}) \\
&\quad -t(-x^{n+1} W_{-2n} + x^1 W_0 + x^1 \sum_{k=1}^n x^k W_{-2k}) - v(\sum_{k=1}^n x^k W_{-2k}).
\end{aligned}$$

Similarly, using the recurrence relation

$$\begin{aligned}
W_{-n+5} &= rW_{-n+4} + sW_{-n+3} + tW_{-n+2} + uW_{-n+1} + vW_{-n} \\
\Rightarrow W_{-n} &= \frac{1}{v}W_{-n+5} - \frac{u}{v}W_{-n+1} - \frac{t}{v}W_{-n+2} - \frac{s}{v}W_{-n+3} - \frac{r}{v}W_{-n+4}
\end{aligned}$$

i.e.

$$uW_{-n+1} = W_{-n+5} - rW_{-n+4} - sW_{-n+3} - tW_{-n+2} - vW_{-n}$$

we obtain

$$\begin{aligned}
ux^n W_{-2n} &= x^n W_{-2n+4} - rx^n W_{-2n+3} - sx^n W_{-2n+2} - tx^n W_{-2n+1} - vx^n W_{-2n-1} \\
ux^{n-1} W_{-2n+2} &= x^{n-1} W_{-2n+6} - rx^{n-1} W_{-2n+5} - x^{n-1} s W_{-2n+4} - tx^{n-1} W_{-2n+3} - vx^{n-1} W_{-2n+1} \\
ux^{n-2} W_{-2n+4} &= x^{n-2} W_{-2n+8} - rx^{n-2} W_{-2n+7} - sx^{n-2} W_{-2n+6} - tx^{n-2} W_{-2n+5} - vx^{n-2} W_{-2n+3} \\
ux^{n-3} W_{-2n+6} &= x^{n-3} W_{-2n+10} - rx^{n-3} W_{-2n+9} - sx^{n-3} W_{-2n+8} - tx^{n-3} W_{-2n+7} - vx^{n-3} W_{-2n+5} \\
&\vdots \\
ux^4 W_{-8} &= x^4 W_{-4} - rx^4 W_{-5} - sx^4 W_{-6} - tx^4 W_{-7} - vx^4 W_{-9} \\
ux^3 W_{-6} &= x^3 W_{-2} - rx^3 W_{-3} - sx^3 W_{-4} - tx^3 W_{-5} - vx^3 W_{-7} \\
ux^2 W_{-4} &= x^2 W_0 - rx^2 W_{-1} - sx^2 W_{-2} - tx^2 W_{-3} - vx^2 W_{-5} \\
ux^1 W_{-2} &= x^1 W_2 - rx^1 W_1 - sx^1 W_0 - tx^1 W_{-1} - vx^1 W_{-3}.
\end{aligned}$$

If we add the equations side by side, we get

$$\begin{aligned}
u \sum_{k=1}^n x^k W_{-2k} &= (-x^{n+1} W_{-2n+2} - x^{n+2} W_{-2n} + x^1 W_2 + x^2 W_0 + \sum_{k=1}^n x^{k+2} W_{-2k}) \\
&\quad -r(-x^{n+1} W_{-2n+1} + x^1 W_1 + \sum_{k=1}^n x^{k+1} W_{-2k+1}) \\
&\quad -s(-x^{n+1} W_{-2n} + x^1 W_0 + \sum_{k=1}^n x^{k+1} W_{-2k}) - t(\sum_{k=1}^n x^k W_{-2k+1}) \\
&\quad -v(x^n W_{-2n-1} - x^0 W_{-1} + \sum_{k=1}^n x^{k-1} W_{-2k+1}).
\end{aligned}$$

Since

$$W_{-1} = -\frac{u}{v} W_0 - \frac{t}{v} W_1 - \frac{s}{v} W_2 - \frac{r}{v} W_3 + \frac{1}{v} W_4$$

it follows that

$$\begin{aligned}
u \sum_{k=1}^n x^k W_{-2k} &= (-x^{n+1} W_{-2n+2} - x^{n+2} W_{-2n} + x^1 W_2 + x^2 W_0 + x^2 \sum_{k=1}^n x^k W_{-2k}) \quad (4.2) \\
&\quad -r(-x^{n+1} W_{-2n+1} + x^1 W_1 + x^1 \sum_{k=1}^n x^k W_{-2k+1}) \\
&\quad -s(-x^{n+1} W_{-2n} + x^1 W_0 + x^1 \sum_{k=1}^n x^k W_{-2k}) - t(\sum_{k=1}^n x^k W_{-2k+1}) \\
&\quad -v(x^n W_{-2n-1} - x^0 (-\frac{u}{v} W_0 - \frac{t}{v} W_1 - \frac{s}{v} W_2 - \frac{r}{v} W_3 + \frac{1}{v} W_4) + x^{-1} \sum_{k=1}^n x^k W_{-2k+1}).
\end{aligned}$$

Then, solving system (4.1)-(4.2) the required result of (b) and (c) follow. \square

5 Specific Cases

In this section, for the specific cases of x , we present the closed form solutions (identities) of the sums $\sum_{k=1}^n x^k W_{-k}$, $\sum_{k=1}^n x^k W_{-2k}$ and $\sum_{k=1}^n x^k W_{-2k+1}$ for the specific case of sequence $\{W_n\}$.

5.1 The case $x = 1$

In this subsection, we consider the special case $x = 1$.

The case $x = 1$ of Theorem 4.1 is given in Soykan [21].

5.2 The case $x = -1$

In this subsection we consider the special case $x = -1$.

Taking $r = s = t = u = v = 1$ in Theorem 4.1 (a), (b) and (c), we obtain the following Proposition.

Proposition 5.1. *If $r = s = t = u = v = 1$ then for $n \geq 1$ we have the following formulas:*

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{2}((-1)^n (W_{-n+4} - 2W_{-n+3} + W_{-n+2} - 2W_{-n+1} + W_{-n}) - W_4 + 2W_3 - W_2 + 2W_1 - W_0)$.
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{2}((-1)^n (W_{-2n+3} - 2W_{-2n+2} - W_{-2n+1} + 2W_{-2n} + W_{-2n-1}) - W_4 + 3W_2 + 2W_1 - W_0)$.
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{2}((-1)^n (-W_{-2n+3} + 3W_{-2n+1} + 2W_{-2n} + W_{-2n-1}) - W_4 + 2W_3 + W_2 - 2W_1 - W_0)$.

From the above Proposition, we have the following Corollary which gives linear sum formulas of Pentanacci numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2, P_4 = 4$).

Corollary 5.2. *For $n \geq 1$, Pentanacci numbers have the following properties.*

- (a) $\sum_{k=1}^n (-1)^k P_{-k} = \frac{1}{2}((-1)^n (P_{-n+4} - 2P_{-n+3} + P_{-n+2} - 2P_{-n+1} + P_{-n}) + 1)$.
- (b) $\sum_{k=1}^n (-1)^k P_{-2k} = \frac{1}{2}((-1)^n (P_{-2n+3} - 2P_{-2n+2} - P_{-2n+1} + 2P_{-2n} + P_{-2n-1}) + 1)$.
- (c) $\sum_{k=1}^n (-1)^k P_{-2k+1} = \frac{1}{2}((-1)^n (-P_{-2n+3} + 3P_{-2n+1} + 2P_{-2n} + P_{-2n-1}) - 1)$.

Taking $W_n = Q_n$ with $Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7, Q_4 = 15$ in the above Proposition, we have the following Corollary which presents linear sum formulas of Pentanacci-Lucas numbers.

Corollary 5.3. *For $n \geq 1$, Pentanacci-Lucas numbers have the following properties.*

- (a) $\sum_{k=1}^n (-1)^k Q_{-k} = \frac{1}{2}((-1)^n (Q_{-n+4} - 2Q_{-n+3} + Q_{-n+2} - 2Q_{-n+1} + Q_{-n}) - 7)$.
- (b) $\sum_{k=1}^n (-1)^k Q_{-2k} = \frac{1}{2}((-1)^n (Q_{-2n+3} - 2Q_{-2n+2} - Q_{-2n+1} + 2Q_{-2n} + Q_{-2n-1}) - 9)$.
- (c) $\sum_{k=1}^n (-1)^k Q_{-2k+1} = \frac{1}{2}((-1)^n (-Q_{-2n+3} + 3Q_{-2n+1} + 2Q_{-2n} + Q_{-2n-1}) - 5)$.

Taking $r = 2, s = t = u = v = 1$ in Theorem 4.1 (a), (b) and (c), we obtain the following Proposition.

Proposition 5.4. *If $r = 2, s = t = u = v = 1$ then for $n \geq 1$ we have the following formulas:*

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{3}((-1)^n (W_{-n+4} - 3W_{-n+3} + 2W_{-n+2} - 3W_{-n+1} + 2W_{-n}) - W_4 + 3W_3 - 2W_2 + 3W_1 - 2W_0)$.
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{5}((-1)^n (2W_{3-2n} - 5W_{2-2n} - 2W_{1-2n} + 4W_{-2n} + W_{-2n-1}) - W_4 + 6W_2 + 3W_1 - 3W_0)$.
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{5}((-1)^n (-W_{3-2n} + 6W_{1-2n} + 3W_{-2n} + 2W_{-2n-1}) - 2W_4 + 5W_3 + 2W_2 - 4W_1 - W_0)$.

From the last Proposition, we have the following Corollary which gives linear sum formulas of fifth-order Pell numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5, P_4 = 13$).

Corollary 5.5. *For $n \geq 1$, fifth-order Pell numbers have the following properties:*

- (a) $\sum_{k=1}^n (-1)^k P_{-k} = \frac{1}{3}((-1)^n (P_{-n+4} - 3P_{-n+3} + 2P_{-n+2} - 3P_{-n+1} + 2P_{-n}) + 1)$.
- (b) $\sum_{k=1}^n (-1)^k P_{-2k} = \frac{1}{5}((-1)^n (2P_{3-2n} - 5P_{2-2n} - 2P_{1-2n} + 4P_{-2n} + P_{-2n-1}) + 2)$.

$$(c) \sum_{k=1}^n (-1)^k P_{-2k+1} = \frac{1}{5}((-1)^n (-P_{3-2n} + 6P_{1-2n} + 3P_{-2n} + 2P_{-2n-1}) - 1).$$

Taking $W_n = Q_n$ with $Q_0 = 5, Q_1 = 2, Q_2 = 6, Q_3 = 17, Q_4 = 46$ in the last Proposition, we have the following Corollary which presents linear sum formulas of fifth-order Pell-Lucas numbers.

Corollary 5.6. *For $n \geq 1$, fifth-order Pell-Lucas numbers have the following properties:*

$$(a) \sum_{k=1}^n (-1)^k Q_{-k} = \frac{1}{3}((-1)^n (Q_{-n+4} - 3Q_{-n+3} + 2Q_{-n+2} - 3Q_{-n+1} + 2Q_{-n}) - 11).$$

$$(b) \sum_{k=1}^n (-1)^k Q_{-2k} = \frac{1}{5}((-1)^n (2Q_{3-2n} - 5Q_{2-2n} - 2Q_{1-2n} + 4Q_{-2n} + Q_{-2n-1}) - 19).$$

$$(c) \sum_{k=1}^n (-1)^k Q_{-2k+1} = \frac{1}{5}((-1)^n (-Q_{3-2n} + 6Q_{1-2n} + 3Q_{-2n} + 2Q_{-2n-1}) - 8).$$

Taking $r = s = t = 1, u = 1, v = 2$ in Theorem 4.1 (a), (b) and (c), we obtain the following Proposition.

Proposition 5.7. *If $r = s = t = 1, u = 1, v = 2$ then for $n \geq 1$ we have the following formulas:*

$$(a) \sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{3}((-1)^n (W_{-n+4} - 2W_{-n+3} + W_{-n+2} - 2W_{-n+1} + W_{-n}) - W_4 + 2W_3 - W_2 + 2W_1 - W_0).$$

$$(b) \sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{5}((-1)^n (2W_{-2n+3} - 3W_{-2n+2} - 3W_{-2n+1} + 2W_{-2n} + 2W_{-2n-1}) - W_4 - W_3 + 4W_2 + 4W_1 - W_0).$$

$$(c) \sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{5}((-1)^n (-W_{-2n+3} - W_{-2n+2} + 4W_{-2n+1} + 4W_{-2n} + 4W_{-2n-1}) - 2W_4 + 3W_3 + 3W_2 - 2W_1 - 2W_0).$$

Taking $W_n = J_n$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1, J_4 = 1$ in the last Proposition, we have the following Corollary which presents linear sum formulas of fifth-order Jacobsthal numbers.

Corollary 5.8. *For $n \geq 1$, fifth order Jacobsthal numbers have the following properties:*

$$(a) \sum_{k=1}^n (-1)^k J_{-k} = \frac{1}{3}((-1)^n (J_{-n+4} - 2J_{-n+3} + J_{-n+2} - 2J_{-n+1} + J_{-n}) + 2).$$

$$(b) \sum_{k=1}^n (-1)^k J_{-2k} = \frac{1}{5}((-1)^n (2J_{-2n+3} - 3J_{-2n+2} - 3J_{-2n+1} + 2J_{-2n} + 2J_{-2n-1}) + 6).$$

$$(c) \sum_{k=1}^n (-1)^k J_{-2k+1} = \frac{1}{5}((-1)^n (-J_{-2n+3} - J_{-2n+2} + 4J_{-2n+1} + 4J_{-2n} + 4J_{-2n-1}) + 2).$$

From the last Proposition, we have the following Corollary which gives linear sum formulas of fifth order Jacobsthal-Lucas numbers (take $W_n = j_n$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10, j_4 = 20$).

Corollary 5.9. *For $n \geq 1$, fifth order Jacobsthal-Lucas numbers have the following properties:*

$$(a) \sum_{k=1}^n (-1)^k j_{-k} = \frac{1}{3}((-1)^n (j_{-n+4} - 2j_{-n+3} + j_{-n+2} - 2j_{-n+1} + j_{-n}) - 5).$$

$$(b) \sum_{k=1}^n (-1)^k j_{-2k} = \frac{1}{5}((-1)^n (2j_{-2n+3} - 3j_{-2n+2} - 3j_{-2n+1} + 2j_{-2n} + 2j_{-2n-1}) - 8).$$

$$(c) \sum_{k=1}^n (-1)^k j_{-2k+1} = \frac{1}{5}((-1)^n (-j_{-2n+3} - j_{-2n+2} + 4j_{-2n+1} + 4j_{-2n} + 4j_{-2n-1}) - 1).$$

Taking $W_n = K_n$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10, K_4 = 20$ in the last proposition, we have the following corollary which presents linear sum formula of modified fifth order Jacobsthal numbers.

Corollary 5.10. *For $n \geq 1$, modified fifth order Jacobsthal numbers have the following property:*

$$(a) \sum_{k=1}^n (-1)^k K_{-k} = \frac{1}{3}((-1)^n (K_{-n+4} - 2K_{-n+3} + K_{-n+2} - 2K_{-n+1} + K_{-n}) - 4).$$

$$(b) \sum_{k=1}^n (-1)^k K_{-2k} = \frac{1}{5}((-1)^n (2K_{-2n+3} - 3K_{-2n+2} - 3K_{-2n+1} + 2K_{-2n} + 2K_{-2n-1}) - 17).$$

$$(c) \sum_{k=1}^n (-1)^k K_{-2k+1} = \frac{1}{5}((-1)^n (-K_{-2n+3} - K_{-2n+2} + 4K_{-2n+1} + 4K_{-2n} + 4K_{-2n-1}) - 9).$$

From the last proposition, we have the following corollary which gives linear sum formula of fifth-order Jacobsthal Perrin numbers (take $W_n = Q_n$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8, Q_4 = 16$).

Corollary 5.11. For $n \geq 1$, fifth-order Jacobsthal Perrin numbers have the following property:

- (a) $\sum_{k=1}^n (-1)^k Q_{-k} = \frac{1}{3}((-1)^n (Q_{-n+4} - 2Q_{-n+3} + Q_{-n+2} - 2Q_{-n+1} + Q_{-n}) - 5)$.
- (b) $\sum_{k=1}^n (-1)^k Q_{-2k} = \frac{1}{5}((-1)^n (2Q_{-2n+3} - 3Q_{-2n+2} - 3Q_{-2n+1} + 2Q_{-2n} + 2Q_{-2n-1}) - 19)$.
- (c) $\sum_{k=1}^n (-1)^k Q_{-2k+1} = \frac{1}{5}((-1)^n (-Q_{-2n+3} - Q_{-2n+2} + 4Q_{-2n+1} + 4Q_{-2n} + 4Q_{-2n-1}) - 8)$.

Taking $W_n = S_n$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2, S_4 = 4$ in the proposition, we have the following corollary which presents linear sum formula of adjusted fifth-order Jacobsthal numbers.

Corollary 5.12. For $n \geq 1$, adjusted fifth-order Jacobsthal numbers have the following property:

- (a) $\sum_{k=1}^n (-1)^k S_{-k} = \frac{1}{3}((-1)^n (S_{-n+4} - 2S_{-n+3} + S_{-n+2} - 2S_{-n+1} + S_{-n}) + 1)$.
- (b) $\sum_{k=1}^n (-1)^k S_{-2k} = \frac{1}{5}((-1)^n (2S_{-2n+3} - 3S_{-2n+2} - 3S_{-2n+1} + 2S_{-2n} + 2S_{-2n-1}) + 2)$.
- (c) $\sum_{k=1}^n (-1)^k S_{-2k+1} = \frac{1}{5}((-1)^n (-S_{-2n+3} - S_{-2n+2} + 4S_{-2n+1} + 4S_{-2n} + 4S_{-2n-1}) - 1)$.

From the last proposition, we have the following corollary which gives linear sum formula of modified fifth-order Jacobsthal-Lucas numbers (take $W_n = R_n$ with $R_0 = 5, R_1 = 1, R_2 = 3, R_3 = 7, R_4 = 15$).

Corollary 5.13. For $n \geq 1$, modified fifth-order Jacobsthal-Lucas numbers have the following property:

- (a) $\sum_{k=1}^n (-1)^k R_{-k} = \frac{1}{3}((-1)^n (R_{-n+4} - 2R_{-n+3} + R_{-n+2} - 2R_{-n+1} + R_{-n}) - 7)$.
- (b) $\sum_{k=1}^n (-1)^k R_{-2k} = \frac{1}{5}((-1)^n (2R_{-2n+3} - 3R_{-2n+2} - 3R_{-2n+1} + 2R_{-2n} + 2R_{-2n-1}) - 11)$.
- (c) $\sum_{k=1}^n (-1)^k R_{-2k+1} = \frac{1}{5}((-1)^n (-R_{-2n+3} - R_{-2n+2} + 4R_{-2n+1} + 4R_{-2n} + 4R_{-2n-1}) - 12)$.

Taking $r = 2, s = 3, t = 5, u = 7, v = 11$ in Theorem 4.1 (a), (b) and (c), we obtain the following proposition.

Proposition 5.14. If $r = 2, s = 3, t = 5, u = 7, v = 11$ then for $n \geq 1$ we have the following formulas:

- (a) $\sum_{k=1}^n (-1)^k W_{-k} = \frac{1}{9}((-1)^n (W_{-n+4} - 3W_{-n+3} - 5W_{-n+1} - 2W_{-n}) - W_4 + 3W_3 + 5W_1 + 2W_0)$.
- (b) $\sum_{k=1}^n (-1)^k W_{-2k} = \frac{1}{73}((-1)^n (8W_{-2n+3} - 13W_{-2n+2} - 38W_{-2n+1} - 36W_{-2n} - 33W_{-2n-1}) + 3W_4 - 14W_3 + 4W_2 + 23W_1 + 15W_0)$.
- (c) $\sum_{k=1}^n (-1)^k W_{-2k+1} = \frac{1}{73}((-1)^n (+3W_{-2n+3} - 14W_{-2n+2} + 4W_{-2n+1} + 23W_{-2n} + 88W_{-2n-1}) - 8W_4 + 13W_3 + 38W_2 + 36W_1 + 33W_0)$.

From the last proposition, we have the following corollary which gives linear sum formulas of 5-primes numbers (take $W_n = G_n$ with $G_0 = 0, G_1 = 0, G_2 = 0, G_3 = 1, G_4 = 2$).

Corollary 5.15. For $n \geq 1$, 5-primes numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k G_{-k} = \frac{1}{9}((-1)^n (G_{-n+4} - 3G_{-n+3} - 5G_{-n+1} - 2G_{-n}) + 1)$.
- (b) $\sum_{k=1}^n (-1)^k G_{-2k} = \frac{1}{73}((-1)^n (8G_{-2n+3} - 13G_{-2n+2} - 38G_{-2n+1} - 36G_{-2n} - 33G_{-2n-1}) - 8)$.
- (c) $\sum_{k=1}^n (-1)^k G_{-2k+1} = \frac{1}{73}((-1)^n (+3G_{-2n+3} - 14G_{-2n+2} + 4G_{-2n+1} + 23G_{-2n} + 88G_{-2n-1}) - 3)$.

Taking $W_n = H_n$ with $H_0 = 5, H_1 = 2, H_2 = 10, H_3 = 41, H_4 = 150$ in the last proposition, we have the following corollary which presents linear sum formulas of Lucas 5-primes numbers.

Corollary 5.16. For $n \geq 1$, Lucas 5-primes numbers have the following properties:

- (a) $\sum_{k=1}^n (-1)^k H_{-k} = \frac{1}{9}((-1)^n (H_{-n+4} - 3H_{-n+3} - 5H_{-n+1} - 2H_{-n}) - 7)$.

$$(b) \sum_{k=1}^n (-1)^k H_{-2k} = \frac{1}{73}((-1)^n (8H_{-2n+3} - 13H_{-2n+2} - 38H_{-2n+1} - 36H_{-2n} - 33H_{-2n-1}) + 37).$$

$$(c) \sum_{k=1}^n (-1)^k H_{-2k+1} = \frac{1}{73}((-1)^n (+3H_{-2n+3} - 14H_{-2n+2} + 4H_{-2n+1} + 23H_{-2n} + 88H_{-2n-1}) - 50).$$

From the last proposition, we have the following corollary which gives linear sum formulas of modified 5-primes numbers (take $W_n = E_n$ with $E_0 = 0, E_1 = 0, E_2 = 0, E_3 = 1, E_4 = 1$).

Corollary 5.17. *For $n \geq 1$, modified 5-primes numbers have the following properties:*

$$(a) \sum_{k=1}^n (-1)^k E_{-k} = \frac{1}{9}((-1)^n (E_{-n+4} - 3E_{-n+3} - 5E_{-n+1} - 2E_{-n}) + 2).$$

$$(b) \sum_{k=1}^n (-1)^k E_{-2k} = \frac{1}{73}((-1)^n (8E_{-2n+3} - 13E_{-2n+2} - 38E_{-2n+1} - 36E_{-2n} - 33E_{-2n-1}) - 11).$$

$$(c) \sum_{k=1}^n (-1)^k E_{-2k+1} = \frac{1}{73}((-1)^n (+3E_{-2n+3} - 14E_{-2n+2} + 4E_{-2n+1} + 23E_{-2n} + 88E_{-2n-1}) + 5).$$

5.3 The case $x = i$

In this subsection, we consider the special case $x = i$.

Taking $r = s = t = u = v = 1$ in Theorem 4.1, we obtain the following proposition.

Proposition 5.18. *If $r = s = t = u = v = 1$ then for $n \geq 1$ we have the following formulas:*

$$(a) \sum_{k=1}^n i^k W_{-k} = \frac{1}{1-i}(i^n(-iW_{-n+4} + (1+i)W_{-n+3} - (1-2i)W_{-n+2} - 2W_{-n+1} - iW_{-n}) + iW_4 - (1+i)W_3 + (1-2i)W_2 + 2W_1 + iW_0).$$

$$(b) \sum_{k=1}^n i^k W_{-2k} = \frac{1}{3-3i}(i^n(W_{-2n+3} - (3+i)W_{-2n+2} + (1+2i)W_{-2n+1} + (2-2i)W_{-2n} - (1-2i)W_{-2n-1}) + (1-2i)W_4 - (2-2i)W_3 + (2+3i)W_2 - 2W_1 - (3-4i)W_0).$$

$$(c) \sum_{k=1}^n i^k W_{-2k+1} = \frac{1}{3-3i}(i^n(-(2+i)W_{-2n+3} + (2+2i)W_{-2n+2} + (3-2i)W_{-2n+1} + 2iW_{-2n} + W_{-2n-1}) - W_4 + (3+i)W_3 - (1+2i)W_2 - (2-2i)W_1 + (1-2i)W_0).$$

From the above Proposition, we have the following Corollary which gives sum formulas of Pentanacci numbers (take $W_n = P_n$ with $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2, P_4 = 4$).

Corollary 5.19. *For $n \geq 1$, Pentanacci numbers have the following properties.*

$$(a) \sum_{k=1}^n i^k P_{-k} = \frac{1}{1-i}(i^n(-iP_{-n+4} + (1+i)P_{-n+3} - (1-2i)P_{-n+2} - 2P_{-n+1} - iP_{-n}) + 1).$$

$$(b) \sum_{k=1}^n i^k P_{-2k} = \frac{1}{3-3i}(i^n(P_{-2n+3} - (3+i)P_{-2n+2} + (1+2i)P_{-2n+1} + (2-2i)P_{-2n} - (1-2i)P_{-2n-1}) - i).$$

$$(c) \sum_{k=1}^n i^k P_{-2k+1} = \frac{1}{3-3i}(i^n(-(2+i)P_{-2n+3} + (2+2i)P_{-2n+2} + (3-2i)P_{-2n+1} + 2iP_{-2n} + P_{-2n-1}) + (-1+2i)).$$

Taking $W_n = Q_n$ with $Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7, Q_4 = 15$ in the above Proposition, we have the following Corollary which presents sum formulas of Pentanacci-Lucas numbers.

Corollary 5.20. *For $n \geq 1$, Pentanacci-Lucas numbers have the following properties.*

$$(a) \sum_{k=1}^n i^k Q_{-k} = \frac{1}{1-i}(i^n(-iQ_{-n+4} + (1+i)Q_{-n+3} - (1-2i)Q_{-n+2} - 2Q_{-n+1} - iQ_{-n}) + (-2+7i)).$$

$$(b) \sum_{k=1}^n i^k Q_{-2k} = \frac{1}{3-3i}(i^n(Q_{-2n+3} - (3+i)Q_{-2n+2} + (1+2i)Q_{-2n+1} + (2-2i)Q_{-2n} - (1-2i)Q_{-2n-1}) + (-10+13i)).$$

$$(c) \sum_{k=1}^n i^k Q_{-2k+1} = \frac{1}{3-3i}(i^n(-(2+i)Q_{-2n+3} + (2+2i)Q_{-2n+2} + (3-2i)Q_{-2n+1} + 2iQ_{-2n} + Q_{-2n-1}) + (6-7i)).$$

Corresponding sums of the other fifth order generalized Pentanacci numbers can be calculated similarly.

References

- [1] Akbulak, M., Öteleş. A., On the sum of Pell and Jacobsthal numbers by matrix method, Bulletin of the Iranian Mathematical Society, 40 (4), 1017-1025, 2014.
- [2] Cook C. K., Bacon, M. R., Some identities for Jacobsthal and Jacobsthal-Lucas numbers satisfying higher order recurrence relations, Annales Mathematicae et Informaticae, 41, 27–39, 2013.
- [3] Frontczak, R., Sums of Tribonacci and Tribonacci-Lucas Numbers, International Journal of Mathematical Analysis, 12 (1), 19-24, 2018.
- [4] Gökbaş, H., Köse, H., Some Sum Formulas for Products of Pell and Pell-Lucas Numbers, Int. J. Adv. Appl. Math. and Mech. 4(4), 1-4, 2017.
- [5] Hansen., R.T., General Identities for Linear Fibonacci and Lucas Summations, Fibonacci Quarterly, 16(2), 121-28, 1978.
- [6] Koshy, T., Fibonacci and Lucas Numbers with Applications, A Wiley-Interscience Publication, New York, 2001.
- [7] Koshy, T., Pell and Pell-Lucas Numbers with Applications, Springer, New York, 2014.
- [8] Melham, R. S., Some Analogs of the Identity $F_n^2 + F_{n+1}^2 = F_{2n+1}^2$, Fibonacci Quarterly, 305-311, 1999.
- [9] Natividad, L. R., On Solving Fibonacci-Like Sequences of Fourth, Fifth and Sixth Order, International Journal of Mathematics and Computing, 3 (2), 2013.
- [10] Parpar, T., k'ncü Mertebeden Rekürans Bağıntısının Özellikleri ve Bazı Uygulamaları, Selçuk Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, 2011.
- [11] Rathore, G.P.S., Sikhwal, O., Choudhary, R., Formula for finding nth Term of Fibonacci-Like Sequence of Higher Order, International Journal of Mathematics And its Applications, 4 (2-D), 75-80, 2016.
- [12] Sloane, N.J.A., The on-line encyclopedia of integer sequences. Available: <http://oeis.org/>
- [13] Soykan, Y., On Summing Formulas For Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers, Advances in Research, 20(2), 1-15, 2019.
- [14] Soykan, Y., Corrigendum: On Summing Formulas for Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers, Advances in Research, 21(10), 66-82, 2020. DOI: 10.9734/AIR/2020/v21i1030253
- [15] Soykan, Y., On Summing Formulas for Horadam Numbers, Asian Journal of Advanced Research and Reports 8(1): 45-61, 2020, DOI: 10.9734/AJARR/2020/v8i130192.
- [16] Soykan, Y., Generalized Fibonacci Numbers: Sum Formulas, Journal of Advances in Mathematics and Computer Science, 35(1), 89-104, 2020, DOI: 10.9734/JAMCS/2020/v35i130241.
- [17] Soykan Y., Generalized Tribonacci Numbers: Summing Formulas, Int. J. Adv. Appl. Math. and Mech. 7(3), 57-76, 2020.
- [18] Soykan, Y., Summing Formulas For Generalized Tribonacci Numbers, Universal Journal of Mathematics and Applications, 3(1), 1-11, 2020. ISSN 2619-9653, DOI: <https://doi.org/10.32323/ujma.637876>
- [19] Soykan, Y., On Sum Formulas for Generalized Tribonacci Sequence, Journal of Scientific Research & Reports, 26(7), 27-52, 2020. ISSN: 2320-0227, DOI: 10.9734/JSRR/2020/v26i730283
- [20] Soykan, Y., Summation Formulas For Generalized Tetranacci Numbers, Asian Journal of Advanced Research and Reports, 7(2), 1-12, 2019. doi.org/10.9734/ajarr/2019/v7i230170.

-
- [21] Soykan, Y., Sum Formulas For Generalized Fifth-Order Linear Recurrence Sequences, *Journal of Advances in Mathematics and Computer Science*, 34(5), 1-14, 2019; Article no.JAMCS.53303, ISSN: 2456-9968, DOI: 10.9734/JAMCS/2019/v34i530224.
- [22] Soykan, Y., Linear Summing Formulas of Generalized Pentanacci and Gaussian Generalized Pentanacci Numbers, *Journal of Advanced in Mathematics and Computer Science*, 33(3): 1-14, 2019.
- [23] Soykan, Y., On Summing Formulas of Generalized Hexanacci and Gaussian Generalized Hexanacci Numbers, *Asian Research Journal of Mathematics*, 14(4), 1-14, 2019; Article no.ARJOM.50727.
- [24] Soykan, Y., A Study On Sum Formulas of Generalized Sixth-Order Linear Recurrence Sequences, *Asian Journal of Advanced Research and Reports*, 14(2), 36-48, 2020. DOI: 10.9734/AJARR/2020/v14i230329
- [25] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, *Communications in Mathematics and Applications*, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102
- [26] Soykan, Y., On Generalized Pentanacci and Gaussian Generalized Pentanacci Numbers, *Asian Research Journal of Mathematics*, 16(9), 102-121, 2020. DOI: 10.9734/ARJOM/2020/v16i930224
- [27] Soykan, Y., Properties of Generalized Fifth-Order Pell Numbers, *Asian Research Journal of Mathematics*, 15(3), 1-18, 2019.
- [28] Soykan, Y., Polatlı, E.E., A Note on Fifth Order Jacobsthal Numbers, *IOSR Journal of Mathematics (IOSR-JM)*, 17(2), 01-23, 2021. DOI: 10.9790/5728-1702010123
- [29] Soykan, Y., A Study On Generalized 5-primes Numbers, *Journal of Scientific Perspectives*, 4(3), 185-202, 2020,, DOI: <https://doi.org/10.26900/jsp.4.017>.
- [30] Öteleş, A., Akbulak, M., A Note on Generalized k-Pell Numbers and Their Determinantal Representation, *Journal of Analysis and Number Theory*, 4(2), 153-158, 2016.
- [31] Waddill, M. E., The Tetranacci Sequence and Generalizations, *Fibonacci Quarterly*, 9-20, 1992.