



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>
Eprints ID : 15826

To link to this article : 10.1111/fwb.12606
URL: <http://dx.doi.org/10.1111/fwb.12606>

To cite this version : Cornut, Julien and Ferreira, Verónica and Gonçalves, Ana Lúcia and Chauvet, Eric and Canhoto, Cristina *Fungal alteration of the elemental composition of leaf litter affects shredder feeding activity*. (2015) *Freshwater Biology*, vol. 60 (n° 9). pp. 1755-1771. ISSN 0046-5070

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

Fungal alteration of the elemental composition of leaf litter affects shredder feeding activity

JULIEN CORNUT^{*†‡}, VERÓNICA FERREIRA[‡], ANA LÚCIA GONÇALVES[§], ERIC CHAUVET^{*†} AND CRISTINA CANHOTO[§]

^{*}Université de Toulouse, INP, UPS, EcoLab (Laboratoire Ecologie Fonctionnelle et Environnement), Toulouse, France

[†]CNRS, EcoLab, Toulouse, France

[‡]MARE - Marine and Environmental Sciences Centre, Department of Life Sciences, University of Coimbra, Coimbra, Portugal

[§]Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, Coimbra, Portugal

SUMMARY

1. Leaf litter from riparian vegetation provides the main source of matter and energy for food webs of small forest streams. Shredding macroinvertebrates mostly feed on this litter when it has been colonised and conditioned by microorganisms, especially by aquatic hyphomycetes. Since shredders feed selectively, they must make foraging decisions based on the physical and chemical characteristics of the food resource, which can change depending on the identity of fungal species.
2. Here, we addressed the effect of changes in fungal assemblage structure on the elemental composition of oak (*Quercus robur*) leaf litter and how variation in litter quality affects the feeding of a stream shredder. Leaf discs were incubated in microcosms for 2 weeks, inoculated with various fungal assemblages comprised of three species each, and offered to a shredder (*Schizopelex festiva*, Trichoptera: Sericostomatidae) as food.
3. This shredder ate more leaves with a high mycelial biomass, which depended on fungal assemblage composition. Leaf litter conditioned by different fungal assemblages resulted in different litter N and P concentrations. Mycelial biomass was positively related to litter P concentration, with the lowest and highest P concentrations differing by 40% at most, but not to litter N concentration, even though the lowest and highest N concentrations differed by as much as 35%. The caddisfly larvae ate more leaves with a low C/P ratio.
4. These findings suggest a key role of litter P concentration in eliciting fungal conditioning effects on shredder-mediated litter decomposition.

Keywords: ecosystem functioning, fungal community composition, leaf litter stoichiometry, *Schizopelex festiva*, species identity

Introduction

One challenge that all organisms face is to acquire sufficient energy and elements to maintain bioenergetic processes (Frost *et al.*, 2005). A growing body of evidence indicates that organisms meet an imbalanced mixture of elements from their resources (Sturner & Elser, 2002; Frost *et al.*, 2005). Elemental constraints can therefore strongly affect consumer growth and reproduction

(Bruning, 1991; Sturner & Schulz, 1998; Aerts & Chapin, 2000; Smith, 2002; Frost *et al.*, 2005), with important consequences for the dynamics of populations (Elser *et al.*, 1998; Chen *et al.*, 2004), interspecific interactions (DeMott & Gulati, 1999; Denno & Fagan, 2003) and key ecosystem processes (Sturner *et al.*, 1997; Cebrian, 1999; Frost *et al.*, 2005).

In small woodland streams, the primary source of carbon (C) and energy for aquatic food webs is terrestrially

derived organic matter supplied by the riparian vegetation, whose shade also limits instream primary production (Vannote *et al.*, 1980). Leaf litter decomposition in these streams is therefore an important ecosystem process, which depends on the activity of shredding invertebrates and microorganisms (Hieber & Gessner, 2002; Pascoal & Cássio, 2004). The latter, especially aquatic hyphomycetes, convert leaf C into biomass and immobilise nutrients, which increases the palatability of leaf detritus for shredding invertebrates (Gessner, Dobson & Chauvet, 1999; Graça, 2001). Therefore, the microbially conditioned leaf litter often allows a higher survival, growth rate and/or reproductive output and thus enhances shredder fitness (Graça, 2001).

Early studies showed that aquatic hyphomycete species differ in degradative capabilities when they are inoculated individually onto leaf litter (Suberkropp, Arsuffi & Anderson, 1983; Suberkropp & Arsuffi, 1984; Butler & Suberkropp, 1986). This suggests that communities dominated by distinct fungal species would decompose leaf litter at different rates and therefore differentially improve the food quality of leaf detritus. Beyond interspecific variations due to those qualitative differences in enzymatic capabilities, Danger & Chauvet (2013) recently reported a strong plasticity in the elemental composition of several aquatic hyphomycete species.

Overall, these points could have important consequences for consumer–resource interactions, the supply of C and nutrients up the food web, and nutrient recycling. Detritivores are able to discriminate among leaves colonised by different fungal species (Arsuffi & Suberkropp, 1984; Gonçalves *et al.*, 2014). Detritivores and primary consumers, in general, are among the organisms most affected by variation in resource quality. Detritivores face a world where the resources (mainly leaf litter) they consume are often of poor quality. Factors such as leaf toughness, structural constituents or secondary leaf compounds, and low nitrogen (N) concentration can affect detritivore performance (Schlief & Mutz, 2006; Danger *et al.*, 2012). Several mechanisms for overcoming low nutritional quality have been suggested (Sterner & Hessen, 1994). Consumers could actively select resources with higher concentrations of the limiting mineral nutrient (Kimmerer & Potter, 1987; Butler, Suttle & Neill, 1989) or could increase their uptake of food with low nutritional quality to access sufficient amounts of the limiting nutrient (Fink & von Elert, 2006). However, Frost & Elser (2002) suggested that such compensatory mechanisms probably cause additional energetic costs, which reduce their adaptive value. To our knowledge,

the mechanisms underlying feeding preferences of detritivores have been poorly explored.

Here, we assessed whether changes in the composition of aquatic hyphomycete assemblages would influence the nutrient concentration of decomposing oak leaf litter, and whether these variations in litter quality would affect the feeding activity of a stream detritivore. We hypothesised that variation in leaf litter resource quality, defined here as leaf litter elemental composition, would change consumer feeding performance, with consequences for the rate of leaf litter breakdown as a major ecological process in detritus-based headwater streams. We tested this hypothesis by conducting an experimental laboratory study, in which the structure of litter-associated fungal assemblages was manipulated in microcosms simulating a stream environment. The conditioned litter was then fed to a stream detritivore, the larva of the shredder *Schizopelex festiva* (Trichoptera: Sericostomatidae). We predicted that fungal assemblages comprising aquatic hyphomycete species that exhibit high degradative capabilities (Zemek *et al.*, 1985) would also better enhance leaf litter nutritional value, resulting in an increased shredder feeding activity and litter decomposition rates.

Methods

Microcosm set-up and fungal assemblages

Leaf litter, fungal assemblages and microcosms. Oak leaves were collected at abscission in autumn 2008 using a suspended net in the riparian zone of the Alzeau river, Montagne Noire (south-western France; 02°13'23" E, 43°25'51" N; altitude 743 m a.s.l.) and dried at room temperature. Before use, leaves were soaked in distilled water for 3 h and ~8000 leaf discs were cut from them with a 12-mm Ø cork borer, avoiding primary and secondary veins. Once cut, leaf discs were oven-dried (65 °C, >24 h), weighed individually (± 0.01 mg) and distributed among 63 mass classes of 0.1 mg range (from 2.10–2.20 to 8.20–8.30 mg). Eight mass classes, among which two thickness levels were distinguished, were selected: 3.30–3.40, 3.40–3.50 mg [Thin, Early assemblage treatment]; 3.50–3.60, 3.60–3.70 mg [Thin, Late assemblage]; 3.90–4.00, 4.00–4.10 mg (Thick, Early assemblage); and 4.10–4.20, 4.20–4.30 mg (Thick, Late assemblage).

Leaf discs were conditioned in laboratory microcosms designed to simulate the stream environment. Each microcosm consisted of a 1-L glass flask with a sealed cap through which passed three tubes: one tube was connected to an air pump and allowed continuous aeration

($\sim 670 \text{ mL min}^{-1}$) that created turbulence and kept the leaf discs in permanent agitation, the second tube allowed the aseptic filling of the microcosm with 500 mL of new nutrient solution every 2 days, and the third tube allowed the aseptic drainage of medium and recovery of conidial suspensions (Fig. 1). The nutrient solution contained sterile distilled water in which was dissolved $75.5 \text{ mg L}^{-1} \text{ CaCl}_2$, $10 \text{ mg L}^{-1} \text{ MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.5 g L^{-1} morpholino-propane sulphonic acid, $5.5 \text{ mg L}^{-1} \text{ K}_2\text{HPO}_4$ and $100 \text{ mg L}^{-1} \text{ KNO}_3$; pH was adjusted to 7.0 (Dang, Chauvet & Gessner, 2005).

Six species of aquatic hyphomycetes, frequently reported to be dominant in leaf litter in streams, were used to condition the leaf discs to be later offered to the caddisfly larvae: *Anguillospora filiformis* (noted as *Anguillospora*), *Articulospora tetracladia* (*Articulospora*), *Clavariopsis aquatica* (*Clavariopsis*), *Tetrachaetum elegans* (*Tetrachaetum*), *Tetracladium marchalianum* (*Tetracladium*) and *Tricladium chaetocladium* (*Tricladium*) (Gessner *et al.*, 1993; Suberkropp, 2001; Ferreira *et al.*, 2006a). The fungal strains were isolated from single conidia trapped in naturally occurring foam or released from leaf litter accumulations, collected from streams in southern France.

Growing colonies were kept at room temperature ($\sim 20 \text{ }^\circ\text{C}$), in 9-cm \varnothing Petri dishes with $\sim 10 \text{ mL}$ of sterile growth medium (2% malt and 2% agar), until they were used to induce conidial production.




Fig. 1 Laboratory microcosm designed to simulate stream conditions and used to inoculate and grow aquatic fungi axenically on leaf discs. Each microcosm consisted of a 1-L glass flask equipped with three sealed tubing systems running through the cap: one tube was connected to an air pump and allowed continuous aeration that created turbulence and kept the leaf discs in permanent agitation (air pump connection), the second tube allowed the aseptic filling of the microcosm (aseptic inlet), and the third tube allowed the aseptic drainage of medium and recovery of conidial suspensions (aseptic outlet).

Conidial inoculations (< 1 day old) were produced at $15 \text{ }^\circ\text{C}$, under light conditions, by agitation of agar plugs taken from the leading edge of 14- to 30-day-old colonies in 25 mL of nutrient solution (prepared as described previously). An aliquot of each specific conidial suspension, based on conidial density, was used to inoculate the microcosms. Eight microcosms were inoculated with an assemblage composed of aquatic hyphomycete species tending to be early colonisers (Early assemblage: *Anguillospora*, *Tetrachaetum* and *Tricladium*), and eight microcosms were inoculated with an assemblage composed of aquatic hyphomycete species tending to be late colonisers (Late assemblage: *Articulospora*, *Clavariopsis* and *Tetracladium*) (Gessner *et al.*, 1993; Nikolcheva, Cockshutt & Bärlocher, 2003). The use of only three species allowed us to assess the importance of the dominant species identity on the decomposition process (Ferreira & Chauvet, 2012) and, more specifically, the extent to which this affected the food quality of leaf detritus. For each assemblage type (Early and Late), there were four dominance treatments: an even treatment, in which microcosms were inoculated with even proportions of conidia from each species (i.e. 33.3%), and three uneven treatments in which microcosms were inoculated with species 1, species 2 or species 3 as dominant (i.e. 60% and 20% for each of the two other species, cf. Fig. 2; Ferreira & Chauvet, 2012).

Conditioning. Five groups of 155 discs were set apart from the selected mass classes, weighed ($\pm 0.01 \text{ mg}$), placed inside each microcosm with 150 mL of distilled water and autoclaved (20 min at $121 \text{ }^\circ\text{C}$). In parallel, three groups of five leaf discs of the selected mass classes were weighed ($\pm 0.01 \text{ mg}$), placed inside glass tubes with 10 mL of distilled water, given the same treatment and used to create a correction factor for mass loss due to leaching during sterilisation. After cooling to room temperature, the water from the microcosms ($n = 20$) was replaced with 500 mL of a sterile nutrient solution (described above) added to microcosms through the tubing system, which was otherwise closed with a clamp, and aerated for 24 h to allow further leaching. Nutrient solution was changed and the microcosms inoculated with conidial suspensions. Eight microcosms were inoculated with an early assemblage, and eight were inoculated with a late assemblage; a total of $\sim 155\,000$ conidia were inoculated into each microcosm (Fig. 2). For the first 6 h, the microcosms were aerated for periods of 20 min interspaced by periods of 20-min quiescence; after this, microcosms were aerated continuously until the end of the experiment. The nutrient solution was




Fig. 2 Experimental design for conditioning oak leaf discs and contribution of each aquatic hyphomycete species to the initial conidial inoculum (%). Fungal species: *Anguillospora filiformis* (*Anguillospora*), *Articulospora tetracladia* (*Articulospora*), *Clavariopsis aquatica* (*Clavariopsis*), *Tetrachaetum elegans* (*Tetrachaetum*), *Tetracladium marchalianum* (*Tetracladium*) and *Tricladium chaetocladium* (*Tricladium*).

replaced after 24 h and then every 2 days for the duration of the experiment (14 days). Microcosms were incubated at 15 °C under a 12-h light:12-h dark photoperiod. All manipulations of microcosms took place under aseptic conditions in a sterile laminar flow cabinet.

After 2 weeks, the 155 leaf discs from each microcosm were sampled. Ninety discs were promptly used in the invertebrate experiment. Three sets of five leaf discs were used for respirometry trials and then frozen at -20 °C, lyophilised, weighed (± 0.01 mg) and used for ergosterol extraction. Three sets of 10 leaf discs were

used for determination of DM, ground and used for determinations of phosphorus (P), N and C concentrations. Four microcosms (i.e. two microcosms per thickness level) were kept sterile and were used as controls for abiotic decomposition.

Mass loss and leaf physical–chemical characteristics. Specific leaf disc area (SLA) was determined for 96 discs as the area-to-mass ratio, and results were expressed as $\text{mm}^2 \text{mg}^{-1}$. Toughness was determined for the same leaf discs with a penetrometer after discs had been soaked in distilled water for 1 h, with results expressed as the mass (g) required to force an iron rod ($1.89 \text{ mm}^2 \varnothing$) through the discs (Graça, Bärlocher & Gessner, 2005).

To characterise initial litter chemical composition before and after autoclaving, three sets of 10 air-dried and autoclaved litter discs were oven-dried (105°C , 24 h) to allow estimation of initial DM and ground with a mortar and pestle (0.5 mm powder size) and subsamples weighed ($\pm 0.01 \text{ mg}$) to be analysed for C, N [IRMS Thermo Delta V advantage with a Flash EA (1112 series)], and P concentrations (APHA, 1998). Results were expressed as percentage of dry mass (% DM). After the end of the experiment, three sets of 10 leaf discs from each microcosm were oven-dried (105°C , 24 h), weighed ($\pm 0.01 \text{ mg}$) to determine remaining dry mass (DMr) calculated as the difference between initial mass and final mass, and used to estimate leaf litter elemental quality after conditioning.

Fungal sporulation. Each of the seven times the nutrient solutions were changed, 40 mL of the suspension from each microcosm was sampled, stored in 50-mL Falcon tubes and preserved with 2 mL 37% formalin until processed. When preparing filters for conidial counting and identification, 150 μL of polyethylene glycol p-(1,1,3,3-tetramethylbutyl)-phenyl ether (Triton X-100, 0.5%) was added to the suspension and mixed with a magnetic stirring bar to ensure a uniform distribution of conidia, and an aliquot of the suspension was filtered (25 mm \varnothing , pore size 5 μm , Millipore SMWP; Millipore Corp., Bedford, MA, U.S.A.). Filters were stained with 0.05% trypan blue in 60% lactic acid, and spores were identified and counted under a compound microscope at $320\times$ (Leitz Diaplan, Wetzlar, Germany) according to Graça *et al.* (2005). Cumulative conidial production over time was calculated by summing the numbers of conidia produced at the current and preceding sampling dates. The total conidial mass was calculated by multiplying the number of conidia from each species by the average mass of individual conidia obtained from the literature (Chauvet &

Suberkropp, 1998) or calculated from biovolume data (Bärlocher & Schweizer, 1983). Results were expressed as μg of conidia released mg^{-1} initial leaf DM.

Fungal respiration. At the end of the experiment, three subsets of five discs from each microcosm were used to determine fungal oxygen consumption rates using a closed six-channel dissolved oxygen measuring system (Strathkelvin 929 System, North Lanarkshire, U.K.) connected to a computer. The oxygen electrodes were calibrated against a saturated solution of sodium sulphite in 0.01 M sodium borate (0% O_2) and a 100% O_2 -saturated nutrient solution at 15°C . Leaf discs were incubated in 3-mL chambers filled with a sterile 100% O_2 -saturated nutrient solution, stirred with a magnetic bar, and kept at 15°C by circulation of water originating from a temperature-controlled water bath.

Additional chambers without leaf discs were used as controls. After a 1-h trial, leaf discs were enclosed in small sterile zip lock bags and promptly frozen at -20°C for later DM determination and ergosterol extraction. Oxygen consumption rates were determined by the difference in the O_2 concentration in the sample and the control (chamber with no leaf discs) over a 20-min interval during which O_2 consumption over time was linear, corrected for chamber volume, time and disc mass. Results were expressed as $\text{mg O}_2 \text{g}^{-1} \text{leaf DM h}^{-1}$.

Mycelial biomass. The three subsets of five leaf discs from each microcosm, frozen after the respirometry trial, were lyophilised, promptly weighed ($\pm 0.01 \text{ mg}$) to determine DM, and used for ergosterol extraction as a surrogate for mycelial biomass (Gessner & Chauvet, 1993; Graça *et al.*, 2005). Lipid extraction and saponification were carried out in 10 mL KOH/methanol (8 g L^{-1}) at 80°C for 30 min. The extract was then purified by solid phase extraction (Waters Oasis HLB 3-cc cartridges; Waters Corp., Milford, MA, U.S.A.; Graça *et al.*, 2005) and quantified with high-performance liquid chromatography (HPLC; Dionex DX-120, Sunnyvale, CA, U.S.A.) by measuring absorbance at 282 nm. The HPLC system was equipped with a LiChroCART 250-4 LiChrospher 100 RP-18 (5 μm) column (Merck, Darmstadt, Germany) maintained at 33°C . The mobile phase was 100% methanol, and the flow rate was set to 1.4 mL min^{-1} . Fungal biomass was estimated from ergosterol using either specific ergosterol–fungal DM conversion factors (*Articulospora*, 4.2 mg ergosterol g^{-1} fungal DM; *Anguillospora*, 10.9; *Clavariopsis*, 4.6; *Tetracladium*, 4.6; *Tetrachaelum*, 4.7) or a general 5.5 mg ergosterol g^{-1} fungal DM

conversion factor (*Tricladium*; no specific conversion factors were available) (Gessner & Chauvet, 1993; Suberkropp, Gessner & Chauvet, 1993), applied to the relative contribution of each species to total ergosterol mass based on specific relative contribution to total reproductive output. Results were expressed as mg mycelial biomass g^{-1} leaf DM.

Consumption test. Here, we tested whether changes in the composition of aquatic hyphomycete assemblages, identity of the dominant species and leaf thickness would influence the nutrient concentration of decomposing leaf litter. We also evaluated whether this variation in nutrient concentrations affected the feeding activity of shredders by offering leaf discs inoculated in the 20 microcosms considered above (two thickness levels \times two assemblage types \times four dominance types + four sterile controls) to the detritivore *Schizopelex festiva*.

Similar-sized, early-instar larvae were collected from depositional areas in Ribeira de Múceres, Serra do Carumulo (central Portugal; 40°32'01" N, 08°09'15" W; altitude 210 m a.s.l.), during winter 2010, and transported to the laboratory in plastic containers with stream water and sand. Individuals were kept in plastic containers with stream sand and aerated stream water, fed *ad libitum* chestnut litter conditioned in the stream for 2 weeks, and acclimated at 15 °C under a 12-h light:12-h dark photoperiod for 2 weeks before being used. The diameter of the case opening (CO, mm) of each individual was measured under a stereoscopic microscope at 16 \times (Wild M38, Heerbrugg, Switzerland), at the beginning of the experiment, and individual DM (g) was estimated by the application of the regression model $\text{DM} = 0.0002 \times \text{CO}^{3.2068}$ ($r^2 = 0.92$, $P < 10^{-3}$, $n = 30$). For the feeding experiment, the invertebrates were allocated to plastic containers (7 cm diameter \times 8.5 cm high), with a standard amount (two tea spoons; ~19 g) of ashed stream sediment (<0.5 mm; 550 °C, 8 h) covering the bottom, and 200 mL of filtered stream water (47 mm \varnothing , pore size 0.7 μm , Millipore APFF04700; Millipore Corp.), aerated for the duration of the experiment (Graça *et al.*, 2005). One pre-measured *S. festiva* larva was allocated to each of 300 plastic containers (20 treatments \times 15 replicates) and allowed access to three conditioned leaf discs, while three other leaf discs were enclosed in a fine mesh bag that hung from the plastic containers' edge and served as a control for mass loss due to leaching and microbial activity. After 48 h, when remaining litter in most containers was <50% of initial disc area, the remaining litter mass was recovered, oven-dried (105 °C; 24 h) and weighed (± 0.01 mg). The litter

DM eaten by the larva during the elapsed time (L_e , g) was calculated as $C_F - E_F$, where C_F is the final DM (g) of control litter discs and E_F is the final DM (g) of litter exposed to the larva (this assumes that control and exposed litter had the same initial mass after the conditioning step). Relative consumption rates (g leaf DM g^{-1} individual DM h^{-1}) were calculated as $L_e / (\text{DM}_{\text{II}} \times t)$, where DM_{II} is the invertebrate initial DM (g) and t is the time elapsed (h).

Data analysis. Phosphorus and N concentrations of initial and autoclaved oak leaf discs, and initial SLA, were compared between thin and thick discs by t -tests. A three-way factorial ANOVA was used to assess differences in mycelial biomass, respiration rate and percentage of leaf mass loss at the end of the conditioning period and relative consumption rate of *S. festiva* larvae among treatments [leaf thickness (i.e. Thin or Thick), assemblage type (i.e. Early or Late) and dominance treatment as the main effects] (Zar, 1996). When significant differences were detected among treatments, Tukey's HSD tests were then carried out for *post hoc* pairwise comparisons. No statistical analysis was performed on total conidial production and conidial biomass production because there was only one value per microcosm. The relationships between the specific relative conidial production and conidial biomass production after 2 weeks of incubation and the specific relative conidial abundance in the inocula were assessed across treatments within each assemblage type by linear regression. The relationship between mycelial biomass (expressed as log-transformed data) across dominance treatments and relative consumption rate of *S. festiva* larvae was assessed using simple linear regression analysis. Similarly, relationships between mycelial biomass across dominance treatments and relative consumption rate of *S. festiva* larvae and elemental composition ratios of leaf litter were examined with simple linear regression analyses. Data were log-transformed to improve homoscedasticity when necessary. All analyses were performed with STATISTICA 6 software (StatSoft Inc., 2001). For all analyses, a significance threshold of $P = 0.05$ was used.

Results

Initial litter chemical composition, toughness and specific leaf area

The thinnest leaf discs (mass classes 3.30–3.40, 3.40–3.50; and 3.50–3.60, 3.60–3.70 mg) had on average 10.6%

higher initial P concentration than leaf discs exhibiting the highest thickness (mass classes 3.90–4.00, 4.00–4.10 mg; and 4.10–4.20, 4.20–4.30 mg), although the difference was not significant (Table 1; $P = 0.23$). Leaching by autoclaving affected thick and thin leaf discs differently, with greater losses in P concentration observed for the thinnest leaf discs (52.2%) compared with the thickest ones (29.6%; Table 1). Although leaching increased the relative difference in P concentration between the two sets of leaf discs to 24.9%, this still remained non-significant (Table 1, $P = 0.11$). Leaf disc thickness did not significantly affect initial N concentration ($P = 0.08$) in spite of a relative difference of 11.2% between thickness levels (Table 1). However, leaching by autoclaving significantly increased the relative difference in N concentration between the two sets of leaf discs to 19.7% (Table 1; $P = 0.01$). Carbon concentrations before and after autoclaving were similar between thin and thick

leaves (Table 1, <0.3% difference; $P = 0.38$ and 0.55 , respectively). Based on a set of leaf discs randomly chosen, disc toughness was negatively related to SLA (linear regression, $P < 0.001$, $r^2 = 0.27$, $n = 96$, leaf disc mass from 2.2 to 6.7 mg or leaf disc SLA from 51.88 to 16.96 mm² mg⁻¹). However, the relationship remained weak in spite of a large difference between the extreme leaf disc mass or SLA values.

Aquatic hyphomycete assemblages. Overall, conidial biomass production from early assemblages was consistently higher than that from late assemblages, whatever the disc thickness (i.e. 1.4 and 1.5 times higher in the thin and thick discs, respectively; Table 2). Manipulation of the specific number of conidia inoculated into the microcosms allowed us to anticipate the identity of the dominant species in the *Anguillospora* (Early assemblage), *Clavariopsis* (Late assemblage) and *Tetracladium*

Table 1 Phosphorus (P), nitrogen (N) and carbon (C) concentrations (mean \pm 1 SE) of initial and autoclaved thin and thick oak leaf discs. Grey-shaded text shows early assemblages and unshaded late assemblages

Thickness	Litter state	P (% DM)	N (% DM)	C (% DM)
Thin	Initial non autoclaved	0.027 \pm 0.000	0.86 \pm 0.01	46.72 \pm 0.09
	Initial autoclaved	0.011 \pm 0.002	0.87 \pm 0.07	47.44 \pm 0.15
	Initial non autoclaved	0.030 \pm 0.003	0.93 \pm 0.06	47.24 \pm 0.14
	Initial autoclaved	0.017 \pm 0.003	1.04 \pm 0.07	48.07 \pm 0.24
Thick	Initial non autoclaved	0.024 \pm 0.002	0.96 \pm 0.07	47.15 \pm 0.01
	Initial autoclaved	0.017 \pm 0.003	1.18 \pm 0.02	47.77 \pm 0.11
	Initial non autoclaved	0.028 \pm 0.001	1.06 \pm 0.07	47.11 \pm 0.20
	Initial autoclaved	0.019 \pm 0.001	1.20 \pm 0.10	47.48 \pm 0.12

Table 2 Total production of conidial biomass and relative contribution of the leaf-associated fungal species to the total production of conidial biomass at two leaf thickness levels and two assemblage types for the four dominance types over 2 weeks. Grey-shaded text shows early assemblages and unshaded late assemblages. Fungal species: *Anguillospora filiformis* (*Anguillospora*), *Articulospora tetracladia* (*Articulospora*), *Clavariopsis aquatica* (*Clavariopsis*), *Tetrachaetum elegans* (*Tetrachaetum*), *Tetracladium marchalianum* (*Tetracladium*) and *Tricladium chaetocladium* (*Tricladium*)

Thickness	Dominance type	Total conidial production ($\mu\text{g mg}^{-1}$ DMi)	<i>Anguillospora</i> (%)	<i>Tetrachaetum</i> (%)	<i>Tricladium</i> (%)	<i>Articulospora</i> (%)	<i>Clavariopsis</i> (%)	<i>Tetracladium</i> (%)
Thin	Even	9.71	47	43	9	–	–	–
	<i>Anguillospora</i>	8.78	52	41	7	–	–	–
	<i>Tetrachaetum</i>	8.88	52	43	5	–	–	–
	<i>Tricladium</i>	9.46	46	33	21	–	–	–
	Even	5.02	–	–	–	12	38	51
	<i>Articulospora</i>	6.01	–	–	–	7	41	52
	<i>Clavariopsis</i>	8.21	–	–	–	5	54	41
	<i>Tetracladium</i>	7.05	–	–	–	3	27	70
Thick	Even	9.20	61	32	7	–	–	–
	<i>Anguillospora</i>	10.62	67	27	7	–	–	–
	<i>Tetrachaetum</i>	7.99	54	40	5	–	–	–
	<i>Tricladium</i>	8.90	49	27	23	–	–	–
	Even	4.58	–	–	–	10	38	52
	<i>Articulospora</i>	6.13	–	–	–	9	37	54
	<i>Clavariopsis</i>	8.29	–	–	–	5	50	45
	<i>Tetracladium</i>	4.74	–	–	–	8	9	83

(Late assemblage) treatments for both thickness levels. Early assemblages were dominated by *Anguillospora* (Table 2), even when this species was represented by only 20% conidia in the initial inocula (see Methods). Conidial biomass production by *Tetrachaetum* across treatments was also not related to its abundance in the inocula ($r^2 = 0.37$; $P = 0.11$), and *Tetrachaetum* generally codominated with *Anguillospora* (Table 2). *Tricladium* conidial production across treatments was related to its initial abundance ($r^2 = 0.93$; $P < 0.001$), but was the species contributing the least to the total conidial pool within each treatment. Late assemblages were dominated by *Tetracladium*, except for *Clavariopsis* treatments where *Clavariopsis* dominated. Conidial biomass production by *Tetracladium* across treatments was related to its abundance in the inocula ($r^2 = 0.81$; $P < 0.01$). *Articulospora* was consistently outcompeted by the other two species, even in the *Articulospora* treatments. Nevertheless, each of the three species tended to produce relatively more conidial biomass in the microcosms where they were inoculated with the highest number of conidia (Table 2).

Litter mass loss. Mass loss of oak leaf discs after incubation in microcosms for 2 weeks varied between 13.2% (Thin, Early assemblage, Even) and 49.7% (Thin, Late assemblage, *Tetracladium*). Interestingly, leaf decomposition by late assemblages was much faster than that pro-

motivated by early assemblages, in both leaf thickness levels ($P < 0.001$; Tables 3 & 4). Decomposition rate was affected by species dominance identity within thickness level (i.e. Thin and Thick; $P = 0.006$) but also by dominant species identity within assemblage type (i.e. Early versus Late; $P < 0.001$; Tables 3 & 4). Mass loss was negatively affected by litter thickness ($P < 0.001$, Tables 3 & 4).

Fungal biomass, respiration and leaf litter quality for consumers. After 2-week conditioning, fungal biomass on leaf discs varied between 83 (Thin, Early assemblage, *Anguillospora*) and 389 mg g⁻¹ leaf DM (Thin, Late assemblage, *Articulospora*) (Table 3). Higher fungal biomass accrual (~three times) was obtained in microcosms where leaf discs were inoculated with late assemblages ($P < 0.001$; Tables 3 & 4), with the difference being more pronounced between assemblage types in the thin than in the thick discs ($P < 0.001$; Tables 3 & 4). Interestingly, the increase in disc thickness led to a significant increase in fungal biomass in microcosms with early assemblages, whereas it resulted in a decrease in fungal biomass with late assemblages ($P < 0.001$; Table 4). Mycelial biomass accrual was affected by changes in identity of the dominant species: thin discs in microcosms inoculated with *Articulospora* as the dominant species had higher values than thin discs in microcosms inoculated with *Anguillospora* or *Tetrachaetum* as the dominant species (Table 3).

Table 3 Mycelial biomass, fungal activity, dry mass remaining and elemental composition (mean \pm 1 SE) of oak leaf discs at two leaf thickness levels and two assemblage types for the four dominance types over 2 weeks. Grey-shaded text shows early assemblages and unshaded late assemblages. Treatments with the same letter are not significantly different (Tukey's HSD, $P > 0.05$). Fungal species: *Anguillospora filiformis* (*Anguillospora*), *Articulospora tetracladia* (*Articulospora*), *Clavariopsis aquatica* (*Clavariopsis*), *Tetrachaetum elegans* (*Tetrachaetum*), *Tetracladium marchalianum* (*Tetracladium*) and *Tricladium chaetocladium* (*Tricladium*)

Thickness	Dominance type	Mycelial biomass (mg g ⁻¹ leaf DM)	Fungal activity (mg O ₂ g ⁻¹ DM h ⁻¹)	DMr (%)	P (% DM)	N (% DM)	C (% DM)
Thin	Even	101 \pm 9a	0.71 \pm 0.04a	86.76 \pm 0.56e	0.106 \pm 0.006abc	1.68 \pm 0.02ab	48.40 \pm 0.09a
	<i>Anguillospora</i>	83 \pm 9a	0.82 \pm 0.07ab	81.89 \pm 1.02de	0.097 \pm 0.007ab	1.84 \pm 0.06ab	47.99 \pm 0.39a
	<i>Tetrachaetum</i>	86 \pm 3a	0.86 \pm 0.02abc	80.95 \pm 1.69de	0.093 \pm 0.000a	1.72 \pm 0.11ab	46.86 \pm 0.95a
	<i>Tricladium</i>	104 \pm 9a	1.09 \pm 0.10abcde	80.54 \pm 1.71d	0.138 \pm 0.007efg	1.89 \pm 0.06ab	50.30 \pm 2.52a
	Even	296 \pm 5def	1.36 \pm 0.07cdef	50.95 \pm 0.38a	0.135 \pm 0.004efg	1.95 \pm 0.07ab	47.64 \pm 0.13a
	<i>Articulospora</i>	389 \pm 21g	1.82 \pm 0.16f	53.43 \pm 2.10ab	0.155 \pm 0.002g	1.87 \pm 0.03ab	48.23 \pm 0.09a
	<i>Clavariopsis</i>	346 \pm 15fg	1.51 \pm 0.13ef	52.39 \pm 0.83a	0.139 \pm 0.003efg	1.46 \pm 0.32a	40.42 \pm 5.99a
	<i>Tetracladium</i>	318 \pm 10ef	1.41 \pm 0.11def	50.31 \pm 0.78a	0.132 \pm 0.010defg	1.93 \pm 0.12ab	48.30 \pm 0.46a
Thick	Even	88 \pm 15a	1.01 \pm 0.07abcde	77.81 \pm 0.36d	0.108 \pm 0.002abcd	1.47 \pm 0.19a	47.68 \pm 0.72a
	<i>Anguillospora</i>	114 \pm 9a	1.01 \pm 0.02abcde	76.82 \pm 0.34d	0.106 \pm 0.003abc	1.80 \pm 0.12ab	48.28 \pm 0.37a
	<i>Tetrachaetum</i>	108 \pm 3a	0.94 \pm 0.04abcd	80.91 \pm 0.68de	0.095 \pm 0.004ab	1.50 \pm 0.23a	44.51 \pm 3.95a
	<i>Tricladium</i>	134 \pm 4a	1.14 \pm 0.09abcde	81.50 \pm 2.08de	0.124 \pm 0.004cdef	1.93 \pm 0.06ab	48.15 \pm 0.30a
	Even	284 \pm 7cde	1.23 \pm 0.03bcde	59.07 \pm 0.27bc	0.138 \pm 0.005efg	1.78 \pm 0.17ab	48.03 \pm 0.24a
	<i>Articulospora</i>	238 \pm 5bc	1.82 \pm 0.21f	65.12 \pm 0.40c	0.145 \pm 0.005fg	2.24 \pm 0.19b	50.45 \pm 2.84a
	<i>Clavariopsis</i>	261 \pm 7bcd	1.13 \pm 0.09abcde	63.79 \pm 1.69c	0.119 \pm 0.005abcde	2.07 \pm 0.11ab	47.51 \pm 0.88a
	<i>Tetracladium</i>	228 \pm 18b	1.22 \pm 0.10abcde	52.31 \pm 0.93a	0.120 \pm 0.002bcdef	1.83 \pm 0.02ab	47.94 \pm 0.24a

Table 4 Summary table for three-way ANOVAs performed on mycelial biomass and fungal respiration associated with oak leaf discs, litter mass loss over 2 weeks and relative consumption rate of *Schizopelex festiva* caddisfly larvae

	d.f.	Mycelial biomass		Fungal respiration		Litter mass loss		Relative consumption rate of <i>S. festiva</i>	
		F	P	F	P	F	P	F	P
Intercept	1	4261.5	<0.001	2374.0	<0.001	54557.6	<0.001	3477.0	<0.001
Thickness level (Thickness)	1	0.5	0.496	0.0	0.841	18.4	<0.001	4.1	0.043
Assemblage type (Assemblage)	1	327.0	<0.001	99.5	<0.001	1818.3	<0.001	255.4	<0.001
Dominance type (Dominance)	3	1.4	0.251	7.0	0.001	6.9	0.001	8.6	<0.001
Thickness × Assemblage	1	160.2	<0.001	11.4	0.002	97.6	<0.001	7.4	0.007
Thickness × Dominance	3	0.8	0.499	1.5	0.232	4.9	0.006	1.3	0.277
Assemblage × Dominance	3	4.9	0.007	9.3	<0.001	13.7	<0.001	6.4	<0.001
Thickness × Assemblage × Dominance	3	10.4	<0.001	0.5	0.716	10.2	<0.001	1.4	0.240
Error	32								

Fungal respiration after incubation for 2 weeks was strongly affected by changes in identity of the dominant species in the inoculum ($P = 0.001$; Table 4) and varied between $0.71 \text{ mg O}_2 \text{ g}^{-1} \text{ DM h}^{-1}$ (Thin, Early assemblage, Even) and $1.82 \text{ mg O}_2 \text{ g}^{-1} \text{ DM h}^{-1}$ (Thin, Late assemblage, *Articulospora* and Thick, Late assemblage, *Articulospora*). Overall, leaf litter thickness did not affect respiration rate ($P = 0.84$; Tables 3 & 4), and higher values for fungal respiration were consistently obtained in microcosms inoculated with late assemblages ($P < 0.001$; Table 3). As previously shown for fungal biomass, the increase of leaf litter thickness led to a significantly higher fungal respiration in microcosms with early assemblages, conversely to late assemblages ($P = 0.002$; Table 4). Fungal respiration showed a significant posi-


tive correlation with mycelial biomass ($r^2 = 0.66$; $P < 0.001$).

Leaf litter conditioned by different fungal assemblages resulted in different N and P concentrations on leaves, with a tendency for higher concentrations when leaf discs were inoculated with the late assemblage ($P < 0.001$ for P, $P = 0.03$ for N).

Consumption of leaf discs by a stream detritivore

Larval mortality during the short-term feeding assay was very low (1.0%). Larval consumption rate averaged $0.071 \text{ g leaf DM g}^{-1} \text{ individual DM h}^{-1}$ across treatments, but there were clear differences between disc thickness levels, fungal assemblages and dominance

Fig. 3 Relative consumption rate (RCR) of *Schizopelex festiva* caddisfly larvae for thin and thick oak leaf discs, for each fungal assemblage type and species dominance treatment. Grey bars represent early assemblages, and open bars represent late assemblages. Treatments within each thickness level were compared with three-way ANOVA, followed by Tukey's HSD; different letters indicate significant differences among treatments ($P < 0.05$). Fungal species: *Anguillospora filiformis* (*Anguillospora*), *Articulospora tetracladia* (*Articulospora*), *Clavariopsis aquatica* (*Clavariopsis*), *Tetrachaetum elegans* (*Tetrachaetum*), *Tetracladium marchalianum* (*Tetracladium*) and *Tricladium chaetocladium* (*Tricladium*).



types (Table 4). The relative consumption rate of caddisfly larvae was two-three times higher on late than on early assemblages for both thickness levels. Interestingly, for both thin and thick discs, species dominance identity was the prime factor affecting larval consumption within early but not within late assemblages (Fig. 3). In parallel, fungal biomass ($r^2 = 0.71$; $P < 0.001$) and fungal activity measured as oxygen consumption ($r^2 = 0.62$; $P < 0.001$) consistently explained the relative consumption rate of caddisfly larvae (Figs 3 & 4; Table 3).

Relationship between leaf litter quality and invertebrate consumption rate

There was a strong and positive relationship between relative consumption rate of *S. festiva* larvae and mycelial biomass across all treatments ($r^2 = 0.90$, $P < 0.001$; Fig. 4). Caddisfly larvae consumption rate and mycelial biomass were also related to litter P concentration ($r^2 = 0.63$, $P < 0.001$ and $r^2 = 0.63$, $P < 0.001$, respectively), with P concentration differing between thickness and fungal assemblage treatments by 40% at most. In contrast, caddisfly larvae consumption rate and mycelial biomass were not significantly related to litter N concentration ($r^2 = 0.23$, $P = 0.06$ and $r^2 = 0.07$, $P = 0.31$, respectively), even though N concentration differed between thickness and fungal assemblage treatments by




Fig. 4 Relative consumption rate (RCR) of *Schizopelex festiva* caddisfly larvae as a function of mycelial biomass. Grey symbols represent early assemblages, and open circles represent late assemblages. Line and corresponding r^2 - and P -values are shown for simple linear regression.

as much as 35%. Caddisfly larvae consumption rate and mycelial biomass were strongly related to C/P leaf litter ratio ($r^2 = 0.71$, $P < 0.001$ and $r^2 = 0.72$, $P < 0.001$, respectively; Figs 5a, 6a). In contrast, although being significantly related to caddisfly larvae consumption rate ($r^2 = 0.39$, $P = 0.01$; Fig. 5b), C/N leaf litter ratio was not related to mycelial biomass ($r^2 = 0.22$, $P = 0.07$; Fig. 6b). Caddisfly larvae consumption rate and mycelial biomass were weakly related to N/P leaf litter ratio ($r^2 = 0.26$, $P = 0.04$ and $r^2 = 0.38$, $P < 0.01$, respectively; Figs 5c, 6c).

Discussion

Changes in community structure as an early outcome of anthropogenic induced stresses are expected to precede a reduction in species richness (Ellison *et al.*, 2005; Hillebrand, Bennett & Cadotte, 2008). Recent laboratory experiments and field studies have shown that changes in dissolved nutrient concentration and temperature induce shifts in fungal species abundance (Dang *et al.*, 2005; Artigas, Romani & Sabater, 2008; Ferreira & Chauvet, 2011). While these studies have suggested that changes in abundance of particular species could strongly alter nutrient transfer efficiency in food webs, the relationship between leaf litter elemental composition and changes in associated fungal community structure has not been investigated. Thus, it is relevant to assess whether these changes have the potential to affect ecosystem processes. Here, we show how changes in decomposing litter quality, due solely to differences in the species composition of the fungal assemblage, can affect the feeding activity of a stream detritivore, which might ultimately affect its growth and survival (Arsuffi & Suberkropp, 1986; Chung & Suberkropp, 2009). Our data suggest that changes in assemblage structure have a strong influence on the feeding activity of *S. festiva* caddisfly larvae through changes in fungal biomass accumulation and on the elemental composition of its resource.

Manipulation of the relative abundances of fungal species inoculated did not determine the identity of the dominant species in all fungal assemblages, but only in six of 16. Nevertheless, we obtained sixteen different fungal assemblages that differed in the relative proportion of each species. The strong discrepancies between the specific relative conidial production over the incubation time and that initially present in the inoculum might be attributed to differences in conidial traits (e.g. shape and size) likely to influence the probability of their settlement and germination (Dang, Gessner &




Fig. 5 Relative consumption rate (RCR) of *Schizopelex festiva* cadisfly larvae as a function of leaf litter (a) C/P, (b) C/N and (c) N/P molar ratios. Grey symbols represent early assemblages, and open circles represent late assemblages. Lines and corresponding r^2 - and P -values are shown for simple linear regression.




Fig. 6 Leaf litter (a) C/P, (b) C/N and (c) N/P molar ratios as a function of mycelial biomass. Grey symbols represent early assemblages, and open circles represent late assemblages. Lines and corresponding r^2 - and P -values are shown for simple linear regression.

Chauvet, 2007; Kearns & Bärlocher, 2008). Moreover, aquatic hyphomycete assemblage structure in our study was based on specific reproductive output, which might not necessarily reflect specific mycelial biomass in the

leaf litter. A recurrent and unique problem with microbial systems is the difficulty of keeping track of individual species. Recent advances in molecular approaches, although prone to uncertainty, can circumvent such obstacles by allowing the quantification of species contribution to total fungal biomass (Bärlocher, 2007, 2010; Krauss *et al.*, 2011).

Our study contrasts with other studies suggesting functional redundancy among aquatic hyphomycete assemblages (Bärlocher & Graça, 2002; Pascoal, Cássio & Marvanová, 2005; Ferreira, Gulis & Graça, 2006b; Ferreira & Chauvet, 2012). Conversely, they agree with studies that have demonstrated the important role played by the identity of dominant species in the relationship between community evenness and ecosystem processes (McKie *et al.*, 2008; Gonçalves & Canhoto, 2009; Swan, Gluth & Horne, 2009).

Thereby, our results demonstrated that experimental changes in fungal community structure [i.e. assemblage type (Early versus Late), species identity and evenness] resulted in an increased nutritional quality of leaf litter which, in turn, translated into an increased proportion of leaf litter consumed by a detritivore. Changes in fungal community structure associated with leaf litter can alter the nutritional quality of leaf detritus via a number of mechanisms that are not mutually exclusive and that may act synergistically.

First, changes in fungal species composition can translate into an increased biomass and activity of fungi associated with leaf detritus, which are often differentially ingested by invertebrate detritivores (Suberkropp, 1992) or assimilated at a higher efficiency than non-living detritus (Graça, Maltby & Calow, 1993a). It is widely accepted that the degree of conditioning (usually defined by measures of microbial biomass or activity) is often the main factor affecting resource quality and constitutes the standard by which all types of shredders discriminate among leaves. The diversity of feeding patterns exhibited by shredders reported by numerous studies (e.g. Suberkropp *et al.*, 1983; Suberkropp & Arsuffi, 1984; Butler & Suberkropp, 1986; Arsuffi & Suberkropp, 1989; Friberg & Jacobsen, 1994), with a few others reporting ambiguous or less well-defined patterns, suggests that this factor alone does not determine foraging behaviour. For example, there was no correlation between palatability and protein increase or mass loss (as indicator of overall enzymatic activity; Bärlocher & Kendrick, 1973). Graça, Maltby & Calow (1993b, 1994) reported that no leaf parameter was correlated with resource preference of *Gammarus pulex* in experiments where leaf discs were conditioned by different pure cul-

tures of individual fungal species. Similarly, consumption by *Gammarus tigrinus* of leaf discs conditioned by six pure cultures of individual aquatic hyphomycete species was not correlated with leaf protein, phenols, lipids or ergosterol concentrations (Rong, Sridhar & Bärlocher, 1995).

Second, heterotrophic microorganisms contain high concentrations of nutrients (Sterner & Elser, 2002; Cross *et al.*, 2005), which may directly affect the growth and production of consumers (Iversen, 1974; Urabe, Clasen & Sterner, 1997). Leaf litter nutrient concentration has been shown to increase shredder consumption rates in headwater streams (Iversen, 1974; Irons, Oswood & Bryant, 1988).

Regression analysis in our study indicated that leaf litter quality was affected by the amount of fungal biomass associated with the leaf discs. There was quite strong variation in fungal biomass, depending on different fungal assemblages (e.g. 4.7 times between the two extreme values). In line with previous studies, fungal colonisation led to nutrient immobilisation (Gulis & Suberkropp, 2003), which in our case translated into lower C/P ratios, without significantly affecting C/N ratios.

Interestingly, several studies that documented foraging preferences of caddisfly larvae (i.e. three families, five genera; Suberkropp *et al.*, 1983; Arsuffi & Suberkropp, 1984; Suberkropp & Arsuffi, 1984; Arsuffi & Suberkropp, 1986) reported that all caddisfly species examined had similar preferences when offered leaf litter colonised by the same aquatic hyphomycete species. This similarity among trichopteran contrasts strongly with different ranked responses of different groups of detritivores (i.e. Amphipoda, Plecoptera, Diptera and Gastropoda) to leaves colonised by various fungi (Arsuffi & Suberkropp, 1989), and our findings suggest that stoichiometric imbalances between consumers and resources might affect consumer feeding activity.

Heterotrophic microorganisms also contain biochemicals, such as polyunsaturated fatty acids (PUFA) (Brett & Mueller-Navarra, 1997; Arce Funck *et al.*, 2015), which are almost exclusively synthesised by autotrophic organisms. Animals can convert from one form of PUFA to another through elongation and desaturation, but very few can synthesise PUFA *de novo*. Accordingly, most consumers depend on PUFA supplied through their dietary intakes to meet their physiological requirements (Arts, Brett & Kainz, 2009). Polyunsaturated fatty acids are essential for invertebrates owing to their involvement in a wide range of physiological processes, such as regulation of cell membrane properties or as precursors to many hormones (Brett & Mueller-Navarra, 1997).

Despite the pivotal role played by aquatic hyphomycetes as the main basal food sources in stream food webs, data on their lipid composition are extremely scarce (but see Cargill *et al.*, 1985 and Arce Funck *et al.*, 2015). Accordingly, further investigation in this direction might deepen our understanding of the role of aquatic hyphomycetes in forested headwater stream ecosystems.

Lastly, changes in dominance/evenness and/or identity of species among assemblages are likely to result in qualitative differences in enzymatic capabilities among species belonging to early- or late-successional stages and may alter leaf detritus degradation and, in turn, modulate the digestibility of detritus to consumers (Bärlocher, 1985; Sinsabaugh, Linkins & Benfield, 1985; Suberkropp, 1992). Based on a study by Güsewell & Gessner (2009), and given the N/P ratio of the nutrient solution adopted in our study (31.3 on a molar basis) and the overall nutrient supply used in our microcosms, we assume that the fungal assemblages were likely to be N-limited, constraining them to maintain N content in their own biomass at a basal metabolic level. This was apparent from the absence of a significant relationship between C/N leaf litter ratio and mycelial biomass. In contrast, P limitation for fungi is expected to be reached at higher N/P ratios than that used here (Güsewell & Gessner, 2009). Accordingly, the negative relationship between C/P leaf litter ratio and mycelial biomass among treatments was mainly due to P immobilisation by fungi, which might be related to an increase in fungal decomposition activity and biomass. In addition, fungi growing in P-unlimited conditions, as was the case here, might store P in excess in their own biomass, mainly in membranes or in vacuoles (Beever & Burns, 1980), thereby reducing C/P leaf litter ratios. It is important, however, to note that these results have been obtained with assemblages of heterotrophic decomposer assemblages solely composed of fungi. Hodge, Robinson & Fitter (2000) proposed that bacteria should dominate decomposition processes on substrates with low C/N ratios, while the contribution of fungi should prevail on substrates with high C/N ratios, due to the contrasting N requirements of these organisms (but see Hieber & Gessner, 2002). Similarly, fungi are known to have lower nutrient requirements and have lower metabolic activity than bacteria (Hieber & Gessner, 2002; Cross *et al.*, 2005). Accordingly, the relative P requirements of fungi should be lower than those of bacteria (Smith, 2002), so that fungi are expected to become P-limited at higher N/P ratios than bacteria. Consequently, it is likely that the presence of bacteria in our experiment altered the observed patterns regarding the elemental composition

of leaf detritus, potentially leading to changes in the feeding activity of shredder-mediated litter decomposition.

Relatively few studies have explored stoichiometric detritivore–resource relationships in running waters (Cross *et al.*, 2003, 2005; Hladyz *et al.*, 2009; Danger *et al.*, 2012, 2013; Lauridsen *et al.*, 2012, 2014). In this study, litter C/P was a better predictor of caddisfly larvae feeding performance than C/N or N/P, suggesting that P rather than N might have been limiting for these consumers. Accordingly, our results differ from those of Ferreira *et al.* (2010), which indicated that invertebrates prefer litter with higher N than P concentration, and also with the notion that litter N concentration generally determines invertebrate litter choice because it is often the limiting nutrient for primary consumers (Evans-White, Stelzer & Lamberti, 2005). In contrast, our results agree with the growth rate hypothesis, which proposes that consumer growth rather tends to be limited by the resource P concentration (Elser *et al.*, 2003).

Organisms require a mixture of energy, vitamins, biochemicals and minerals to grow and reproduce. Imbalanced supply of one or more of these elements slows the growth of animals (Elser *et al.*, 2000a, 2003; Frost & Elser, 2002) and alters their physiology, life history and behaviour (Sterner & Elser, 2002; Frost *et al.*, 2005). While we did not assess the elemental body composition of our invertebrates, Cross *et al.* (2003) and Hladyz *et al.* (2009) found similar molar C/N ratios for shredders: 6.7 [mean for 11 shredder species including six genera: *Leuctra* and *Tallaperla* (Plecoptera), *Tipula* (Diptera), *Fattigia*, *Lepidostoma* and *Pycnopsyche* (Trichoptera), the mean also included multiple size classes for *Tallaperla* and *Fattigia*] and 6.7 [mean for four shredder species: *Protonemura meyeri* (Plecoptera), *Gammarus duebeni* (Amphipoda), *Halesus radiatus* (Trichoptera) and *Potamophylax cingulatus* (Trichoptera)] or 6.5 (mean for two Trichoptera species: *Halesus radiatus* and *Potamophylax cingulatus*), respectively, which suggests very narrow constraints on tissue N concentration across even distantly related and geographically separated taxa (Hladyz *et al.*, 2009). Molar C/P ratios were also quite similar in both studies [498 and 487 (values from caddisfly larvae only) on average, respectively]. In accordance with the elemental body composition values of caddisfly shredders from the literature (Cross *et al.*, 2003; Hladyz *et al.*, 2009), our results show that C/N imbalances between *S. festiva* caddisfly larvae and oak leaf litter were relatively high (i.e. range from 4.0 to 5.7 times) and, therefore, consumers were particularly limited by N. In parallel, C/P imbalances between consumers and resources were relatively low when compared to C/N (i.e. range from 1.5

to 2.6 times) and may explain why shredders responded more strongly to litter C/P rather than C/N ratio.

Meeting nutrient requirements is particularly challenging for leaf-shredding invertebrates because their food resources exhibit a dramatic imbalance from their body composition (Cross *et al.*, 2003, 2005). According to the principles of ecological stoichiometry, the nutritional requirements of a given consumer are defined by its nutrient concentration and relative growth rate (Sterner & Elser, 2002; Cross *et al.*, 2003). Consequently, consumers (or specific life stages) with high body N or P concentration and high growth rates require food that is high in N or P, to maintain optimal growth (Cross *et al.*, 2003). Here, we show that the higher sensitivity regarding food P rather than N content in *S. festiva* caddisfly larvae may be explained by its early life stage. High growth rates during early stages of development are correlated with a high content of (P-rich) ribosomal RNA and associated high body P demand (Elser *et al.*, 2000b). These results are in accordance with predictions of ecological stoichiometry and have considerable empirical support, as underlined by Cross *et al.* (2003), who suggested that differences in ontogeny, life history strategy and relative allocation of structural biomolecules may all contribute to changes regarding the nutritional requirements of a given consumer.

In conclusion, we demonstrated that changes in fungal community structure affected the elemental composition of leaves, which significantly affected the feeding performance of *S. festiva* caddisfly larvae. Stimulated (Late assemblages) or reduced (Early assemblages) feeding activities of caddisfly larvae were related to changes in aquatic hyphomycete community structure. We showed that microbial identity can regulate leaf litter decomposition. Changes in consumption rates can potentially alter growth and mortality rates, and body elemental composition, and therefore influence shredder population dynamics, which will directly affect the processes in which these organisms are involved (e.g. litter decomposition, nutrient cycling). The relationship between fungi and invertebrates under changing environments remains to be explored in further studies to identify cascading effects on aquatic food-web structure and related ecosystem processes.

Acknowledgments

This work was completed at IMAR-CMA, Department of Life Sciences, University of Coimbra with funding by the Portuguese Foundation for Science and Technology (FCT) to VF (References SFRH/BPD/34368/2006, SFRH/

BPD/76482/2011, programme POPH/FSE) and to ALG (Reference SFRH/BD/47089/2008, SFRH/BPD/94820/2013), and by the National Center for Scientific Research (CNRS) to JC, as well as by the Pessoa exchange programme between Portugal and France (PHC 22808 TB) from the French Ministry of Foreign and European Affairs. We are grateful to Michael Danger, Manuel A.S. Graça and Mark Gessner whose comments led to significant improvements of the manuscript, and Elsa Rodrigues, Cristina Docal and Ana Virgínia Lirio for their expert technical assistance. Comments provided by three anonymous referees on an early version of the manuscript were most appreciated.

Author contribution

J.C., V.F., A.L.G., E.C. and C.C. designed the experiments; J.C., V.F. and A.L.G. carried out the experiments; and J.C., V.F., A.L.G., E.C. and C.C. wrote the manuscript.

References

- Aerts R. & Chapin F.S. (2000) The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. *Advances in Ecological Research*, **30**, 1–67.
- APHA (1998) Phosphorus: automated ascorbic acid reduction method, 4500-P, F. In: *Standard Methods for the Examination of Water and Wastewater* (20th ed.), (Ed. M.A.H. Franson), pp. 148–149. American Public Health Association, Washington, DC.
- Arce Funck J., Bec A., Perrière F., Felten V. & Danger M. (2015) Aquatic hyphomycetes: a potential source of polyunsaturated fatty acids in detritus-based stream food webs. *Fungal Ecology*, **13**, 205–210.
- Arsuffi T.L. & Suberkropp K. (1984) Leaf processing capabilities of aquatic hyphomycetes: interspecific differences and influence on shredder feeding preferences. *Oikos*, **42**, 144–154.
- Arsuffi T.L. & Suberkropp K. (1986) Growth of two stream caddisflies (Trichoptera) on leaves colonized by different fungal species. *Journal of the North American Benthological Society*, **5**, 297–305.
- Arsuffi T.L. & Suberkropp K. (1989) Selective feeding by shredders on leaf-colonizing stream fungi: comparison of macroinvertebrate taxa. *Oecologia*, **79**, 30–37.
- Artigas J., Romání A.M. & Sabater S. (2008) Effect of nutrients on the sporulation and diversity of aquatic hyphomycetes on submerged substrata in a Mediterranean stream. *Aquatic Botany*, **88**, 32–38.
- Arts M.T., Brett M.T. & Kainz M.J. (2009) *Lipids in Aquatic Ecosystems*. Springer, Dordrecht.
- Bärlocher F. (1985) The role of fungi in the nutrition of stream invertebrates. *Botanical Journal of the Linnean Society*, **91**, 83–94.

- Bärlocher F. (2007) Molecular approaches applied to aquatic hyphomycetes. *Fungal Biology Reviews*, **21**, 19–24.
- Bärlocher F. (2010) Molecular approaches promise a deeper and broader understanding of the evolutionary ecology of aquatic hyphomycetes. *Journal of the North American Benthological Society*, **29**, 1027–1041.
- Bärlocher F. & Graça M.A.S. (2002) Exotic riparian vegetation lowers fungal diversity but not leaf decomposition in Portuguese streams. *Freshwater Biology*, **47**, 1123–1135.
- Bärlocher F. & Kendrick B. (1973) Fungi and food preferences of *Gammarus pseudolimnaeus*. *Archiv für Hydrobiologie*, **72**, 501–516.
- Bärlocher F. & Schweizer M. (1983) Effects of leaf size and decay rate on colonization by aquatic hyphomycetes. *Oikos*, **41**, 205–210.
- Beever R.E. & Burns D.J.W. (1980) Phosphorus uptake and utilization by fungi. *Advances in Botanical Research*, **8**, 127–219.
- Brett M.T. & Mueller-Navarra D.C. (1997) The role of highly unsaturated fatty acids in aquatic foodweb processes. *Freshwater Biology*, **38**, 483–499.
- Bruning K. (1991) Effects of phosphorus limitation on the epidemiology of a chytrid phytoplankton parasite. *Freshwater Biology*, **25**, 405–417.
- Butler N.M., Suttle C.A. & Neill W.E. (1989) Discrimination by freshwater zooplankton between single algal cells differing in nutritional status. *Oecologia*, **78**, 368–372.
- Butler S.K. & Suberkropp K. (1986) Aquatic hyphomycetes on oak leaves: comparison of growth, degradation and palatability. *Mycologia*, **78**, 922–928.
- Cargill A.S., Cummins K.W., Hanson B.J. & Lowry R.R. (1985) The role of lipids as feeding stimulants for shredding aquatic insects. *Freshwater Biology*, **15**, 455–464.
- Cebrian J. (1999) Patterns in the fate of production in plant communities. *The American Naturalist*, **154**, 449–468.
- Chauvet E. & Suberkropp K. (1998) Temperature and sporulation of aquatic hyphomycetes. *Applied and Environmental Microbiology*, **64**, 1522–1525.
- Chen Y., Williams K.A., Harbaugh B.K. & Bell M.L. (2004) Effects of tissue phosphorus and nitrogen in *Impatiens wallerana* on western flower thrips (*Frankliniella occidentalis*) population levels and plant damage. *HortScience*, **39**, 545–550.
- Chung N. & Suberkropp K. (2009) Effects of aquatic fungi on feeding preferences and bioenergetics of *Pycnopsyche gentilis* (Trichoptera: Limnephilidae). *Hydrobiologia*, **630**, 257–269.
- Cross W.F., Benstead J.P., Frost P.C. & Thomas S.A. (2005) Ecological stoichiometry in freshwater benthic systems: recent progress and perspectives. *Freshwater Biology*, **50**, 1895–1912.
- Cross W.F., Benstead J.P., Rosemond A.D. & Wallace J.B. (2003) Consumer-resource stoichiometry in detritus-based streams. *Ecology Letters*, **6**, 721–732.
- Dang C.K., Chauvet E. & Gessner M.O. (2005) Magnitude and variability of process rates in fungal diversity-litter decomposition relationships. *Ecology Letters*, **8**, 1129–1137.
- Dang C.K., Gessner M.O. & Chauvet E. (2007) Influence of conidial traits and leaf surface structure on attachment success of aquatic hyphomycetes on leaf litter. *Mycologia*, **99**, 24–32.
- Danger M., Arce Funck J., Devin S., Heberle J. & Felten V. (2013) Phosphorus content in detritus controls life-history traits of a detritivore. *Functional Ecology*, **27**, 807–815.
- Danger M. & Chauvet E. (2013) Elemental composition and degree of homeostasis of fungi: are aquatic hyphomycetes more like metazoans, bacteria or plants? *Fungal Ecology*, **6**, 453–457.
- Danger M., Cornut J., Elger A. & Chauvet E. (2012) Effects of burial on leaf litter quality, microbial conditioning and palatability to three shredder taxa. *Freshwater Biology*, **57**, 1017–1030.
- DeMott W.R. & Gulati R.D. (1999) Phosphorus limitation in *Daphnia*: evidence from a long term study of three hypereutrophic Dutch lakes. *Limnology and Oceanography*, **44**, 1557–1564.
- Denno R.F. & Fagan W.F. (2003) Might nitrogen limitation promote omnivory among carnivorous arthropods? *Ecology*, **84**, 2522–2531.
- Ellison A.M., Bank M.S., Clinton B.D., Colburn E.A., Elliott K., Ford C.R. *et al.* (2005) Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment*, **9**, 479–486.
- Elser J.J., Acharya K., Kyle M., Cotner J., Makino W., Markow T. *et al.* (2003) Growth rate-stoichiometry couplings in diverse biota. *Ecology Letters*, **6**, 936–943.
- Elser J.J., Chrzanowski T.H., Sterner R.W. & Mills K.H. (1998) Stoichiometric constraints on food-web dynamics: a whole-lake experiment on the Canadian Shield. *Ecosystems*, **1**, 120–136.
- Elser J.J., Fagan W.F., Denno R.F., Dobberfuhl D.R., Folarin A., Huberty A. *et al.* (2000a) Nutritional constraints in terrestrial and freshwater food webs. *Nature*, **408**, 578–580.
- Elser J.J., Sterner R.W., Gorokhova E., Fagan W.F., Markow T.A., Cotner J.B. *et al.* (2000b) Biological stoichiometry from genes to ecosystems. *Ecology Letters*, **3**, 540–550.
- Evans-White M.A., Stelzer R.S. & Lamberti G.A. (2005) Taxonomic and regional patterns in benthic macroinvertebrate elemental composition in streams. *Freshwater Biology*, **50**, 1786–1799.
- Ferreira V. & Chauvet E. (2011) Synergistic effects of water temperature and dissolved nutrients on litter decomposition and associated fungi. *Global Change Biology*, **17**, 551–564.
- Ferreira V. & Chauvet E. (2012) Changes in dominance among species in aquatic hyphomycete assemblages do not affect litter decomposition rates. *Aquatic Microbial Ecology*, **66**, 1–11.

- Ferreira V., Eloşegi A., Gulis V., Pozo J. & Graça M.A.S. (2006a) Eucalyptus plantations affect fungal communities associated with leaf litter decomposition in Iberian streams. *Archiv für Hydrobiologie*, **166**, 467–490.
- Ferreira V., Gonçalves A.L., Godbold D.L. & Canhoto C. (2010) Effect of increased atmospheric CO₂ on the performance of an aquatic detritivore through changes in water temperature and litter quality. *Global Change Biology*, **16**, 3284–3296.
- Ferreira V., Gulis V. & Graça M.A.S. (2006b) Whole-stream nitrate addition affects litter decomposition and associated fungi but not invertebrates. *Oecologia*, **149**, 718–729.
- Fink P. & von Elert E. (2006) Physiological responses to stoichiometric constraints: nutrient limitation and compensatory feeding in a freshwater snail. *Oikos*, **115**, 484–494.
- Friberg N. & Jacobsen D. (1994) Feeding plasticity of two detritivore-shredders. *Freshwater Biology*, **32**, 133–142.
- Frost P.C. & Elser J.J. (2002) Growth responses of littoral mayflies to the phosphorus content of their food. *Ecology Letters*, **5**, 232–240.
- Frost P.C., Evans-White M.A., Finkel Z.V., Jensen T.C. & Matzek V. (2005) Are you what you eat? Physiological constraints on organismal stoichiometry in an elementally imbalanced world. *Oikos*, **109**, 18–28.
- Gessner M.O. & Chauvet E. (1993) Ergosterol-to-biomass conversion factors for aquatic hyphomycetes. *Applied and Environmental Microbiology*, **59**, 502–507.
- Gessner M.O., Dobson M. & Chauvet E. (1999) A perspective on leaf litter breakdown in streams. *Oikos*, **85**, 377–384.
- Gessner M.O., Thomas M., Jean-Louis A.M. & Chauvet E. (1993) Stable successional patterns of aquatic hyphomycetes on leaves decaying in a summer cool stream. *Mycological Research*, **97**, 163–172.
- Gonçalves A.L. & Canhoto C. (2009) Decomposition of eucalyptus and alder mixtures: responses to variation in evenness. *Fundamental and Applied Limnology/Archiv für Hydrobiologie*, **173**, 293–303.
- Gonçalves A.L., Chauvet E., Bärlocher F., Graça M.A.S. & Canhoto C. (2014) Top-down and bottom-up control of litter decomposers in streams. *Freshwater Biology*, **59**, 2172–2182.
- Graça M.A.S. (2001) The role of invertebrates on leaf litter decomposition in streams: a review. *International Review Hydrobiology*, **86**, 383–393.
- Graça M.A.S., Bärlocher F. & Gessner M.O. (2005) *Methods to Study Litter Decomposition: A Practical Guide*. Springer-Verlag, New York.
- Graça M.A.S., Maltby L. & Calow P. (1993a) Importance of fungi in the diet of *Gammarus pulex* and *Asellus aquaticus*. II. Effects on growth, reproduction and physiology. *Oecologia*, **96**, 304–309.
- Graça M.A.S., Maltby L. & Calow P. (1993b) Importance of fungi in the diet of *Gammarus pulex* and *Asellus aquaticus* I: feeding strategies. *Oecologia*, **93**, 139–144.
- Graça M.A.S., Maltby L. & Calow P. (1994) Comparative ecology of *Gammarus pulex* (L.) and *Asellus aquaticus* (L.) I: population dynamics and microdistribution. *Hydrobiologia*, **281**, 155–162.
- Gulis V. & Suberkropp K. (2003) Effect of inorganic nutrients on relative contribution of fungi and bacteria to carbon flow from submerged decomposing leaf litter. *Microbial Ecology*, **45**, 11–19.
- Güsewell S. & Gessner M.O. (2009) N: P ratios influence litter decomposition and colonization by fungi and bacteria in microcosms. *Functional Ecology*, **23**, 211–219.
- Hieber M. & Gessner M.O. (2002) Contribution of stream detritivores, fungi, and bacteria to leaf breakdown based on biomass estimates. *Ecology*, **83**, 1026–1038.
- Hillebrand H., Bennett D.M. & Cadotte M.W. (2008) Consequences of dominance: a review of evenness effects on local and regional ecosystem processes. *Ecology*, **89**, 1510–1520.
- Hladyz S., Gessner M.O., Giller P.S., Pozo J. & Woodward G. (2009) Resource quality and stoichiometric constraints on stream ecosystem functioning. *Freshwater Biology*, **54**, 957–970.
- Hodge A., Robinson D. & Fitter A. (2000) Are microorganisms more effective than plants at competing for nitrogen? *Trends in Plant Science*, **5**, 304–308.
- Irons J.G., Oswood M.W. & Bryant J.P. (1988) Consumption of leaf detritus by a stream shredder: influence of tree species and nutrient status. *Hydrobiologia*, **160**, 53–61.
- Iversen T.M. (1974) Ingestion and growth in *Sericostoma personatum* (Trichoptera) in relation to the nitrogen content of ingested leaves. *Oikos*, **25**, 278–282.
- Kearns S.G. & Bärlocher F. (2008) Leaf surface roughness influences colonization success of aquatic hyphomycete spore. *Fungal Ecology*, **1**, 13–18.
- Kimmerer T.W. & Potter D.A. (1987) Nutritional quality of specific tissues and selective feeding by a specialist leaf-miner. *Oecologia*, **71**, 548–551.
- Krauss G.J., Solé M., Krauss G., Schlosser D., Wesenberg D. & Bärlocher F. (2011) Fungi in freshwaters: ecology, physiology and biochemical potential. *FEMS Microbiology Reviews*, **35**, 620–651.
- Lauridsen R.B., Edwards F.K., Bowes M.J., Woodward G., Hildrew A.G., Ibbotson A.T. *et al.* (2012) Consumer-resource elemental imbalances in a nutrient-rich stream. *Freshwater Science*, **31**, 408–422.
- Lauridsen R.B., Edwards F.K., Cross W.F., Woodward G., Hildrew A.G. & Jones J.I. (2014) Consequences of inferring diet from feeding guilds when estimating and interpreting consumer-resource stoichiometry. *Freshwater Biology*, **59**, 1497–1508.
- McKie B.G., Woodward G., Hladyz S., Nistorescu M., Preda E., Popescu C. *et al.* (2008) Ecosystem functioning in stream assemblages from different regions: contrasting responses to variation in detritivore richness, evenness and density. *Journal of Animal Ecology*, **77**, 495–504.

- Nikolcheva L.G., Cockshutt A.M. & Bärlocher F. (2003) Determining diversity of freshwater fungi on decaying leaves: comparison of traditional and molecular approaches. *Applied and Environmental Microbiology*, **69**, 2548–2554.
- Pascoal C. & Cássio F. (2004) Contribution of fungi and bacteria to leaf litter decomposition in a polluted river. *Applied and Environmental Microbiology*, **70**, 5266–5273.
- Pascoal C., Cássio F. & Marvanová L. (2005) Anthropogenic stress may affect aquatic hyphomycete diversity more than leaf decomposition in a low-order stream. *Archiv für Hydrobiologie*, **162**, 481–496.
- Rong Q., Sridhar K.R. & Bärlocher F. (1995) Food selection in three leaf-shredding stream invertebrates. *Hydrobiologia*, **316**, 173–181.
- Schlief J. & Mutz M. (2006) Palatability of leaves conditioned in streams affected by mine drainage: a feeding experiment with *Gammarus pulex* (L.). *Hydrobiologia*, **563**, 445–452.
- Sinsabaugh R.L., Linkins A.E. & Benfield E.F. (1985) Cellulose digestion and assimilation by three leaf-shredding aquatic insects. *Ecology*, **66**, 1464–1471.
- Smith V.H. (2002) Effects of resource supplies on the structure and function of microbial communities. *Antonie van Leeuwenhoek*, **81**, 99–106.
- StatSoft Inc. (2001) Statistica 6.0: Electronic Statistics Textbook. Available at: <http://www.statsoft.com/textbook/stathome.html>.
- Sterner R.W. & Elser J.J. (2002) *Ecological Stoichiometry: the Biology of Elements From Molecules to the Biosphere*. Princeton University Press, Princeton, USA.
- Sterner R.W., Elser J.J., Fee E.J., Guildford S.J. & Chrzanowski T.H. (1997) The light:nutrient ratio in lakes: the balance of energy and materials affects ecosystem structure and process. *The American Naturalist*, **150**, 663–684.
- Sterner R.W. & Hessen D.O. (1994) Algal nutrient limitation and the nutrition of aquatic herbivores. *Annual Review of Ecology, Evolution, and Systematics*, **25**, 1–29.
- Sterner R.W. & Schulz K.L. (1998) Zooplankton nutrition: recent progress and a reality check. *Aquatic Ecology*, **32**, 261–279.
- Suberkropp K. (1992) Interactions with invertebrates. In *The Ecology of Aquatic Hyphomycetes*. (Bärlocher F. ed), pp. 118–134. Springer, Berlin.
- Suberkropp K. (2001) Fungal growth, production, and sporulation during leaf decomposition in two streams. *Applied and Environmental Microbiology*, **67**, 5063–5068.
- Suberkropp K. & Arsuffi T.L. (1984) Degradation, growth, and changes in palatability of leaves colonized by six aquatic hyphomycete species. *Mycologia*, **76**, 398–407.
- Suberkropp K., Arsuffi T.L. & Anderson J.P. (1983) Comparison of degradative ability, enzymatic activity, and palatability of aquatic hyphomycetes grown on leaf litter. *Applied and Environmental Microbiology*, **46**, 237–244.
- Swan C.M., Gluth M.A. & Horne C.L. (2009) Leaf litter species evenness influences nonadditive breakdown in a headwater stream. *Ecology*, **90**, 1650–1658.
- Urabe J., Clasen J. & Sterner R.W. (1997) Phosphorus limitation of *Daphnia* growth: is it real? *Limnology and Oceanography*, **42**, 1436–1443.
- Vannote R.L., Minshall G.W., Cummins K.W., Sedell J.R. & Cushing C.E. (1980) The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, **37**, 130–137.
- Zar J.H. (1996) *Biostatistical Analysis*, 3rd edn. Prentice-Hall International, Englewood Cliffs, NJ.
- Zemek J., Marvanová L., Kuniak L. & Kadlecíková B. (1985) Hydrolytic enzymes in aquatic hyphomycetes. *Folia Microbiologica*, **30**, 363–371.