

## Short Communication

# On the South Atlantic distribution of *Callorhynchus callorhynchus* (Holocephali: Callorhynchidae)

By H. L. López<sup>1</sup>, N. A. San Román<sup>2</sup> and E. E. Di Giacomo<sup>3</sup>

<sup>1</sup>Departamento Científico Zoología Vertebrados, Museo de La Plata, Buenos Aires, Argentina; <sup>2</sup>Centro Austral de Investigaciones Científicas (CADIC-CONICET), Tierra del Fuego, Argentina; <sup>3</sup>Instituto de Biología Marina y Pesquera Almirante Storni, Río Negro, Argentina

The four species of the family Callorhynchidae are restricted to the southern hemisphere (Didier 1995) and have the most littoral-related distribution among holocephalans.

*Callorhynchus callorhynchus* was first recorded in Argentine waters by Vaillant in 1891 sub *C. antarcticus*, and from the mouth of the Santa Cruz River (Menni et al. 1984). Information on biology, fishery, biomass and distribution of *C. callorhynchus* is available in Bellisio et al. (1979), Menni and Gosztanyi (1982), Menni and López (1984), Di Giacomo and Perier (1991, 1994, 1996), Di Giacomo (1992), and Di Giacomo et al. (1994).

The northern Atlantic limit of *C. callorhynchus* is in the Brazilian state of Sao Paulo (Figuereido 1977). Other reports from the Atlantic were those for the Río de la Plata (Rios and Pereira 1986), and for several localities along the Bonarensan and Patagonian littoral (Bellisio et al. 1979; Menni et al. 1981; Menni and López 1984). On the basis of the results of the JAMARC expeditions (1976–79), Nakamura et al. (1986) reported a distribution which ranged from the Argentine-Chilean continental shelf to Chiloe Island, but not including the Beagle Channel.

Rios and Pereira (1986) recorded this species for the Río de la Plata and for the littoral off the coasts of Uruguay and Argentina, on sandy and muddy bottoms at depths of 10–80 m. Menni and López (1984) assigned *C. callorhynchus* to the group of the Atlantic inner shelf mixed fish fauna occurring at temperatures of 5–10°C and depths of 22–115 m, and noted an association with *Stromateus brasiliensis* and *Discopyge tshudii*. Di Giacomo (1992) reported on the vertical distribution of *C. callorhynchus* in the Argentine Sea, recording a depth limit of 116 m. On examining its reproductive biology in San Matias Gulf (Northern Patagonia), Di Giacomo and Perier (1994) collected egg cases at depths of from 20 to 40 m to 104 m, which would reflect the migration to shallow waters to breed. Subsequent reports for the gulf by divers and personal observation by one of the authors confirms that breeding takes place along the littoral at depths of 18–25 m.

*Callorhynchus callorhynchus* is herein first reported for the Beagle Channel on the basis of material collected at Puerto España (Argentine sector of the channel). Specimens (MLP 9285 and 9286), one 430 mm total length (TL) male and one 500 mm TL female were caught at 15 m depth by trawling nets during cruises performed within the framework of the project 'Biological basis for management of mixed catches of "centolla" and "centellón" crabs in the Beagle Channel' (G. Lavrich, collector).

The spelling herein applied for the generic name *Callorhynchus* follows Didier's (1995) nomenclatorial considerations.

The occurrence of *C. callorhynchus* in the Argentine sector

of the Beagle Channel confirms the broad distribution range of this species, adding new elements to preceding zoogeographical proposals regarding the South American Atlantic littoral (e.g. Balech 1954; López 1963).

## References

- Balech, E., 1954: División Zoogeográfica del litoral sudamericano. Rev. Biol. Marina, Valparaiso 4, 231–238.
- Bellisio, N. B.; López, R. B.; Torno, A., 1979: Peces marinos patagónicos. Secretaría de Intereses Marítimos, Argentina, 279 pp.
- Di Giacomo, E. E., 1992: Distribución de la población de pez gallo (*Callorhynchus Callorhynchus*) en el golfo San Matias, Argentina. Frente Marítimo 12, Sec. A, 113, 118.
- Di Giacomo, E. E.; Parma, A. M.; Orensanz, J. M., 1994: Food consumption by the cockfish, *Callorhynchus callorhynchus* (Holocephali, Callorhynchidae), from Patagonia (Argentina). Environ. Biol. Fishes 40, 199–211.
- Di Giacomo, E. E.; Perier, M. R., 1991: Evaluación de la biomasa y explotación comercial de pez gallo (*Callorhynchus Callorhynchus*) en el golfo San Matias. Frente Marítimo 9, Sec. A, 7–13.
- Di Giacomo, E. E.; Perier, M. R., 1994: Reproductive biology of the cockfish, *Callorhynchus callorhynchus* (Holocephali: Callorhynchidae), in Patagonian waters (Argentina). Fish. Bull. 92, 531–539.
- Di Giacomo, E. E.; Perier, M. R., 1996: Feeding habits of cockfish, *Callorhynchus callorhynchus* (Holocephali: Callorhynchidae), in Patagonian waters (Argentina). Mar. Freshw. Res. 47, 801–808.
- Didier, D. A., 1995: Phylogenetic systematics of extant Chimaeroid fishes (Holocephali, Chimaeroidae). Am. Mus. Novit. 3119, New York.
- Figuereido, J. L., 1977: Manual de peixes marinhos do Sudeste do Brasil. I. Introdução, raias e quimeras. Sao Paulo, Brasil: Museu de Zoologia, University of Sao Paulo, 104 pp.
- López, R. B., 1963: Problemas sobre la distribución geográfica de los peces marinos sudamericanos. Rev. Mus. Arg. C. Nat. 'B. Rivadavia', Hidrobiol. 1 (3), 111–135.
- Menni, R. C.; Gosztanyi, A. E., 1982: Benthic and semidemersal fish associations in the Argentine Sea. Studies on Neotropical Fauna and Environment 17, 1–29.
- Menni, R. C.; López, H. L., 1984: Distributional patterns of Argentine marine fishes. Physis, Sec. A, 42 (103), 71–85.
- Menni, R. C.; López, H. L.; García, M. L., 1981: Lista comentada de las especies de peces colectadas durante la campaña V del B/L "Shinkai Maru" en el Mar Argentino (25/8–15/09/78). En: Campañas de Investigación realizadas en el Mar Argentino por los B/L "Shinkai Maru" y "Walter Herwig" y el B/P "Marburg". Años 1978 y 1979. Resultados de la parte Argentina: INIDEP No. 383, pp. 267–280. Inst. Invest. y Desarrollo Pesquero (INIDEP) (Ed. V. Angelescu), Mar del Plata.
- Menni, R. C.; Ringuelet, R. A.; Aramburu, R. H., 1984: Peces marinos de la Argentina y Uruguay. Buenos Aires: Ed. Hemisferio Sur, 359 pp.
- Nakamura, Y.; Inada, T.; Takeda, M.; Hatanaka, H., 1986: Important fishes trawled off Patagonia. In: Japan Marine Fishery Resource Research Centre (Jamarc). Ed. J. Nakamura, 369 pp, Tokyo.
- Rios, E. C.; Pereira, J., 1986: 'Pez elefante', un pez malacófago. Com. Soc. Malac. Uruguay 6 (51), 427–428.
- Authors' address: Hugo L. López, Departamento Científico Zoología, Vertebrados, Museo de La Plata, - CIC, paseo del Bosque s/n 1900 La Plata, Buenos Aires, Argentina