



SCCS Working Paper WP-SCCS 2014-01

Sleipner CO₂ securely stored deep beneath seabed, in spite of unexpected Hugin fracture discovery

13th January 2014

Authors:

Prof Stuart Haszeldine

Dr Vivian Scott, Dr Simon Shackley, Dr Stuart Gilfillan,

Dr Leslie Mabon, Dr Gareth Johnson

Scottish Carbon Capture and Storage (SCCS)

School of GeoSciences, University of Edinburgh, EH9 3JW, Scotland, UK s.haszeldine@ed.ac.uk

No competing commercial or business interests. Funded by EPSRC, NERC, EU, SFC

Summary

General readers of Nature may now think that the proposition to store carbon dioxide in deep geological strata is doomed to fail (Monastersky 2013). This is far from the case, as a more balanced review could easily have pointed out. It is now important to provide an alternative perspective, based on published information, that geological storage of CO₂ by deep injection for CCS is both sufficiently secure, and knowable in its environmental impacts. Furthermore, research has shown that there is good support from many parts of the public, although qualified, for CCS as an essential part of a response to the threat of global climate change and ocean acidification.

Sleipner CO₂ securely stored deep beneath seabed, in spite of unexpected Hugin fracture discovery

There is definitely an essential role for journalistic media to challenge CCS developers to explain and to justify their proposition. However, the Monastersky news review seeks to link spatially disconnected observations as cause and inevitable consequence, with no analysis of the evidence, or the feasibility. The ECO₂ project has attempted to conflate a seabed fracture seeping shallow CH₄ with future leakage of CO₂ from the Sleipner project deep storage site 25km distant. This is not the first sensationalist news comment from the ECO₂ project (Reitz 2012), but we are still waiting, several years after “discovery”, for the peer-reviewed evidence to demonstrate the implied causality. The framing of these recent assertions needs to be formally challenged as a science process.

The Sleipner project examined here is the first purpose-engineered sub-sea geological CO₂ storage site in the world. Commencing in 1996, the Norwegian oil company Statoil has injected about 1 million tons of CO₂ per year, driven by the commercial imperative to avoid offshore carbon taxes combined with the ambition to behave in a more sustainable way.

The progress of this injection has been closely monitored by eight seismic reflection surveys, at approximately 2 year intervals. These high resolution records clearly show that CO₂ is being retained within the intended sandstone formation, with no sign of leakage (Chadwick et al 2011). The ascent and lateral dispersion of this CO₂ plume can be accurately modelled and mass-balanced (Cavanagh & Haszeldine 2014), demonstrating that CO₂ security is achieved and understood.

The ECO₂ project has obtained interesting and high resolution images of a fracture, open at the seabed, emitting small volume seeps of water and dissolved gases - which is not unusual in hydrocarbon provinces, (Judd and Hovland 2009). The fracture is observed on 1996 seismic reflection surveys for region around the Sleipner oilfield (Furre et al 2013), and so pre-dates engineered CO₂ injection. The feature was discovered separately in 2011 by a University of Bergen cruise, which named this the “Hugin Fracture”; that was visited twice more in 2012 by the ECO₂ project to obtain high resolution sonar images (Reitz 2012). Published images show the Hugin Fracture is about 5m wide and 3.5km long, as a zig-zag of ENE and NW linear subsets (Baumberger 2013).

However, post-glacial fractures or faulting are not particularly new, and can have several causes. Post glacial ice sheet unloading, stress relief and rebound, forms fault scarps onshore (Lundqvist & Lagerback 1976). Additionally, seabed subsidence is well documented due to hydrocarbon extraction (Mes 1990), as are linear fissures and chains of sea bed pockmarks due to dewatering of deeper sands or ascent of deep gas (Judd and Hovland 2009). In this case (Furre et al 2013), the Hugin

Fracture overlies the edge of a shallow buried sub-glacial tunnel valley, which is likely to be partially filled with sand, and contains shallow methane. The fracture may have formed as a compaction drape over more resistant sand.

It is therefore not surprising that the Hugin Fracture seeps fresh water and biogenic methane of shallow provenance (Reitz 2012); this is what would be expected. No CO₂ from injection at Sleipner has been detected. Traces of thermogenic (deeper) methane are stated to be present (Schaps 2012), although these are common around this part of the North Sea (Hegglund 1997). The overall seepage is tiny, about 1 tonne of gas per year (Retiz 2012). Methane can be derived, through geological time, vertically or laterally from beneath the Utsira CO₂ storage reservoir, migrating vertically, then laterally through a complex spaghetti of glacial tunnel valley conduits. Clear fingerprinting of tracer gases could assist, but the identification of specific methane sources is elusive. Consequently the article title and graphic (Monastersky 2013 Fig 1) juxtaposing stored CO₂ and fracture leakage, is entirely misleading.

ECO₂ have specifically conflated the discovery of a seabed fracture with the possibility of leakage from the Utsira Sand CO₂ store for the Sleipner project. We consider that is rather disingenuous. Even though disclaimers are written into the article, the subliminal suggestion of leakage is being made, to the point of claims that are neither supported nor quantified: *"We are saying it is very likely some thing will come out in the end,"* says Klaus Wallmann, ECO₂ coordinator.

However ECO₂ provide no depth profile beneath the fracture, indicating the downwards extent, which is what we would like to see, to support assertions of connectivity and potential leakage. Additional information might include chemical analyses of emitted gases, and especially their isotopic signatures of H, C, O and fingerprints of trace noble gases, which are diagnostic of specific CH₄ or CO₂ origins (Gilfillan et al 2011). The geological setting is important to remember: the Utsira storage reservoir is overlain by primary seal mudrock from seabed to 800 mbsl, so there is still 550 m of mudrock sealing between the engineered CO₂ store and the supposed base of the Hugin Fracture at 150 mbsl. In other regions of the North Sea, older and more compacted mudrocks have clearly retained natural CO₂ accumulations for tens of millions of years (Lu et al 2009). That is much longer than the thousands of years required here for the purpose of carbon abatement. We would like to see a much better assessment of why one fracture of limited depth, far distant from the well-known CO₂ plume, is expected to be a problem, when it can be seen that the huge additional over-engineering of the mudrock topseal above Utsira Sand is, according to all the evidence to date, adequately retaining all CO₂. The story can readily be spun in the opposite direction: viz "CO₂ securely stored in spite of unexpected discovery". This indicates an unbalance in the way that the ECO₂ findings are reported. The ECO₂ hypotheses are highly speculative, and an interpretation is offered, but is unsupported by data. What is required is an academically balanced and impartial discussion.

The short discussion about vertical chimneys and pipes (Monastersky 2013), which could act as conduits for rapid gas or water escape through sediments overlying the Utsira sand CO₂ reservoir, is also very old news (Hegglund 1997, Zweigel 2000, Judd & Hovland 2009). To bring something additional, ECO₂ could be looking to demonstrate that vertical fluid chimneys from the Utsira CO₂ reservoir are active, rather than dormant, and link CH₄ or CO₂ migration pathways to seabed features which currently seep. These are potentially much greater potential conduits for vertical CO₂ movement than the Hugin Fracture singularity. Fluid chimneys around the Sleipner site were recognized before injection started (Hegglund 1997, Zweigel 2000), and sequential surveys during CO₂ injection enable confident statements to be made that there is no change in gas above the CO₂ reservoir. This demonstrates the value of baseline data and appropriate monitoring.

The article then correctly states that even if CO₂ leaks, it does not seem to be a problem, and finally drifts off into vague economic speculation without any clear relevance or conclusion. What the focus should be, is that more than 60% of injected CO₂ is inevitably trapped as small bubbles in sediment pores by residual saturation (Goater et al 2013) and cannot leak. The other 40% of CO₂ is structurally trapped far away from the Hugin Fracture. Even if leakage occurred, this CO₂ could migrate for only short distances, as it rapidly dissolves by interaction with subsurface porewater; this has been quantified by geochemical studies of natural CO₂ accumulations (Gilfillan et al 2009). Even if CO₂ does leak to the surface (or seabed), then on-land it is dispersed by wind, such that the effects on human health are minimal (Roberts et al 2011). On the North Sea shelf, subsea CO₂ leakage would be dispersed by ocean currents, such that direct local environmental impacts are modelled to be insignificant (Blackford et al 2009).

Monastersky (2013) chooses to quote that that any leakage “*would be a disaster for public opinion*”. A deeper dive into the evidence, provided by social science researchers on the same ECO₂ project (Mabon et al 2014), suggests that leakage – whilst of course undesirable – need not in itself be a show-stopper for the geological storage of CO₂. Many members of the public do understand that environmental systems are complex and that scientists and engineers do not know everything. What publics want to see is not a reassurance that a site will *not* leak, rather that adequate monitoring, control and remediation procedures are in place *should* any leak be discovered (Brunsting et al, 2012; Mabon et al, 2014). Indeed, attempts to convince publics that a site will *never* leak can induce skepticism, and may even reduce public support for CO₂ storage if an impression is created that scientists and/or developers have given insufficient attention to the possibility of an unexpected scenario occurring (Howell et al, 2014).

Furthermore, although it is true that offshore CO₂ storage removes some of the factors that can drive public opposition to CCS (van Noorden 2010) – effects on real estate prices, perceived immediate risks to human health – it is important to caution against over-stating the claim in Monastersky (2013) that offshore CCS will be ‘easier’ in terms of public acceptability. Many drivers of public perception, such as place-attachment and perceptions of fairness, do not correspond easily to land-sea boundaries. To say that offshore CO₂ storage is always going to be ‘easier’ in terms of public acceptability thus flattens the very complex, dynamic and contingent nature of public perceptions of energy infrastructure.

Prudent, stage-by-stage testing (as opposed to excessive precaution which means that nothing is ever achieved) is what is needed. Then, as more is learned about the risks, and how they can be managed, the endeavour can be scaled-up. For leakage of CO₂ from engineered storage to have any effect on stimulating climate warming, then leakage would need to be at the scale of a percentage a year or greater (Teng & Tondeur 2007, Stone et al 2009); that requires such a large number of mistakes to occur at a majority of CO₂ storage sites that the proposition becomes unfeasible. The Hugin Fracture, seeping at 1 tonne/day (Reitz 2012) falls well below that threshold. As a long-term risk, geological CO₂ storage will out-perform most types of carbon mitigation, such as forestry.

In summary, we find that Monastersky (2013) takes an unduly negative view of the Sleipner CO₂ storage experiment, and the quoted researchers offer a mixture of alarmism and innuendo. A seabed fracture has been imaged by a novel technique, and forms one of many fluid conduits beneath the shallow North Sea. There is no evidence of methane movement in the overburden, or unplanned CO₂ migration from the reservoir, during the Sleipner CO₂ storage project history. The multiple barriers to CO₂ leakage – residual saturation in the reservoir, thick and multiple seal rocks, dissolution in overburden porewaters, laterally extensive migration dispersion in overburden tunnel valleys, and lack

of environmental impact modelled for CO₂ at the seabed, all provide good reasons for public confidence. The abundance of baseline and monitoring data, also give confidence that irregularities can be detected, and solved, long before events become problems. Many things are not perfectly explained. Many things could go wrong. But nothing has. Therefore, it is very encouraging that the immature science, which pragmatically selected the Utsira sand reservoir as storage for the Sleipner project, has proven robust. The subsequent 18 years of global endeavor gives high confidence in the ability of geo-science and engineering systems to identify good CO₂ storage, to operate that safely, and monitor secure retention (Scott et al 2012). We remain skeptical of science by press-release, and wait to evaluate the publication of convincing data on the Hugin Fracture; ECO₂ have to put up or shut up; the Sleipner project seems to be storing CO₂ securely in the Utsira reservoir.

References

- Baumberger T., Denny A.R., Pedersen R.B. (2013) High-resolution interferometric synthetic aperture sonar (HISAS 1030). ECO₂ WP11 Workshop, page 19. <http://oceanrep.geomar.de/22459/1/ECO2%20D11.2.pdf>
- Blackford J., Jones N., Proctor R., Holt J., Widdicombe S., Lowe D. & Rees A. 2009. An initial assessment of the potential environmental impact of CO₂ escape, from marine carbon capture and storage systems. Proceedings of the Institution of Mechanical Engineers, Part A, Journal of Power and Energy 223(A3): 269-280.
- Brunsting, S, Mastop, J, Pol, M, Kaiser, M, Zimmer, R, Shackley, S, Mabon, L, Howell, R (2012) SiteChar Deliverable 8.2: Trust building and raising public awareness' ECN: Amsterdam. <http://www.sitechar-co2.eu/SciPublicationsData.aspx?IdPublication=314&IdType=557>
- Cavanagh A.J., and Haszeldine R.S. (2014) The Sleipner storage site: capillary flow modeling of a layered CO₂ plume requires fractured shale barriers within the Utsira Formation. International Journal of Greenhouse Gas Technology. In press.
- Chadwick, R. A., Noy, D., Arts, R., Eiken, O., (2011) Latest time-lapse seismic data from Sleipner yield new insights into CO₂ plume development. GHGT-9 Proceedings, Energy Procedia 1, 2103-2110.
- Furre A., Janbu A.D., & Hagen S. (2013) Characterisation of Sub-sea CO₂ Storage Complexes - Focus on the Greater Sleipner Area. Technologies for Sustainable Use of the Deep Sub-surface. European Association of Geoscientists and Engineers, 30 September - 4 October 2013 Pau, France. DOI: 10.3997/2214-4609.20131603
- Gilfillan, S.M.V., Lollar, B.S., Holland, G., Blagburn, D., Stevens, S., Schoell, M., Cassidy, M., Ding, Z., Zhou, Z., Lacrampe-Couloume, G., Ballentine C.J., 2009. Solubility trapping in formation water as dominant CO₂ sink in natural gas fields. Nature 458, 614-618.
- Gilfillan, S.M.V., Wilkinson, M., Haszeldine, R.S., Shipton, Z., Nelson, S.T. and Poreda, R.J. (2011) He and Ne as tracers of natural CO₂ migration up a fault from a deep reservoir, International Journal of Greenhouse Gas Control, 5, no. 6, p. 1507-1516 DOI:10.1016/j.iggc.2011.08.008
- Goater A.L., Bijeljic B., Blunt M.J. 2013 Dipping open aquifers—The effect of top-surface topography and heterogeneity on CO₂ storage efficiency. International Journal of Greenhouse Gas Control 17,318–331
- Heggland, R., 1997, Detection of gas migration from a deep source by the use of exploration 3D seismic data. Marine Geology 137, p 41-47.
- Howell, R, Shackley, S, Mabon, L, Ashworth, P, and Jeanneret, T (2014) 'Engaging the public with low-carbon energy technologies: results from a Scottish large group process' Energy Policy In Press <http://dx.doi.org/10.1016/j.enpol.2013.11.041>
- Judd A G, Hovland M. (2009) Seabed Fluid Flow: The Impact of Geology, Biology and the Marine Environment. Cambridge Univ Press.
- Lu J, Wilkinson M, Haszeldine RS, Fallick AE. Geology (2009), Long-term performance of a mudrock seal in natural CO₂ storage v37; no. 1; p. 35–38
- Lundqvist J. and Lagerback R (1976) The Pärve Fault: A late-glacial fault in the Precambrian of Swedish Lapland Geologiska Föreningen i Stockholm Förhandlingar, 98, 45-51.
- Mabon, L, Shackley, S, and Bower-Bir, N (2014) 'Perceptions of sub-seabed carbon dioxide storage in Scotland and implications for policy: A qualitative study' Marine Policy 45: 9-15.

- Mes M.J. (1990) Ekofisk Reservoir Pressure Drops and Seabed Subsidence. 6241-MS, Offshore Technology Conference, 7 May-10 May 1990, Houston, Texas
- Monastersky R (2013) Seabed scars raise questions over carbon storage plan. *Nature*, 504, 309 – 310.
- Reitz A. (2012) North Sea fracture discovered. ECO₂ news release 2 Oct 2012
<http://www.eco2-project.eu/newsarticles/items/north-sea-fracture-discovered.html>
- Roberts, J. J. Wood, R. A. and Haszeldine R. S. (2011) Assessing the health risks of natural CO₂ seeps in Italy. *Proceedings of the National Academy of Sciences of the United States of America* vol. 108 no. 40 16545-16548
DOI: 10.1073/pnas.1018590108
- Scott, V., Gilfillan, S.M.V., Markusson, N., Chalmers, H., Haszeldine, R.S. Last chance for carbon capture and storage. (2012) *Nature Climate Change* 3, 105-111 DOI: 10.1038/nclimate1695.
- Schaps K (2012) Subsea ravine leaks a new headache for carbon capture. Reuters, 14 Sept 2012
<http://uk.reuters.com/article/2012/09/17/ccs-nsea-fracture-idUKL5E8KACES20120917>
- Stone, E., Lowe, J. A., Shine, K.P. (2009) The impact of carbon capture and storage on climate. *Energy Environ. Sci.*, 2, 81-91 DOI: 10.1039/B807747A
- Teng, F. and Tondeur, D. (2007) Efficiency of Carbon storage with leakage: Physical and economical approaches. *Energy*, Volume 32, Issue 4, p 540–548.
- Van Noorden R. (2010) Carbon sequestration: Buried trouble. *Nature* 463, 871-873.
- van der Vegt, P., Janszen, A., Moscariello, A., (2012) Tunnel valleys: current knowledge and future perspectives, in Huuse, M., Redfern, J., Le Heron, D. P., Dixon, R. J., Moscariello, A. & Craig, J. (eds), *Glaciogenic Reservoirs and Hydrocarbon Systems*, Geological Society, London, Special Publications, 368, p 75-97.
- Zweigel, P. (2000) Seismic anomaly maps in the Neogene, of Sleipner area. SINTEF Report, open access.
<http://www.sintef.no/Projectweb/IK-23430000-SACS/Technical-reports/Other-technical-reports/>

This is a full version of a shorter article submitted to *Nature* (Contributions)

To be published around 20 Jan 2014

<http://www.nature.com/nature/authors/gta/commsarising.html>