

Circle valued Morse theory and Novikov homology

Andrew Ranicki*

*Department of Mathematics and Statistics
University of Edinburgh, Scotland, UK*

arXiv:math.AT/0111317 v1 30 Nov 2001

*Lecture given at the:
Summer School on High-dimensional Manifold Topology
Trieste, 21 May – 8 June 2001*

LNS

*aar@maths.ed.ac.uk

Abstract

Traditional Morse theory deals with real valued functions $f : M \rightarrow \mathbb{R}$ and ordinary homology $H_*(M)$. The critical points of a Morse function f generate the Morse-Smale complex $C^{MS}(f)$ over \mathbb{Z} , using the gradient flow to define the differentials. The isomorphism $H_*(C^{MS}(f)) \cong H_*(M)$ imposes homological restrictions on real valued Morse functions. There is also a universal coefficient version of the Morse-Smale complex, involving the universal cover \widetilde{M} and the fundamental group ring $\mathbb{Z}[\pi_1(M)]$.

The more recent Morse theory of circle valued functions $f : M \rightarrow S^1$ is more complicated, but shares many features of the real valued theory. The critical points of a Morse function f generate the Novikov complex $C^{Nov}(f)$ over the Novikov ring $\mathbb{Z}((z))$ of formal power series with integer coefficients, using the gradient flow of the real valued Morse function $\overline{f} : \overline{M} = f^*\mathbb{R} \rightarrow \mathbb{R}$ on the infinite cyclic cover to define the differentials. The Novikov homology $H_*^{Nov}(M)$ is the $\mathbb{Z}((z))$ -coefficient homology of \overline{M} . The isomorphism $H_*(C^{Nov}(f)) \cong H_*^{Nov}(M)$ imposes homological restrictions on circle valued Morse functions.

Chapter 1 reviews real valued Morse theory. Chapters 2,3,4 introduce circle valued Morse theory and the universal coefficient versions of the Novikov complex and Novikov homology, which involve the universal cover \widetilde{M} and a completion $\widehat{\mathbb{Z}[\pi_1(M)]}$ of $\mathbb{Z}[\pi_1(M)]$. Chapter 5 formulates an algebraic chain complex model (in the universal coefficient version) for the relationship between the $\mathbb{Z}((z))$ -module Novikov complex $C^{Nov}(f)$ of a circle valued Morse function $f : M \rightarrow S^1$ and the \mathbb{Z} -module Morse-Smale complex $C^{MS}(f_N)$ of the real valued Morse function $f_N = \overline{f}| : M_N = \overline{f}^{-1}[0, 1] \rightarrow [0, 1]$ on a fundamental domain of the infinite cyclic cover \overline{M} .

Keywords: circle valued Morse theory, Novikov complex, Novikov homology

AMS numbers: 57R70, 55U15

Contents

1	Introduction	1
2	Real valued Morse theory	4
3	The Novikov complex	8
4	Novikov homology	10
5	The algebraic model for circle valued Morse theory	14
	References	21

1 Introduction

The Morse theory of circle valued functions $f : M \rightarrow S^1$ relates the topology of a manifold M to the critical points of f , generalizing the traditional theory of real valued Morse functions $M \rightarrow \mathbb{R}$. However, the relationship is somewhat more complicated in the circle valued case than in the real valued case, and the roles of the fundamental group $\pi_1(M)$ and of the choice of gradient-like vector field v are more significant (and less well understood).

The Morse-Smale complex $C = C^{MS}(M, f, v)$ is defined geometrically for a real valued Morse function $f : M^m \rightarrow \mathbb{R}$ and a suitable choice of gradient-like vector field $v : M \rightarrow \tau_M$. In general, there is a $\mathbb{Z}[\pi]$ -coefficient Morse-Smale complex for each group morphism $\pi_1(M) \rightarrow \pi$, with

$$C_i = \mathbb{Z}[\pi]^{c_i(f)}$$

if there are $c_i(f)$ critical points of index i . The differentials $d : C_i \rightarrow C_{i-1}$ are defined by counting the \tilde{v} -gradient flow lines in the cover \tilde{M} of M classified by $\pi_1(M) \rightarrow \pi$. In the simplest case $\pi = \{1\}$ this is just $\tilde{M} = M$, and if $p \in M$ is a critical point of index i and $q \in M$ is a critical point of index $i-1$ the (p, q) -coefficient in d is the number $n(p, q)$ of lines from p to q , with sign chosen according to orientations. The homology of the Morse-Smale complex is isomorphic to the ordinary homology of M

$$H_*(C^{MS}(M, f, v)) \cong H_*(M)$$

so that

- (a) the critical points of f can be used to compute $H_*(M)$,
- (b) $H_*(M)$ provides lower bounds on the number of critical points in any Morse function $f : M \rightarrow \mathbb{R}$, which must have at least as many critical points of index i as there are \mathbb{Z} -module generators for $H_i(M)$ (Morse inequalities).

Basic real valued Morse theory is reviewed in Chapter 2.

In the last 40 years there has been much interest in the Morse theory of circle valued functions $f : M^m \rightarrow S^1$, starting with the work of Stallings [36], Browder and Levine [3], Farrell [8] and Siebenmann [35] on the characterization of the maps f which are homotopic to the projections of fibre bundles over S^1 : these are the circle valued Morse functions without any critical points.

About 20 years ago, Novikov ([17],[18],[19],[20] (pp. 194–199)) was motivated by problems in physics and dynamical systems to initiate the general Morse theory of closed 1-forms, including circle valued functions $f : M \rightarrow S^1$ as the most important special case. The new idea was to use the *Novikov ring* of formal power series with an infinite number of positive coefficients and a finite number of negative coefficients

$$\mathbb{Z}((z)) = \mathbb{Z}[[z]][z^{-1}] = \left\{ \sum_{j=-\infty}^{\infty} n_j z^j \mid n_j \in \mathbb{Z}, n_j = 0 \text{ for all } j < k, \text{ for some } k \right\}$$

as a counting device for the gradient flow lines of the real valued Morse function $\bar{f} : \bar{M} = f^*\mathbb{R} \rightarrow \mathbb{R}$ on the (non-compact) infinite cyclic cover \bar{M} of M , with the indeterminate z

corresponding to the generating covering translation $z : \overline{M} \rightarrow \overline{M}$. For f the number of gradient flow lines starting at a critical point $p \in M$ is finite in the generic case. On the other hand, for \overline{f} the number of gradient flow lines starting at a critical point $\overline{p} \in \overline{M}$ may be infinite in the generic case, so the counting methods for real and circle valued Morse theory are necessarily different.

The *Novikov complex* $\widehat{C} = C^{Nov}(M, f, v)$ is defined for a circle valued Morse function $f : M^m \rightarrow S^1$ and a suitable choice of gradient-like vector field $v : M \rightarrow \tau_M$. In general, there is a $\widehat{\mathbb{Z}[\Pi]}$ -coefficient Novikov complex for each factorization of $f_* : \pi_1(M) \rightarrow \pi_1(S^1) = \mathbb{Z}$ as $\pi_1(M) \rightarrow \Pi \rightarrow \mathbb{Z}$, with $\widehat{\mathbb{Z}[\Pi]}$ a completion of $\mathbb{Z}[\Pi]$, with

$$\widehat{C}_i = \widehat{\mathbb{Z}[\Pi]}^{c_i(f)}$$

if there are $c_i(f)$ critical points of index i . The differentials $d : C_i \rightarrow C_{i-1}$ are defined by counting the \widetilde{v} -gradient flow lines in the cover \widetilde{M} of M classified by $\pi_1(M) \rightarrow \Pi$. The construction of the Novikov complex for arbitrary $\widehat{\mathbb{Z}[\Pi]}$ is described in Chapter 3. In the simplest case

$$\Pi = \mathbb{Z}, \mathbb{Z}[\Pi] = \mathbb{Z}[z, z^{-1}], \widehat{\mathbb{Z}[\Pi]} = \mathbb{Z}((z)), \widetilde{M} = \overline{M} = f^*\mathbb{R}.$$

For a critical point $\overline{p} \in \overline{M}$ of index i and a critical point $\overline{q} \in \overline{M}$ of an index $i - 1$ the $(\overline{p}, \overline{q})$ -coefficients in d is

$$\widehat{n}(\overline{p}, \overline{q}) = \sum_{j=k}^{\infty} n(\overline{p}, z^j \overline{q}) z^j \in \mathbb{Z}((z))$$

with $n(\overline{p}, z^j \overline{q})$ the signed number of \overline{v} -gradient flow lines of the real valued Morse function $\overline{f} : \overline{M} \rightarrow \mathbb{R}$ from \overline{p} to the translate $z^j \overline{q}$ of \overline{q} , and $k = \lceil \overline{f}(\overline{p}) - \overline{f}(\overline{q}) \rceil$. The convention is that the generating covering translation $z : \overline{M} \rightarrow \overline{M}$ is to be chosen parallel to the downward gradient flow $v : M \rightarrow \tau_M$, with

$$\overline{f}(zx) = \overline{f}(x) - 1 \in \mathbb{R} \quad (x \in \overline{M}).$$

In particular, this means that for $f = 1 : M = S^1 \rightarrow S^1$

$$z : \overline{M} = \mathbb{R} \rightarrow \overline{M} = \mathbb{R}; x \mapsto x - 1.$$

Circle valued Morse theory is necessarily more complicated than real valued Morse theory. The Morse-Smale complex $C^{MS}(M, f : M \rightarrow \mathbb{R}, v)$ is an absolute object, describing M on the chain level, with $c_0(f) > 0$, $c_m(f) > 0$. This is the algebraic analogue of the fact that every continuous function $f : M \rightarrow \mathbb{R}$ on a compact space attains an absolute minimum and an absolute maximum. By contrast, the Novikov complex $C^{Nov}(M, f : M \rightarrow S^1, v)$ is a relative object, measuring the chain level difference between f and the projection of a fibre bundle (= Morse function with no critical points). A continuous function $f : M \rightarrow S^1$ can just go round and round! The connection between the geometric properties of f and the algebraic topology of M is still not yet completely understood, although there has been much progress in the work of Pajitnov, Farber, the author and others.

The *Novikov homology groups* of a space M with respect to a cohomology class $f \in [M, S^1] = H^1(M)$ are defined by

$$H_*^{Nov}(M, f) = H_*(\mathbb{Z}((z)) \otimes_{\mathbb{Z}[z, z^{-1}]} C(\overline{M})) .$$

The homology groups of the Novikov complex are isomorphic to the Novikov homology groups

$$H_*(C^{Nov}(M, f, v)) \cong H_*^{Nov}(M, f) .$$

By analogy with the real valued case :

- (a) the critical points of f can be used to compute $H_*^{Nov}(M, f)$,
- (b) $H_*^{Nov}(M, f)$ provides lower bounds on the number of critical points in any Morse function $f : M \rightarrow S^1$, which must have at least as many critical points of index i as there are $\mathbb{Z}((z))$ -module generators for $H_i^{Nov}(M, f)$ (Morse-Novikov inequalities).

Novikov homology is constructed in Chapter 4, for arbitrary $\widehat{\mathbb{Z}[\Pi]}$ -coefficients..

Novikov conjectured ([1]) that for a generic class of gradient-like vector fields $v \in \mathcal{GT}(f)$ the functions $j \mapsto n(\overline{p}, z^j \overline{q})$ have subexponential growth.

Let $S \subset \mathbb{Z}[z]$ be the subring of the polynomials $s(z)$ such that $s(0) = 1$, which (up to sign) are precisely the polynomials which are invertible in the power series ring $\mathbb{Z}[[z]]$. The localization $S^{-1}\mathbb{Z}[z, z^{-1}]$ of $\mathbb{Z}[z, z^{-1}]$ is identified with the subring of $\mathbb{Z}((z))$ consisting of the quotients $\frac{r(z)}{s(z)}$ with $r(z) \in \mathbb{Z}[z, z^{-1}]$, $s(z) \in S$.

Pajitnov [23],[24] constructed a C^0 -dense subspace $\mathcal{GECT}(f) \subset \mathcal{GT}(f)$ of gradient-like vector fields v for which the differentials in the Novikov complex $C^{Nov}(M, f, v)$ are rational

$$\widehat{n}(\overline{p}, \overline{q}) = \sum_{j=-\infty}^{\infty} n(\overline{p}, z^j \overline{q}) z^j \in S^{-1}\mathbb{Z}[z, z^{-1}] \subset \mathbb{Z}((z))$$

and the functions $j \mapsto n(\overline{p}, z^j \overline{q})$ have polynomial growth. The idea is to cut M along the inverse image $N = f^{-1}(0)$ (assuming $0 \in S^1$ is a regular value of f), giving a fundamental domain

$$(M_N; N, z^{-1}N) = \overline{f}^{-1}([0, 1]; \{0\}, \{1\})$$

for $\overline{f} : \overline{M} \rightarrow \mathbb{R}$, and to then use a kind of cellular approximation theorem to give a chain level approximation to the gradient flow in

$$(f_N, v_N) = (\overline{f}, \overline{v})| : (M_N, f_N, v_N) \rightarrow ([0, 1]; \{0\}, \{1\}) .$$

The mechanism described in Chapter 5 below then gives a chain complex over $S^{-1}\mathbb{Z}[z, z^{-1}]$ inducing $C^{Nov}(M, f, v)$. Hutchings and Lee [10],[11] used a similar method to get enough information from $C^{Nov}(M, f, v)$ for generic v to obtain an estimate on the number of closed v -gradient flow lines $\gamma : S^1 \rightarrow M$.

Farber and Ranicki [7] and Ranicki [31] constructed an ‘algebraic Novikov complex’ in $S^{-1}\mathbb{Z}[z, z^{-1}]$ for any circle Morse valued function $f : M \rightarrow S^1$, using any CW structure on

$N = f^{-1}(0)$, the extension to a CW structure on M_N , and a cellular approximation to the inclusion $z^{-1}N \rightarrow M_N$. The construction is recalled in Chapter 5, including the non simply connected version. In many cases (e.g. for $v \in \mathcal{GCCT}(f)$) this algebraic model does actually coincide with the geometric Novikov complex $C^{Nov}(M, f, v)$.

The Morse-Novikov theory of circle valued functions on finite-dimensional manifolds and Novikov homology have many applications to symplectic topology, Floer homology, and Seiberg-Witten theory (Poźniak [27], Le and Ono [14], Hutchings and Lee [10], [11], ...). Also, circle valued Morse theory on infinite-dimensional manifolds features in the work of Taubes on Casson's homology 3-sphere invariant and gauge theory. However, these notes are not a survey of all the applications of circle valued Morse theory and Novikov homology! They deal exclusively with the basic development in the finite-dimensional case and some of the applications to the classification of manifolds.

2 Real valued Morse theory

This section reviews the real valued Morse theory, which is a prerequisite for circle valued Morse theory. The traditional references Milnor [15], [16] remain the best introductions to real valued Morse theory. Bott [2] gives a beautiful account of the history of Morse theory, including the development of the modern chain complex point of view inspired by Witten.

Let M be a compact differentiable m -dimensional manifold. The *critical points* of a differentiable function $f : M \rightarrow \mathbb{R}$ are the zeros $p \in M$ of the differential $\nabla f : \tau_M \rightarrow \tau_{\mathbb{R}}$. A *Morse function* $f : M \rightarrow \mathbb{R}$ is a differentiable function in which every critical point $p \in M$ is required to be isolated and nondegenerate, meaning that in local coordinates

$$f(p + (x_1, x_2, \dots, x_m)) = f(p) - \sum_{j=1}^i (x_j)^2 + \sum_{j=i+1}^m (x_j)^2$$

with i the index of p . The subspace of Morse functions is C^2 -dense in the space of all differentiable functions $f : M \rightarrow \mathbb{R}$.

A vector field $v : M \rightarrow \tau_M$ is *gradient-like* for f if there exists a Riemannian metric $\langle \cdot, \cdot \rangle$ on M such that

$$\langle v, w \rangle = \nabla f(w) \in \mathbb{R} \quad (w \in \tau_M) .$$


Note that $\langle \cdot, \cdot \rangle$ and ∇f determine v , and that the zeros of v are the critical points of f .

A *v-gradient flow line* $\gamma : \mathbb{R} \rightarrow M$ satisfies

$$\gamma'(t) = -v(\gamma(t)) \in \tau_M(\gamma(t)) \quad (t \in \mathbb{R}) .$$

The minus sign here gives the *downward* gradient flow, so that

$$f(\gamma(s)) > f(\gamma(t)) \text{ if } s < t .$$



The limits

$$\lim_{t \rightarrow -\infty} \gamma(t) = p \quad , \quad \lim_{t \rightarrow \infty} \gamma(t) = q \in M$$

are critical points of f with $f(q) < f(p)$, and if γ is isolated then

$$\text{index}(q) = \text{index}(p) - 1 .$$

For every point $x \in M$ there is a v -gradient flow line $\gamma_x : \mathbb{R} \rightarrow M$ (which is unique up to scaling) such that $\gamma_x(0) = x \in M$. If x is a critical point take γ_x to be the constant path at x .

The *unstable* and *stable* manifolds of a critical point $p \in M$ of index i are the open manifolds

$$W^u(p, v) = \{x \in M \mid \lim_{t \rightarrow -\infty} \gamma_x(t) = p\} \quad , \quad W^s(p, v) = \{x \in M \mid \lim_{t \rightarrow \infty} \gamma_x(t) = p\} .$$


The unstable and stable manifolds are images of immersions $\mathbb{R}^i \rightarrow M$, $\mathbb{R}^{m-i} \rightarrow M$ respectively, which are embeddings near $p \in M$.

The basic results relating a Morse function $f : M^m \rightarrow \mathbb{R}$ to the topology of M concern the inverse images

$$N_a = f^{-1}(a)$$

of the regular values $a \in \mathbb{R}$, which are closed $(m-1)$ -dimensional manifolds, and the cobordisms

$$(M_{a,b}; N_a, N_b) = f^{-1}([a, b]; \{a\}, \{b\}) \quad (a < b) .$$



The results are:

- (i) if $[a, b] \subset \mathbb{R}$ contains no critical values the v -gradient flow determines a diffeomorphism

$$N_b \rightarrow N_a ; x \mapsto \gamma_x((f\gamma_x)^{-1}(a)) ,$$

- (ii) if $[a, b] \subset \mathbb{R}$ contains a unique critical value $f(p) \in (a, b)$, and $p \in M$ is a critical point of index i , then N_b is obtained from N_a by surgery on a tubular neighbourhood $S^{i-1} \times D^{m-i} \subset N_a$ of $S^{i-1} = W^u(p, v) \cap N_a$

$$N_b = N_a \setminus (S^{i-1} \times D^{m-i}) \cup D^i \times S^{m-i-1}$$

with $D^i \times S^{m-i-1} \subset N_b$ a tubular neighbourhood of $S^{m-i-1} = W^s(p, v) \cap N_b$, and $(M_{a,b}; N_a, N_b)$ the trace of the surgery

$$M_{a,b} = N_a \times [0, 1] \cup D^i \times D^{m-i} .$$

Let $\mathcal{GT}(f)$ denote the set of gradient-like vector fields v on M which satisfy the Morse-Smale transversality condition that for any critical points $p, q \in M$ with $\text{index}(p) = i$, $\text{index}(q) = j$ the submanifolds $W^u(p, v)^i$, $W^s(q, v)^{m-j} \subset M^m$ intersect transversely in an $(i - j)$ -dimensional submanifold $W^u(p, v) \cap W^s(q, v) \subset M$. The subspace $\mathcal{GT}(f)$ is dense in the space of gradient-like vector fields for f .

Suppose that the Morse function $f : M \rightarrow \mathbb{R}$ has $c_i(f)$ critical points of f of index i , and that the critical points $p_0, p_1, p_2, \dots \in M$ are arranged to satisfy

$$\text{index}(p_0) \leq \text{index}(p_1) \leq \text{index}(p_2) \leq \dots , f(p_0) < f(p_1) < f(p_2) < \dots .$$

A choice of $v \in \mathcal{GT}(f)$ determines a handle decomposition of M

$$M = \bigcup_{i=0}^m \bigcup_{c_i(f)} D^i \times D^{m-i}$$

with one i -handle $h^i = D^i \times D^{m-i}$ for each critical point of index i .

The *Morse-Smale complex* $C^{MS}(M, f, v)$ is defined for a Morse-Smale pair $(f : M \rightarrow \mathbb{R}, v \in \mathcal{GT}(f))$ and a regular cover \widetilde{M} of M with group of covering translations π , to be the based f.g. free $\mathbb{Z}[\pi]$ -module chain complex with

$$d : C^{MS}(M, f, v)_i = \mathbb{Z}[\pi]^{c_i(f)} \rightarrow C^{MS}(M, f, v)_{i-1} = \mathbb{Z}[\pi]^{c_{i-1}(f)} ; \tilde{p} \mapsto \sum_{\tilde{q}} n(\tilde{p}, \tilde{q}) \tilde{q}$$

with $n(\tilde{p}, \tilde{q}) \in \mathbb{Z}$ the finite signed number of \tilde{v} -gradient flow lines $\tilde{\gamma} : \mathbb{R} \rightarrow \widetilde{M}$ which start at a critical point $\tilde{p} \in \widetilde{M}$ of $\tilde{f} : \widetilde{M} \rightarrow \mathbb{R}$ with index i and terminate at a critical point $\tilde{q} \in \widetilde{M}$ of index $i - 1$. Choose an arbitrary lift of each critical point $p \in M$ of f to a critical point $\tilde{p} \in \widetilde{M}$ of \tilde{f} , obtaining a basis for $C^{MS}(M, f, v)$. The Morse-Smale complex is the cellular chain complex

$$C^{MS}(M, f, v) = C(\widetilde{M})$$

of the CW structure on \widetilde{M} in which the i -cells are the lifts of the i -handles h^i . In particular, the homology of the Morse-Smale complex is the ordinary homology of \widetilde{M}

$$H_*(C^{MS}(M, f, v)) = H_*(\widetilde{M}) .$$

If $(f, v) : M \rightarrow \mathbb{R}$ is modified to $(f', v') : M \rightarrow \mathbb{R}$ by adding a pair of critical points p, q of index $i, i-1$ with $n(p, q, v) = 1$ the Morse-Smale complex $C^{MS}(M, f', v')$ is obtained from $C^{MS}(M, f, v)$ by attaching an elementary chain complex

$$E : \cdots \rightarrow 0 \rightarrow E_i = \mathbb{Z}[\pi] \xrightarrow{1} E_{i-1} = \mathbb{Z}[\pi] \rightarrow 0 \rightarrow \cdots ,$$

with an exact sequence

$$0 \rightarrow C^{MS}(M, f, v) \rightarrow C^{MS}(M, f', v') \rightarrow E \rightarrow 0 .$$

Conversely, if $m \geq 5$ then the Whitney trick applies to realize the elementary moves of Whitehead torsion theory by cancellation of pairs of critical points (or equivalently, handles). This cancellation is the basis of the proofs of the h - and s -cobordism theorems.

The identity $C^{MS}(M, f, v) = C(M)$ (for $\widetilde{M} = M$) gives the *Morse inequalities*

$$c_i(f) \geq b_i(M) + q_i(M) + q_{i-1}(M)$$

with

$$b_i(M) = \dim_{\mathbb{Z}}(H_i(M)/T_i(M)) , \quad q_i(M) = \# T_i(M)$$

the Betti numbers of M , where

$$T_i(M) = \{x \in H_i(M) \mid nx = 0 \text{ for some } n \neq 0 \in \mathbb{Z}\}$$

is the torsion subgroup of $H_i(M)$ and $\#$ denotes the minimum number of generators. Smale used the cancellation of critical points to prove that these inequalities are sharp for $\pi_1(M) = \{1\}$, $m \geq 5$: there exists $(f, v) : M \rightarrow \mathbb{R}$ with the minimum possible number of critical points

$$c_i(f) = b_i(M) + q_i(M) + q_{i-1}(M) .$$

The method is to start with an arbitrary Morse function $f : M \rightarrow \mathbb{R}$, and to systematically cancel pairs of critical points until this is no longer possible.

The Morse-Smale complex $C^{MS}(M, f, v)$ is also defined for a Morse function on an m -dimensional cobordism $f : (M; N, N') \rightarrow ([0, 1]; \{0\}, \{1\})$ with $v \in \mathcal{GT}(f)$. In this case there is a relative handle decomposition

$$M = N \times [0, 1] \cup \bigcup_{i=0}^m \bigcup_{c_i(f)} D^i \times D^{m-i}$$

and $C^{MS}(M, f, v) = C(\widetilde{M}, \widetilde{N})$. The s -cobordism theorem states that for a Morse function f on an h -cobordism $\tau(C^{MS}(M, f, v)) = 0 \in Wh(\pi_1(M))$ if (and for $m \geq 6$ only if) the critical points of f can be stably cancelled in pairs.

3 The Novikov complex

Morse functions $f : M \rightarrow S^1$, gradient-like vector field v , critical points, index, $c_i(f)$, are defined in the same way as for the real valued case in Chapter 1. Again, the subspace of Morse functions is C^2 -dense in the space of all functions $f : M \rightarrow S^1$. But it is harder to decide which pairs of critical points can be cancelled.

A Morse function $f : M \rightarrow S^1$ lifts to a \mathbb{Z} -equivariant Morse function $\bar{f} : \bar{M} = f^*\mathbb{R} \rightarrow \mathbb{R}$ on the infinite cyclic cover

$$\begin{array}{ccc} \bar{M} & \longrightarrow & M \\ \bar{f} \downarrow & & \downarrow f \\ \mathbb{R} & \longrightarrow & S^1 \end{array}$$

Let $z : \bar{M} \rightarrow \bar{M}$ be the downward generating covering translation.

$$\begin{array}{c} \hline \leftarrow \overset{z}{\sim} \hline \bar{M} \\ \hline \downarrow \bar{f} \\ \hline \mathbb{R} \end{array}$$

Let $\mathcal{GT}(f)$ be the space of gradient-like vector fields $v : M \rightarrow \tau_M$ such that a lift $\bar{v} : \bar{M} \rightarrow \tau_{\bar{M}}$ satisfies the Morse-Smale transversality condition. The Novikov complex of a circle valued Morse function is defined by analogy with the Morse-Smale complex of a real valued function, as follows.

Given a ring A and an automorphism $\alpha : A \rightarrow A$ let z be an indeterminate over A with

$$az = z\alpha(a) \quad (a \in A).$$

The α -twisted Laurent polynomial extension of A is the localization of the α -twisted polynomial extension $A_\alpha[z]$ inverting z

$$A_\alpha[z, z^{-1}] = A_\alpha[z][z^{-1}],$$

the ring of polynomials $\sum_{j=-\infty}^{\infty} a_j z^j$ ($a_j \in A$) such that $\{j \in \mathbb{Z} \mid a_j \neq 0\}$ is finite.

The α -twisted Novikov ring of A is the localization of the completion of $A_\alpha[z]$

$$A_\alpha((z)) = A_\alpha[[z]][z^{-1}],$$

the ring of power series $\sum_{j=-\infty}^{\infty} a_j z^j$ ($a_j \in A$) such that $\{j \leq 0 \mid a_j \neq 0\}$ is finite.

Given $f : M \rightarrow S^1$ let \widetilde{M} be a regular cover of \overline{M} , with group of covering translations π . Only the case of connected $M, \overline{M}, \widetilde{M}$ will be considered. Let Π be the group of covering translations of \widetilde{M} over M , so that there is defined a group extension

$$\{1\} \rightarrow \pi \rightarrow \Pi \rightarrow \mathbb{Z} \rightarrow \{1\}$$

with a lift of $1 \in \mathbb{Z}$ to an element $z \in \Pi$ such that the covering translation $z : \widetilde{M} \rightarrow \widetilde{M}$ induces $z : \overline{M} \rightarrow \overline{M}$ on $\overline{M} = \widetilde{M}/\pi$. Thus

$$\Pi = \pi \times_{\alpha} \mathbb{Z} \quad , \quad \mathbb{Z}[\Pi] = \mathbb{Z}[\pi]_{\alpha}[z, z^{-1}] .$$

Write the α -twisted Novikov ring as

$$\widehat{\mathbb{Z}[\Pi]} = \mathbb{Z}[\pi]_{\alpha}((z)) .$$

Choose a lift of each critical point $p \in M$ of f to a critical point $\tilde{p} \in \widetilde{M}$ of \tilde{f} .

The Novikov complex $C^{Nov}(M, f, v)$ of $(f : M \rightarrow S^1, v \in \mathcal{GT}(f))$ is the based f.g. free $\widehat{\mathbb{Z}[\Pi]}$ -module chain complex with

$$d : C^{Nov}(M, f, v)_i = \mathbb{Z}[\pi]_{\alpha}((z))^{c_i(f)} \rightarrow C^{Nov}(M, f, v)_{i-1} = \mathbb{Z}[\pi]_{\alpha}((z))^{c_{i-1}(f)} ;$$

$$\tilde{p} \mapsto \sum_{j=-\infty}^{\infty} \sum_{\tilde{q}} n(\tilde{p}, z^j \tilde{q}) z^j \tilde{q}$$

with $n(\tilde{p}, \tilde{q}) \in \mathbb{Z}$ the finite signed number of \tilde{v} -gradient flow lines $\tilde{\gamma} : \mathbb{R} \rightarrow \widetilde{M}$ which start at a critical point $\tilde{p} \in \widetilde{M}$ of $\tilde{f} : \widetilde{M} \rightarrow \mathbb{R}$ with index i and terminate at a critical point $\tilde{q} \in \widetilde{M}$ of index $i - 1$.

Exercise. Work out $C^{Nov}(S^1, f, v)$ for

$$f : S^1 \rightarrow S^1 ; [t] \mapsto [4t - 9t^2 + 6t^3] \quad (0 \leq t \leq 1) . \quad \square$$

The original definition of Novikov [17],[18] was in the special case

$$\widetilde{M} = \overline{M} \quad , \quad \pi = \{1\} \quad , \quad \Pi = \mathbb{Z} \quad , \quad \alpha = 1 \quad , \quad \widehat{\mathbb{Z}[\Pi]} = \mathbb{Z}((z))$$


when $C^{Nov}(M, f, v)$ is a based f.g. free $\mathbb{Z}((z))$ -module chain complex (as in the Exercise).

Take \widetilde{M} to be the universal cover of M and $\pi = \pi_1(\overline{M})$, $\alpha : \pi \rightarrow \pi$ the automorphism induced by a generating covering translation $z : \overline{M} \rightarrow \overline{M}$, $\Pi = \pi_1(M) = \pi \times_{\alpha} \mathbb{Z}$. This case gives the based f.g. free $\widehat{\mathbb{Z}[\pi_1(M)]}$ -module Novikov complex $C^{Nov}(M, f, v)$ of Pajitnov [22].

There is only one class of Morse functions $f : M \rightarrow S^1$ for which the Novikov complex is easy to compute:

Example. Let M be the mapping torus of a diffeomorphism $h : N \rightarrow N$ of a closed $(m - 1)$ -dimensional manifold

$$M = T(h) = (N \times [0, 1]) / \{(x, 0) \sim (h(x), 1)\} .$$



The fibre bundle projection

$$f : M = T(h) \rightarrow S^1 = [0, 1]/\{0 \sim 1\} ; [x, t] \mapsto [t]$$

has no critical points, so that $C^{Nov}(M, f, v) = 0$ for any $v \in \mathcal{GT}(f)$. \square

4 Novikov homology

The Novikov homology $H_*^{Nov}(M, f; \widehat{\mathbb{Z}[\Pi]})$ is defined for a space M with a map $f : M \rightarrow S^1$ and a factorization of $f_* : \pi_1(M) \rightarrow \pi_1(S^1)$ through a group Π . The relevance of the Novikov complex $C^{Nov}(M, f, v)$ to the Morse theory of a Morse map $f : M \rightarrow S^1$ is immediately obvious. The relevance of the Novikov homology is rather less obvious, even though there are isomorphisms $H_*(C^{Nov}(M, f, v)) \cong H_*^{Nov}(M, f; \widehat{\mathbb{Z}[\Pi]})$!

The R -coefficient homology of a space M is defined for any ring morphism $\mathbb{Z}[\pi_1(M)] \rightarrow R$

$$H_*(M; R) = H_*(C(M; R))$$

using any free $\mathbb{Z}[\pi_1(M)]$ -module chain complex $C(\widetilde{M})$ (e.g. cellular, if M is a CW complex) and $C(M; R) = R \otimes_{\mathbb{Z}[\pi_1(M)]} C(\widetilde{M})$.

Given a group π and an automorphism $\alpha : \pi \rightarrow \pi$ let $\pi \times_{\alpha} \mathbb{Z}$ be the group with elements gz^j ($g \in \pi$, $j \in \mathbb{Z}$), and multiplication by $gz = \alpha(g)z$, so that

$$\mathbb{Z}[\pi \times_{\alpha} \mathbb{Z}] = \mathbb{Z}[\pi]_{\alpha}[z, z^{-1}] .$$

For any map $f : M \rightarrow S^1$ with M connected the infinite cyclic cover $\overline{M} = f^*\mathbb{R}$ is connected if and only if $f_* : \pi_1(M) \rightarrow \pi_1(S^1) = \mathbb{Z}$ is onto, in which case

$$\pi_1(M) = \pi_1(\overline{M}) \times_{\alpha_M} \mathbb{Z}$$

with $\alpha_M : \pi_1(\overline{M}) \rightarrow \pi_1(\overline{M})$ the automorphism induced by a generating covering translation $z : \overline{M} \rightarrow \overline{M}$.

Suppose given a connected space M with a cohomology class $f \in [M, S^1] = H^1(M)$ such that $\overline{M} = f^*\mathbb{R}$ is connected. Given a factorization of the surjection $f_* : \pi_1(M) \rightarrow \pi_1(S^1)$

$$f_* : \pi_1(M) = \pi_1(\overline{M}) \times_{\alpha_M} \mathbb{Z} \rightarrow \Pi \rightarrow \mathbb{Z}$$

let $\pi = \ker(\Pi \rightarrow \mathbb{Z})$ and let $z \in \Pi$ be the image of $z = (0, 1) \in \pi_1(M)$, so that $\Pi = \pi \times_\alpha \mathbb{Z}$ with

$$\alpha : \pi \rightarrow \pi ; g \mapsto z^{-1}gz .$$

The $\widehat{\mathbb{Z}[\Pi]}$ -coefficient *Novikov homology* of (M, f) is

$$H_*^{Nov}(M, f; \widehat{\mathbb{Z}[\Pi]}) = H_*(M; \widehat{\mathbb{Z}[\Pi]}) ,$$

with $\widehat{\mathbb{Z}[\Pi]} = \mathbb{Z}[\pi]_\alpha((z))$.

In the original case

$$\widetilde{M} = \overline{M} , \pi = \{1\} , \Pi = \mathbb{Z} , \widehat{\mathbb{Z}[\Pi]} = \mathbb{Z}((z)) ,$$

and $H_*^{Nov}(M, f; \widehat{\mathbb{Z}[\Pi]})$ may be written as $H_*^{Nov}(M, f)$, or even just $H_*^{Nov}(M)$.

Example 1. The $\mathbb{Z}((z))$ -coefficient cellular chain complex of S^1 is

$$C(S^1; \mathbb{Z}((z))) : \dots \rightarrow 0 \rightarrow \mathbb{Z}((z)) \xrightarrow{1-z} \mathbb{Z}((z))$$

and $1 - z \in \mathbb{Z}((z))$ is a unit, so $H_*^{Nov}(S^1) = 0$. □

Example 2. Let N be a connected finite *CW* complex with cellular \mathbb{Z} -module chain complex $C(N)$, and let $h : N \rightarrow N$ be a self-map with induced \mathbb{Z} -module chain map $h : C(N) \rightarrow C(N)$. The $\mathbb{Z}((z))$ -coefficient cellular chain complex of the mapping torus $T(h)$ with respect to the canonical projection

$$f : T(h) \rightarrow S^1 ; [x, t] \mapsto [t]$$

is the algebraic mapping cone

$$C^+(T(h); \mathbb{Z}((z))) = \mathcal{C}(1 - zh : C(N)((z)) \rightarrow C(N)((z))) .$$

Now $1 - zh$ is a $\mathbb{Z}((z))$ -module chain equivalence, so that

$$H_*^{Nov}(T(h), f) = 0 .$$

The $\mathbb{Z}((z))$ -coefficient cellular chain complex of the mapping torus $T(h)$ with respect to the other projection

$$-f : T(h) \rightarrow S^1 ; [x, t] \mapsto [1 - t]$$

is the algebraic mapping cone

$$C^-(T(h); \mathbb{Z}((z))) = \mathcal{C}(z - h : C(N)((z)) \rightarrow C(N)((z))) .$$

If $h : N \rightarrow N$ is a homotopy equivalence then $z - h$ is a $\mathbb{Z}((z))$ -module chain equivalence, so that

$$H_*^{Nov}(T(h), -f) = 0 ,$$

but in general $H_*^{Nov}(T(h), -f) \neq 0$ – see Example 3 below for an explicit non-zero example. □

Example 3. The Novikov homology groups of the mapping torus $T(2)$ of the double covering map $2 : S^1 \rightarrow S^1$ are

$$\begin{aligned} H_1^{Nov}(T(2), f) &= \mathbb{Z}((z))/(1-2z) = 0, \\ H_1^{Nov}(T(2), -f) &= \mathbb{Z}((z))/(z-2) = \widehat{\mathbb{Z}}_2[1/2] = \widehat{\mathbb{Q}}_2 \neq 0 \end{aligned}$$

with $\widehat{\mathbb{Q}}_2$ the 2-adic field (Example 23.25 of Hughes and Ranicki [9]). The inverse of

$$n = 2^a(2b+1) \in \mathbb{Z}$$

is

$$n^{-1} = z^{-a}(1 - zb + z^2b^2 - z^3b^3 + \dots) \in \mathbb{Z}((z))/(2-z) = \widehat{\mathbb{Q}}_2. \quad \square$$

Theorem. (Novikov [17], [18] for $\pi = \{1\}$, Pajitnov [21])

The Novikov complex $C^{Nov}(M, f, v)$ is $\widehat{\mathbb{Z}[\Pi]}$ -module chain equivalent to $C(M; \widehat{\mathbb{Z}[\Pi]})$, with isomorphisms

$$H_*(C^{Nov}(M, f, v)) \cong H_*^{Nov}(M, f; \widehat{\mathbb{Z}[\Pi]}). \quad \square$$

The chain equivalence $C^{Nov}(M, f, v) \simeq C(M; \widehat{\mathbb{Z}[\Pi]})$ will be described in Chapter 4 below.

The Novikov ring $\mathbb{Z}((z))$ is a principal ideal domain, and $H_*^{Nov}(M, f)$ is the homology of a f.g. free $\mathbb{Z}((z))$ -module chain complex. Thus each $H_i^{Nov}(M, f)$ is a f.g. $\mathbb{Z}((z))$ -module, which splits as free \oplus torsion, by the structure theorem for f.g. modules over a principal ideal domain.

The *Novikov numbers* of any finite CW complex M and $f \in H^1(M)$ are the Betti numbers of Novikov homology

$$b_i^{Nov}(M, f) = \dim_{\mathbb{Z}((z))}(H_i^{Nov}(M, f)/T_i^{Nov}(M, f)), \quad q_i^{Nov}(M, f) = \#T_i^{Nov}(M, f)$$

where

$$T_i^{Nov}(M, f) = \{x \in H_i^{Nov}(M, f) \mid ax = 0 \text{ for some } a \neq 0 \in \mathbb{Z}((z))\}$$

is the torsion $\mathbb{Z}((z))$ -submodule of $H_i^{Nov}(M, f)$, and $\#$ denotes the minimum number of generators.

The *Morse-Novikov inequalities* ([17])

$$c_i(f) \geq b_i^{Nov}(M, f) + q_i^{Nov}(M, f) + q_{i-1}^{Nov}(M, f)$$

are an immediate consequence of the isomorphisms $H_*(C^{Nov}(M, f, v)) \cong H_*^{Nov}(M, f)$, since for any f.g. free chain complex C over a principal ideal domain R

$$\dim_R(C_i) \geq b_i(C) + q_i(C) + q_{i-1}(C)$$

where

$$b_i(C) = \dim_R(H_i(C)/T_i(C)), \quad q_i(C) = \#T_i(C)$$

with

$$T_i(C) = \{x \in H_i(C) \mid rx = 0 \text{ for some } r \neq 0 \in R\}$$

the R -torsion submodule of $H_i(C)$, and $\#$ denoting the minimal number of R -module generators.

Farber [5] proved that the Morse-Novikov inequalities are sharp for $\pi_1(M) = \mathbb{Z}$, $m \geq 6$: for any such manifold there exists a Morse function $f : M \rightarrow S^1$ representing $1 \in [M, S^1] = H^1(M)$ with the minimum possible numbers of critical points

$$c_i(f) = b_i^{Nov}(M, f) + q_i^{Nov}(M, f) + q_{i-1}^{Nov}(M, f) .$$

Again, the method is to start with an arbitrary Morse function $f : M \rightarrow S^1$ in the homotopy class, and to systematically cancel pairs of critical points until this is no longer possible.

When does the Novikov homology vanish?

Proposition (Ranicki [29]) Let A be a ring with an automorphism $\alpha : A \rightarrow A$. A finite f.g. free $A_\alpha[z, z^{-1}]$ -module chain complex C is such that

$$H_*(A_\alpha((z)) \otimes_{A_\alpha[z, z^{-1}]} C) = H_*(A_\alpha((z^{-1})) \otimes_{A_\alpha[z, z^{-1}]} C) = 0$$

if and only if C is A -module chain equivalent to a finite f.g. projective A -module chain complex. \square

Note that for an algebraic Poincaré complex (C, ϕ)

$$H_*(A_\alpha((z)) \otimes_{A_\alpha[z, z^{-1}]} C) = 0 \text{ if and only if } H_*(A_\alpha((z^{-1})) \otimes_{A_\alpha[z, z^{-1}]} C) = 0 ,$$

so the two Novikov homology vanishing conditions can be replaced by just one.

Recall that a space X is *finitely dominated* if there exist a finite CW complex and maps $i : X \rightarrow K$, $j : K \rightarrow X$ such that $ji \simeq 1 : X \rightarrow X$. Wall [37] proved that a CW complex X is finitely dominated if and only if $\pi_1(X)$ is finitely presented and the cellular chain complex $C(\tilde{X})$ of the universal cover \tilde{X} is chain equivalent to a finite f.g. projective $\mathbb{Z}[\pi_1(X)]$ -module chain complex.

In the simply-connected case $\pi_1(\overline{M}) = \{1\}$ the following conditions on a map $f : M \rightarrow S^1$ from an m -dimensional manifold M are equivalent :

- (i) \overline{M} is finitely dominated,
- (ii) \overline{M} is homotopy equivalent to a finite CW complex,
- (iii) $H_*^{Nov}(M, f) = 0$,
- (iv) $b_i^{Nov}(M, f) = q_i^{Nov}(M, f) = 0$,
- (v) $C(\overline{M})$ is chain equivalent to a finite f.g. free \mathbb{Z} -module chain complex,
- (vi) the homology groups $H_*(\overline{M})$ are f.g. \mathbb{Z} -modules.

Browder and Levine [3] used handle exchanges (= the ambient surgery version of the cancellation of adjacent critical points) to prove that (vi) holds if (and for $m \geq 6$ only if) $f : M \rightarrow S^1$ is homotopic to the projection of a fibre bundle.

Farrell [8] and Siebenmann [35] defined a Whitehead torsion obstruction $\Phi(M, f) \in Wh(\pi_1(M))$ for a map $f : M^m \rightarrow S^1$ with finitely dominated $\overline{M} = f^*\mathbb{R}$, such that $\Phi(M, f) = 0$ if (and for $m \geq 6$ only if) f is homotopic to the projection of a fibre bundle.

Theorem (Ranicki [29])

(i) For any finite CW complex M and map $f : M \rightarrow S^1$ the infinite cyclic cover $\overline{M} = f^*\mathbb{R}$ of M is finitely dominated if and only if $\pi_1(\overline{M})$ is finitely presented and

$$H_*^{Nov}(M, f; \mathbb{Z}[\widehat{\pi_1(M)}]) = 0 .$$

(ii) For any Morse map $f : M \rightarrow S^1$ on an m -dimensional manifold M with finitely dominated \overline{M} the torsion of the Novikov complex $\tau(C^{Nov}(M, f, v)) \in K_1(\mathbb{Z}[\widehat{\pi_1(M)}])/I$ determines and is determined by the Farrell-Siebenmann fibering obstruction $\Phi(M, f) \in Wh(\pi_1(M))$, where $I \subseteq K_1(\mathbb{Z}[\widehat{\pi_1(M)}])$ is the subgroup generated by $\pm\pi_1(M)$ and $\tau(1 - zh)$ for square matrices h over $\mathbb{Z}[\widehat{\pi_1(M)}]$. Thus $\tau(C^{Nov}(M, f, v)) \in I$ if (and for $m \geq 6$ only if) f is homotopic to a fibre bundle. \square

See Chapter 22 of Hughes and Ranicki [9] and Chapter 15 of Ranicki [30] for more detailed accounts of the relationship between the torsion of the Novikov complex and the Farrell-Siebenmann fibering obstruction.

See Latour [12] and Pajitnov [22] for circle Morse-theoretic proofs that if $m \geq 6$, \overline{M} is finitely dominated and $\tau(C^{Nov}(M, f, v)) \in I$ then it is possible to pairwise cancel all the critical points of f .

5 The algebraic model for circle valued Morse theory

In many cases the Novikov complex $C^{Nov}(M, f, \alpha)$ of a circle valued Morse function $f : M \rightarrow S^1$ can be constructed from an algebraic model for the \bar{v} -gradient flow in a fundamental domain of the infinite cyclic cover \overline{M} .

An *algebraic fundamental domain* (D, E, F, g, h) consists of finite based f.g. free A -module chain complexes D, E and chain maps $g : D \rightarrow E, h : z^{-1}D \rightarrow E$ of the form

$$\begin{aligned} d_E &= \begin{pmatrix} d_D & c \\ 0 & d_F \end{pmatrix} : E_i = D_i \oplus F_i \rightarrow E_{i-1} = D_{i-1} \oplus F_{i-1} , \\ g &= \begin{pmatrix} 1 \\ 0 \end{pmatrix} : D_i \rightarrow E_i = D_i \oplus F_i , \\ h &= \begin{pmatrix} h_D \\ h_F \end{pmatrix} : z^{-1}D_i \rightarrow E_i = D_i \oplus F_i . \end{aligned}$$

zD	zE	D	\xrightarrow{g}	E	\xleftarrow{zh}	$z^{-1}D$	$z^{-1}E$	$z^{-2}D$
------	------	-----	-------------------	-----	-------------------	-----------	-----------	-----------

Define the *algebraic Novikov complex* \widehat{F} to be the based f.g. free $A_\alpha((z))$ -module chain complex with

$$\begin{aligned} d_{\widehat{F}} &= d_F + zh_F(1 - zh_D)^{-1}c \\ &= d_F + \sum_{j=1}^{\infty} z^j h_F (h_D)^{j-1} c : \widehat{F}_i = (F_i)_\alpha((z)) \rightarrow \widehat{F}_{i-1} = (F_{i-1})_\alpha((z)), \end{aligned}$$

as in Farber and Ranicki [7] and Ranicki [31]. The $A_\alpha((z))$ -module chain map


$$\phi = g - zh = \begin{pmatrix} 1 - zh_D \\ -zh_F \end{pmatrix} : D_\alpha((z)) \rightarrow E_\alpha((z))$$

is a split injection in each degree (since $1 - zh_D$ is an isomorphism), and the inclusions $F_i \rightarrow E_i$ determine a canonical isomorphism of based f.g. free $A_\alpha[z, z^{-1}]$ -module chain complexes

$$\widehat{F} \cong \text{coker}(\phi).$$

Here is how algebraic fundamental domains and the algebraic Novikov complex arise in topology.

Let $f : M \rightarrow S^1$ be a Morse function with regular value $0 \in S^1$.



Cut M along the inverse image

$$N^{m-1} = f^{-1}(0) \subset M$$

to obtain a geometric fundamental domain

$$(M_N; N, z^{-1}N) = \overline{f}^{-1}([0, 1]; \{0\}, \{1\})$$


for the infinite cyclic cover

$$\overline{M} = f^*\mathbb{R} = \bigcup_{j=-\infty}^{\infty} z^j M_N.$$

The restriction

$$f_N = \overline{f}| : (M_N; N, z^{-1}N) \rightarrow ([0, 1]; \{0\}, \{1\})$$

is a real valued Morse function with $v_N = \bar{v} \in \mathcal{GT}(f_N)$.



The cobordism $(M_N; N, z^{-1}N)$ has a handlebody decomposition

$$M_N = N \times [0, 1] \cup \bigcup_{i=0}^m \bigcup_{c_i(f)} D^i \times D^{m-i}$$

with one i -handle for each index i critical point of f . Given a CW structure on N with $c_i(N)$ i -cells use this handlebody decomposition to define a CW structure on M_N with $c_i(N) + c_i(f)$ i -cells. A regular cover \widetilde{M} of \overline{M} with group of covering translations π is a regular cover of M with group of covering translations $\Pi = \pi \times_{\alpha} \mathbb{Z}$ (as before), with

$$\mathbb{Z}[\Pi] = \mathbb{Z}[\pi]_{\alpha}[z, z^{-1}] \quad , \quad \widehat{\mathbb{Z}[\Pi]} = \mathbb{Z}[\Pi]_{\alpha}((z)) \quad .$$

Use a cellular approximation $h : z^{-1}N \rightarrow M_N$ to the inclusion to define an algebraic fundamental domain (D, E, F, g, h) over $A = \mathbb{Z}[\pi]$

$$D = C(\widetilde{N}) \quad , \quad E = C(\widetilde{M}_N) \quad , \quad F = C^{MS}(M_N, f_N, v_N) = C(\widetilde{M}_N, \widetilde{N}) \quad .$$

The mapping cylinder of $h : N \rightarrow M_N$ is a CW complex M'_N with two copies of N as subcomplexes. Identifying these copies there is obtained a CW complex structure on M with $\widehat{\mathbb{Z}[\Pi]}$ -coefficient cellular chain complex

$$C(M; \widehat{\mathbb{Z}[\Pi]}) = \mathcal{C}(\phi)$$

the algebraic mapping cone of the $\widehat{\mathbb{Z}[\Pi]}$ -module chain map

$$\phi = g - zh : D_{\alpha}((z)) \rightarrow E_{\alpha}((z)) \quad ,$$

with


$$d_{\mathcal{C}(\phi)} = \begin{pmatrix} -d_D & 0 & 0 \\ 1 - zh_D & d_D & c \\ -zh_F & 0 & d_D \end{pmatrix} \quad :$$

$$\mathcal{C}(\phi)_i = (D_{i-1} \oplus D_i \oplus F_i)_{\alpha}[z, z^{-1}] \rightarrow \mathcal{C}(\phi)_{i-1} = (D_{i-2} \oplus D_{i-1} \oplus F_{i-1})_{\alpha}[z, z^{-1}] \quad .$$

The algebraic Novikov complex $\widehat{F} = \text{coker}(\phi)$ is a based f.g. free $\widehat{\mathbb{Z}[\Pi]}$ -module chain complex such that

$$\dim_{\widehat{\mathbb{Z}[\Pi]}} \widehat{F}_i = c_i(f) .$$

In many cases $\widehat{F} = C^{Nov}(M, f, v)$, and in even more cases \widehat{F} is simple isomorphic to $C^{Nov}(M, f, v)$.



The philosophy here is that $\mathcal{C}(\phi)$ counts the \bar{v} -gradient flow lines of $\bar{f} : \bar{M} \rightarrow \mathbb{R}$, as follows :

- (i) the $(z^{-1}p, q)$ -coefficient of $h_D : z^{-1}D_i \rightarrow D_i$ counts the number of portions in M_N of the \bar{v} -gradient flow lines which start in $z^{-1}M_N$, enter M_N at $z^{-1}p \in z^{-1}N$, exit at $q \in N$ and end in zM_N ,
- (ii) the $(z^{-1}p, q)$ -coefficient of $h_F : z^{-1}D_i \rightarrow F_i$ counts the number of portions in M_N of the \bar{v} -gradient flow lines which start in $z^{-1}M_N$, enter M_N at $z^{-1}p \in z^{-1}N$ and end at $q \in M_N$,
- (iii) the (p, q) -coefficient of $c : F_i \rightarrow D_{i-1}$ counts the number of portions in M_N of the \bar{v} -gradient flow lines which start at $p \in M_N$, exit at $q \in N$, and end in zM_N .

Then for $j = 1, 2, 3, \dots$ the $(p, z^j q)$ -coefficient of $h_F(h_D)^{j-1}c : F_i \rightarrow z^j F_i$ is the number of the \bar{v} -gradient flow lines which start at $p \in M_N$ and end at $z^j q \in z^j M_N$, crossing the walls $N, zN, \dots, z^{j-1}N$. If such is the case, i.e. if the chain map h is *gradient-like* in the terminology of Ranicki [31], this is just the $(p, z^j q)$ -coefficient of $d_{C^{Nov}(M, f, v)}$, so $\widehat{F} = C^{Nov}(M, f, \alpha)$. Pajitnov [24] constructed a C^0 -dense subspace $\mathcal{GECT}(f) \subset \mathcal{GT}(f)$ of gradient-like vector fields v for which there exist a CW structure N and a gradient-like chain map h . Cornea and Ranicki [4] construct for any $v \in \mathcal{GT}(f)$ a Morse function $f' : M \rightarrow S^1$ arbitrarily close to f with $v' \in \mathcal{GT}(f')$ such that

$$C^{Nov}(M, f', v') = \mathcal{C}(\phi) .$$

The projection

$$p : C(M; \widehat{\mathbb{Z}[\Pi]}) = \mathcal{C}(\phi) \rightarrow \text{coker}(\phi) \cong \widehat{F}$$

is a chain equivalence of based f.g. free $\widehat{\mathbb{Z}[\Pi]}$ -module chain complexes, with torsion

$$\tau(p) = \sum_{i=0}^{\infty} (-)^i \tau(1 - zh_D : (D_i)_\alpha((z)) \rightarrow (D_i)_\alpha((z))) \in K_1(\widehat{\mathbb{Z}[\Pi]}) .$$

If h is a gradient-like chain map the torsion of p is a measure of the number of closed orbits of the v -gradient flow in M , i.e. the closed flow lines $\gamma : S^1 \rightarrow M$ (Hutchings and Lee [10],[11], Pajitnov [24],[26], Schütz [32],[33]).

The algebraic surgery treatment of high-dimensional knot theory in Ranicki [30] gives the following algebraic model for the circle valued Morse function on a knot complement.

Example. Let $k : S^n \subset S^{n+2}$ be a knot with $\pi_1(S^{n+2} \setminus k(S^n)) = \mathbb{Z}$. The complement of a tubular neighbourhood $k(S^n) \times D^2 \subset S^{n+2}$ is an $(n+2)$ -dimensional manifold with boundary

$$(M, \partial M) = (\text{cl.}(S^{n+2} \setminus (k(S^n) \times D^2)), k(S^n) \times S^1)$$

with

$$\pi_1(M) = \mathbb{Z} , \pi_1(\overline{M}) = \{1\} , H_*(M) = H_*(S^1) .$$

Let $f : (M, \partial M) \rightarrow S^1$ be a map representing $1 \in H^1(M) = \mathbb{Z}$, with $f| : \partial M \rightarrow S^1$ the projection. Making f transverse regular at $0 \in S^1$ there is obtained a Seifert surface $N^{n+1} = f^{-1}(0) \subset M$ for k , with $\partial N = k(S^n)$. As before, cut M along N to obtain a fundamental domain $(M_N; N, z^{-1}N)$ for the infinite cyclic cover $\overline{M} = f^*\mathbb{R}$ of M . For any CW structures on N, M_N write the reduced chain complexes as

$$\dot{C}(N) = C(N, \{\text{pt.}\}) , \dot{C}(M_N) = C(M_N, \{\text{pt.}\}) .$$

The inclusions $G : N \rightarrow M_N, H : z^{-1}N \rightarrow M_N$ induce \mathbb{Z} -module chain maps

$$G : \dot{C}(N) \rightarrow \dot{C}(M_N) , H : z^{-1}\dot{C}(N) \rightarrow \dot{C}(M_N)$$

such that $G - H : \dot{C}(N) \rightarrow \dot{C}(M_N)$ is a chain equivalence. The chain map

$$e = (G - H)^{-1}G : \dot{C}(N) \rightarrow \dot{C}(N)$$

is a generalization of the Seifert matrix, such that up to \mathbb{Z} -module chain homotopy

$$1 - e = -(G - H)^{-1}H : \dot{C}(N) \rightarrow \dot{C}(N)$$

and such that there is defined a $\mathbb{Z}[z, z^{-1}]$ -module chain equivalence

$$C(\overline{M}, \mathbb{R}) \simeq C(e + z(1 - e) : \dot{C}(N)[z, z^{-1}] \rightarrow \dot{C}(N)[z, z^{-1}]) .$$

Let $\dot{N} = \text{cl.}(N \setminus D^{n+1})$, for any embedding $D^{n+1} \subset N \setminus \partial N$. For any handlebody decomposition of the $(n+1)$ -dimensional cobordism $(\dot{N}; k(S^n), S^n)$ with $c_i(N)$ i -handles

$$\dot{N} = k(S^n) \times [0, 1] \cup \bigcup_{i=1}^n \bigcup_{c_i(N)} D^i \times D^{n+1-i}$$

there exists a Morse function $f : M \rightarrow S^1$ in the homotopy class $1 \in [M, S^1] = H^1(M) = \mathbb{Z}$ with

$$c_i(f) = c_i(N) + c_{i-1}(N)$$

critical points of index i . In this case the algebraic model for $C^{Nov}(M, f, v)$ has

$$\begin{aligned} D &= C(N) = \mathbb{Z} \oplus \dot{D}, \quad \dot{D}_i = \mathbb{Z}^{c_i(N)}, \\ F &= C^{MS}(M_N, f_N, v_N) = \mathcal{C}(e : \dot{D} \rightarrow \dot{D}), \\ d_F &= \begin{pmatrix} d_{\dot{D}} & e \\ 0 & -d_{\dot{D}} \end{pmatrix} : F_i = \dot{D}_i \oplus \dot{D}_{i-1} \rightarrow F_{i-1} = \dot{D}_{i-1} \oplus \dot{D}_{i-2}, \\ c &= (0 \ 1) : F_i = \dot{D}_i \oplus \dot{D}_{i-1} \rightarrow D_{i-1}, \\ h_D &= 0 : z^{-1}D_i \rightarrow D_i, \\ h_F &= \begin{pmatrix} 1-e \\ 0 \end{pmatrix} : z^{-1}D_i \rightarrow F_i = \dot{D}_i \oplus \dot{D}_{i-1} \end{aligned}$$

with algebraic Novikov complex

$$\begin{aligned} d_{\widehat{F}} &= d_F + \sum_{j=1}^{\infty} z^j h_F (h_D)^{j-1} c \\ &= \begin{pmatrix} d_{\dot{D}} & e + z(1-e) \\ 0 & -d_{\dot{D}} \end{pmatrix} : \widehat{F}_i = (\dot{D}_i \oplus \dot{D}_{i-1})(z) \rightarrow \widehat{F}_{i-1} = (\dot{D}_{i-1} \oplus \dot{D}_{i-2})(z) \end{aligned}$$

such that $H_*(\widehat{F}) = H_*^{Nov}(M, f)$. The short exact sequences of $\mathbb{Z}((z))$ -modules

$$0 \rightarrow H_i(N)((z)) \xrightarrow{e+z(1-e)} H_i(N)((z)) \rightarrow H_i^{Nov}(M, f) \rightarrow 0$$

can be used to express the Novikov numbers $b_i^{Nov}(M, f), q_i^{Nov}(M, f)$ of the knot complement in terms of the Alexander polynomials

$$\Delta_i(z) = \det(e + z(1-e) : H_i(N)[z, z^{-1}] \rightarrow H_i(N)[z, z^{-1}]) \in \mathbb{Z}[z, z^{-1}] \quad (1 \leq i \leq n),$$

generalizing the case $n = 1$ due to Lazarev [13]. For $n \geq 4$ and $\pi_1(M) = \mathbb{Z}$ the following conditions are equivalent :

- (i) the knot fibres, i.e. $f : M \rightarrow S^1$ is homotopic to the projection of a fibre bundle, with no critical points,
- (ii) $b_*^{Nov}(M, f) = q_*^{Nov}(M, f) = 0$, i.e. $H_*^{Nov}(M, f) = 0$,
- (iii) the constant and leading coefficients of $\Delta_*(z)$ are $\pm 1 \in \mathbb{Z}$. □

There is also a more refined version of the algebraic model for circle valued Morse theory, using the noncommutative Cohn localization $\Sigma^{-1}A_\alpha[z, z^{-1}]$ of $A_\alpha[z, z^{-1}]$ inverting the set Σ of square matrices of the form $1 - zh$ for a square matrix h over A . Indeed, the formula for the differentials in the algebraic Novikov complex

$$d_{\widehat{F}} = d_F + zh_F(1 - zh_D)^{-1}c$$

is already defined in $\Sigma^{-1}A_\alpha[z, z^{-1}]$. See Farber and Ranicki [7] and Ranicki [31] for further details of the construction. Farber [6] applied the refinement to obtain improvements of the Morse-Novikov inequalities, using homology with coefficients in flat line bundles instead of Novikov homology. It should be noted that the natural morphism $\Sigma^{-1}A_\alpha[z, z^{-1}] \rightarrow A_\alpha((z))$ is injective for commutative A with $\alpha = 1$, but it is not injective in general (Sheiham [34]).

References

- [1] V. Arnold, *Dynamics of intersection*, Proc. Conf. in Honour of J. Moser, Academic Press, 77–84 (1990)
- [2] R. Bott, *Morse theory indomitable*, Pub. Math. I.H.E.S. 68, 99–114 (1989)
- [3] W. Browder and J. Levine, *Fibering manifolds over the circle*, Comm. Math. Helv. 40, 153–160 (1966)
- [4] O. Cornea and A.A. Ranicki, *Rigidity and glueing for the Morse and Novikov complexes*, e-print <http://arXiv.org/abs/math.AT/0107221>
- [5] M. Farber, *Exactness of the Novikov inequalities*, Functional analysis and its applications 19, 49–59 (1985)
- [6] ———, *Morse-Novikov critical point theory, Cohn localization and Dirichlet units*, e-print <http://arXiv.org/abs/dg-ga/9911157>, Commun. Contemp. Math 1 (1999), 467–495
- [7] ——— and A.A. Ranicki, *The Morse-Novikov theory of circle valued functions and noncommutative localization*, e-print <http://arXiv.org/abs/dg-ga/9812122>, Proc. 1998 Moscow Conference for the 60th Birthday of S. P. Novikov, Tr. Mat. Inst. Steklova 225, 381–388 (1999)
- [8] F.T. Farrell, *The obstruction to fibering a manifold over a circle*, Indiana Univ. J. 21, 315–346 (1971)
- [9] B. Hughes and A.A. Ranicki, *Ends of complexes*, Tracts in Mathematics 123, Cambridge (1996)
- [10] M. Hutchings and Y.-J. Lee, *Circle valued Morse theory and Reidemeister torsion*, e-print <http://arXiv.org/abs/dg-ga/9706012>, Geom. Top. 3, 369–396 (1999)
- [11] ——— and ———, *Circle-valued Morse theory, Reidemeister torsion, and Seiberg-Witten invariants of 3-manifolds*, Topology 38, 861–888 (1999)
- [12] F. Latour, *Existence de 1-formes fermées non singulières dans une classe de cohomologie de de Rham*, Publ. Math. I.H.E.S. 80, 135–193 (1994)
- [13] A.Y. Lazarev, *Novikov homologies in knot theory*, Mat. Zam. 51, 53–57 (1992), English tr. : Math. Notes 51, 259–262 (1992)
- [14] V.H. Le and K. Ono, *Symplectic fixed points, the Calabi invariant and Novikov homology*, Topology 34, 155–176 (1995)
- [15] J. Milnor, *Morse theory*, Annals of Math. Studies 51, Princeton (1963)
- [16] ———, *Lectures on the h-cobordism theorem*, Mathematical Notes 1, Princeton (1965)

- [17] S.P. Novikov, *Multivalued functions and functionals. An analogue of the Morse theory*, Soviet Math. Dokl. 24, 222–226 (1981)
- [18] ———, *The hamiltonian formalism and a multi-valued analogue of Morse theory*, Uspeki Mat. 37, 3–49 (1982). English tr. Russian Math. Surveys 37, 1–56 (1982).
- [19] ———, *Quasiperiodic structures in topology*, in *Topological methods in modern mathematics* (Proc. Milnor 60th Birthday Symposium), 223–233, Publish or Perish (1993)
- [20] ———, *Topology I.*, Encyclopaedia of Math. Sci. 12, Springer (1996)
- [21] A.V. Pajitnov, *On the Novikov complex for rational Morse forms*, Annales de la Faculté de Sciences de Toulouse 4, 297–338 (1995)
- [22] ———, *Surgery on the Novikov Complex*, *K-theory* 10, 323–412 (1996)
- [23] ———, *The incidence coefficients in the Novikov complex are generically rational functions*, *Algebra i Analiz* 9, 92–139 (1997). English tr. St. Petersburg Math. J. 9, 969–1006 (1998)
- [24] ———, *Simple homotopy type of Novikov complex for closed 1-forms and Lefschetz ζ -function of the gradient flow*, e-print <http://arXiv.org/abs/dg-ga/9706014>, *Russian Math. Surveys* 54, 117–170 (1999)
- [25] ———, *Closed orbits of gradient flows and logarithms of non-abelian Witt vectors*, e-print <http://arXiv.org/abs/math.DG/9908010>, *K-theory* 21, 301–324 (2000)
- [26] ———, *Counting closed orbits of gradient flows of circle valued maps*, e-print math.AT/0104273
- [27] M. Poźniak, *Floer homology, Novikov rings and clean intersections*, A.M.S. Translations 196, 119–181 (1999)
- [28] A.A. Ranicki, *The algebraic theory of surgery*, Proc. Lond. Math. Soc. (3) 40, 87–192, 193–283 (1980)
- [29] ———, *Finite domination and Novikov rings*, *Topology* 34, 619–632 (1995)
- [30] ———, *High dimensional knot theory*, Springer Mathematical Monograph, Springer (1998)
- [31] ———, *The algebraic construction of the Novikov complex*, to appear in *Mathematische Annalen*, e-print <http://arXiv.org/abs/math.AT/9903090>
- [32] D. Schütz, *Gradient flows of 1-forms and their closed orbits*, e-print <http://arXiv.org/abs/math.DG/0009055>, to appear in *Forum Math.*
- [33] ———, *One parameter fixed point theory and gradient flows of closed 1-forms*, e-print <http://arXiv.org/abs/math.DG/0104245>, to appear in *K-theory*

- [34] D. Sheiham, *Noncommutative characteristic polynomials and Cohn localisation*, e-print <http://arXiv.org/abs/math.RA/0104158>, J. Lond. Math. Soc. 64, 13–28 (2001)
- [35] L. Siebenmann, *A total Whitehead torsion obstruction to fibering over the circle*, Comm. Math. Helv. 45, 1–48 (1970)
- [36] J. Stallings, *On fibering certain 3-manifolds*, Proc. 1961 Georgia Conf. on the Topology of 3-manifolds, Prentice-Hall, 95–100 (1962)
- [37] C.T.C. Wall, *Finiteness conditions for CW complexes II.*, Proc. Roy. Soc. London Ser. A 295, 129–139 (1965)