



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Association of inbreeding and regional equine leucocyte antigen homozygosity with the prevalence of insect bite hypersensitivity in Old Kladruber horse

Citation for published version:

Vostry, L, Vostra-Vydrova, H, Citek, J, Gorjanc, G & Curik, I 2021, 'Association of inbreeding and regional equine leucocyte antigen homozygosity with the prevalence of insect bite hypersensitivity in Old Kladruber horse', *Animal Genetics*. <https://doi.org/10.1111/age.13075>

Digital Object Identifier (DOI):

[10.1111/age.13075](https://doi.org/10.1111/age.13075)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Animal Genetics

Publisher Rights Statement:

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Association of inbreeding and regional equine leucocyte antigen homozygosity with the prevalence of insect bite hypersensitivity in Old Kladruber horse

L. Vostry* , H. Vostra-Vydrova*[†] , J. Citek^{‡,§}, G. Gorjanc^{||}  and I. Curik** 

*Czech University of Life Sciences, Kamycka 129, Prague 16500, Czech Republic. [†]Institute of Animal Science, Pratelstvi 815, Prague 10400, Czech Republic. [‡]South Bohemia University, Branisovska 31a, Ceske Budejovice 370 05, Czech Republic. [§]Veterinary Research Institute, Hudcova 296/70, Brno 621 00, Czech Republic. ^{||}The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Edinburgh, Midlothian EH259RG, UK. **Faculty of Agriculture, University of Zagreb, Svetošimunska cesta 25, Zagreb 10000, Croatia.

Summary

Inbreeding depression is the reduction of performance caused by mating of close relatives. In livestock populations, inbreeding depression has been traditionally estimated by regression of phenotypes on pedigree inbreeding coefficients. This estimation can be improved by utilising genomic inbreeding coefficients. Here we estimate inbreeding depression for insect bite hypersensitivity (IBH) prevalence, the most common allergic horse disease worldwide, in Old Kladruber horse. In a deep pedigree with 3214 horses (187 genotyped), we used a generalised linear mixed model with IBH phenotype from 558 horses examined between 1996 and 2009 (1368 records). In addition to the classical pedigree information, we used the single-step approach that enabled joint use of pedigree and genomic information to estimate inbreeding depression overall genome and equine leucocyte antigen (*ELA class II*) region. Significant inbreeding depression was observed in all models fitting overall inbreeding coefficients (odds ratio between 1.018 and 1.074, $P < 0.05$) with the exception of Kalinowski's new inbreeding ($P = 0.0516$). The increase of *ELA class II* inbreeding was significantly associated with increased prevalence of IBH (odds ratio 1.018; $P = 0.027$). However, when fitted jointly with the overall inbreeding coefficient, the effect of *ELA class II* inbreeding was not significant (odds ratio 1.016; $P = 0.062$). Overall, the higher *ELA class II* and/or overall inbreeding (pedigree or genomic) was associated with increased prevalence of IBH in Old Kladruber horses. The single-step approach provides an efficient use of all the available pedigree, genomic, and phenotype information for estimation of overall and regional inbreeding effects.

Keywords genomics, horse, inbreeding, inbreeding depression, insect bite hypersensitivity, pedigree

Mating of relatives is unavoidable in closed and finite populations and can lead to inbreeding depression – higher incidence of defects caused by recessive alleles and redistribution of genetic variance within and between populations. Inbreeding depression is a well-documented phenomenon,

Address for correspondence

L. Vostry, Czech University of Life Sciences, Kamycka 129, 16500 Prague, Czech Republic.

E-mail: vostry@af.czu.cz

and

I. Curik, Faculty of Agriculture, University of Zagreb, Svetošimunska cesta 25, 10000 Zagreb, Croatia.

E-mail: icurik@agr.hr

Accepted for publication 22 April 2021

known as the reduction of performance in quantitative traits under constant selection pressure such as fitness and health traits (Crnokrak & Roff 1999; Bittles & Black 2010; Leroy 2014).

In livestock populations, inbreeding depression has been traditionally estimated by regression of phenotypes on pedigree inbreeding coefficients (Baker *et al.* 1945; Leroy 2014). Alternatively, some other pedigree-derived coefficients, for example Kalinowski's new inbreeding coefficient, have been successfully used (Mc Parland *et al.* 2009; Doekes *et al.* 2019; Curik *et al.* 2020). Development of large scale high-throughput genotyping has enabled quantification of individual genomic inbreeding in human (McQuillan *et al.* 2008), wild (Santure *et al.* 2010), and livestock populations

(VanRaden 2008; Curik *et al.* 2014). The substitution of the pedigree with genomic inbreeding coefficient, improves the estimation of inbreeding depression from several perspectives (Kardos *et al.* 2016; Curik *et al.* 2017). In livestock populations, there are often many individuals in the pedigree that have phenotype records but are not genotyped. Legarra *et al.* (2020) showed how to estimate inbreeding coefficients by combining pedigree and genomic relationship matrices, which is commonly referred to as the single-step approach (Legarra *et al.* 2009). Recently, this approach was used in the estimation of inbreeding depression for semen traits in the Basco-Béarnaise dairy sheep breed (Antonios *et al.* 2021).

Classical horse breeds are often bred as closed populations and are known to reach high inbreeding often accompanied by inbreeding depression. Consequently, inbreeding depression has been observed for racing performance (Klemetsdal 1998; Todd *et al.* 2018), reproductive traits (Van Eldik *et al.* 2006; Müller-Unterberg *et al.* 2017), morphological traits (Gómez *et al.* 2009; Vostrý *et al.* 2011), and health traits (Sánchez-Guerrero *et al.* 2019).

The Old Kladruber horse, an old Czech breed established in 1579 that belongs to the Spanish horse group, has been maintained in a closed population for a long time with recent average inbreeding of 0.13 (Vostrá-Vydrová *et al.* 2016). In this population, inbreeding depression has been already observed for melanoma and vitiligo grading (Hofmanová *et al.* 2016) as well as for reproductive traits (Šichtař *et al.* 2017). The breed is affected also by insect bite hypersensitivity (IBH), the most common allergic skin disease caused by the bites of insects of the genus *Culicoides*. There is a common consensus that IBH is genetically controlled, and the starting impulse is a bite by midges of the *Culicoides* spp. and less frequently *Simulium* spp. The subsequent allergic reaction is a result of hypersensitivity to the allergens contained in their saliva (van der Rijt *et al.* 2008). The main symptoms are pruritus (itchiness), scaling, and hair loss. Clinical manifestation of IBH is due to an allergic reaction sharing some similarities with human atopic dermatitis. IBH is appearing in various horse breeds all over the world as a complex disease affected by environmental and genetic factors (Unkel *et al.* 1987; Eriksson *et al.* 2008; Schurink *et al.* 2011).

The impact of inbreeding on the IBH was analysed in Dutch Friesian horse and found non-significant (Schurink *et al.* 2011). In contrast, Andersson *et al.* (2012) have shown that increased homozygosity across the entire equine leucocyte antigen (*ELA*) class II region is associated with a higher risk of IBH. Unfortunately, Andersson *et al.* (2012) did not analyse if overall or regional inbreeding increases the risk of higher IBH.

The aim of this study was to estimate the effects of inbreeding on the prevalence of IBH in the Old Kladruber horse by logistic linear mixed regression from data with a heterogeneous combination of pedigree, genomic, and

phenotypic information. Inbreeding was based on the combined pedigree–genomic single-step approach of Legarra *et al.* (2009, 2020) and compared with the classic pedigree approach. We also estimated the effect of *ELA class II* regional inbreeding to test the results from Andersson *et al.* (2012). With respect to the prevalent heterogeneous combination of pedigree, genomic, and phenotypic information in many livestock populations, the application of single-step approach is empowering.

Materials and methods

Phenotype, pedigree, and genomic information


The analysis was based on 1368 IBH inspections from 558 horses. We assumed that all horses have experienced the same probability of mosquito bites as all phenotyped horses have been kept at the same stud and were exposed to the same environmental factors, including biting insects. The National Stud Kladrub is situated in a temperate climate zone, at an altitude of about 200 m above the sea level in the eastern part of Bohemia, Czech Republic. While conditions are favourable for horses, they are also favourable for biting insects such as *Culicoides*, for example *Culicoides nubeculosus* and *Culicoides obsoletus* (van der Rijt *et al.* 2008). The data on the Old Kladruber horses were provided by the stud book. The inspections were performed visually twice a year (May and October) between 1996 and 2009 by the same person on a binary scale with respect to the prevalence of the typical skin lesions (0 when clinical symptoms were absent and 1 when clinical symptoms were present). Similar IBH scoring system was also used by Grevenhof *et al.* (2007), Peeters *et al.* (2015) and Schurink *et al.* (2011). During the measurement period, 558 horses were inspected several times and 29% of horses (163) showed clinical symptoms of IBH. The pattern of the insect bite hypersensitivity prevalence with respect to the sex and horse age at the inspection is presented in Fig. 1.

This prevalence was much higher compared to 8.1% observed in Swedish-born Icelandic horses (Eriksson *et al.* 2008), 8.8% in Shetland ponies (Schurink *et al.* 2009), 18.2% in Dutch Friesian broodmares (Schurink *et al.* 2011), and 10% in Belgian Warmblood horses (Peeters *et al.* 2015).

The total pedigree was up to 52 generations deep and comprised of 3214 individuals, with mean complete generation equivalent (EQC, Maignel *et al.* 1996) equal to 15.58 for the inspected horses. Overall, 187 horses were genotyped with the Equine SNP70 BeadChip (Illumina), consisting of 65 102 evenly distributed SNPs over the equine genome. This genotyping has been performed as a part of the routine diversity management, recently introduced into the breeding programme.

SNP data were edited and analysed using `PLINK` 1.9 (Chang *et al.* 2015). Only autosomal SNPs with known chromosomal position were analysed with individual call

Figure 1 The number of animals with insect bite hypersensitivity by sex and age at the inspection.



rate >0.9 and SNP call rate >0.9 . The minor allele frequency (MAF) was filtered according to two different approaches. Filtering for $MAF >0.01$ was performed in the calculation of the \mathbf{G} matrix and FG, while filtering for MAF was not performed in the calculation of runs of homozygosity (ROH-based genomic inbreeding coefficients as recommended by Meyerjans *et al.* (2020). To define the parameters of an ROH, we set the minimum density to 0.05 Mb/SNP and the largest gap to 1 Mb. The minimal number of SNPs in each ROH was set to 20. Similar parameters were used by Todd *et al.* (2018). The maximum number of heterozygous SNPs allowed within ROH >5 Mb was 0 and maximum number of missing SNPs allowed within ROH >5 Mb was 1. Unfortunately, only 47 inspected horses were genotyped (seven were affected and 40 were unaffected by IBH) while the remainder of 137 horses were not phenotyped (inspected for IBH). However, all included horses were highly related and were kept in the same environment, on the National Stud in Kladruby nad Labem.

Estimation of pedigree and genomic inbreeding coefficients

We estimated three different pedigree inbreeding coefficients: classical (F_X); classical restricted to five generations (F_{X5}); and Kalinowski's new (F_{NEW}) inbreeding coefficient. When pedigree depth varies considerably, estimation of inbreeding depression can be biased as individual inbreeding coefficient estimates refer to different base populations. To avoid this potential bias, we used the F_{X5} as all phenotyped horses had complete pedigrees for at least five generations. Note that this inbreeding coefficient refers to the recent inbreeding. F_{NEW} is defined as the probability that any allele in an individual is autozygous for the first time and is assumed to better capture inbreeding depression

caused by alleles with large detrimental effects or detrimental alleles that have not been purged (Doekes *et al.* 2019; Curik *et al.* 2020). F_X and F_{X5} were calculated by PEDIG software (Boichard 2002), while F_{NEW} (Kalinowski *et al.* 2000) was calculated by gene dropping with 1 000 000 repetitions as suggested by (Baumung *et al.* 2015) and with corrections implemented in the GRAIN 2.2 software (Doekes *et al.* 2020).

Following the idea presented in Legarra *et al.* (2020) we estimated genomic inbreeding coefficients for all horses, by combining pedigree and genomic information with the single-step approach. This approach projects genomic relationships (\mathbf{G} matrix; VanRaden 2008) to all ungenotyped individuals in a pedigree (\mathbf{A} matrix; Legarra *et al.* 2009; Christensen & Lund 2010). The resulting projected matrix is often denoted as the \mathbf{H} matrix. The projection requires careful consideration to account for different base populations of pedigree and genomic information. To address this, we used one 'meta-founder' for pedigree and expected allele frequency equal to 0.5 (Legarra *et al.* 2015). With this approach, we estimated three genomic inbreeding coefficients as the diagonal of \mathbf{H} matrix minus one: (a) F_H representing whole genome; (b) F_{ELA-H} representing the *ELA class II* region with 87 SNPs between 32 and 34 Mb on *ECA20* (Brinkmeyer-Langford *et al.* 2013); and (c) $F_{noELA-H}$ representing whole genome without the *ELA class II* region. $F_{noELA-H}$ (whole genome SNPs without 87 SNPs covering the *ELA class II* region) was calculated to reduce confounding between overall (F_H) and regional (F_{ELA-H}) inbreeding.

In addition, for the genotyped animals we also calculated overall and regional ROH inbreeding coefficient (McQuillan *et al.* 2008; Curik *et al.* 2014), to enable comparison of the inbreeding level in other horse populations. Calculated ROH inbreeding coefficient ($F_{ROH>5Mb}$) was defined as the proportion of the genome in ROH of a certain minimal length

relative to the whole autosomal genome ($F_{\text{ROH}} = L_{\text{ROH}}/L_{\text{AUTOSOME}}$). *ELA class II* ROH inbreeding coefficient ($F_{\text{ELA-ROH}>5\text{Mb}}$) was defined in the same way but only for the *ELA class II* region. We estimated all genomic inbreeding coefficients with the `R` software, except for ROH inbreeding coefficients that were estimated with `PLINK 1.9` (Chang *et al.* 2015) following settings of Todd *et al.* (2018).

Note that genomic inbreeding coefficients derived from **H** are defined relative to a population with allele frequency of 0.5 for all loci, due to the meta-founder single-step approach (Legarra *et al.* 2015). At this allele frequency, we expect 50% heterozygous loci. Therefore, these inbreeding coefficients express deviations from this expectation and an individual with less (more) than 50% heterozygous loci will have positive (negative) inbreeding coefficient. In contrast, pedigree and ROH based inbreeding coefficients are defined relative to a past base population (founders in the pedigree or conceptual base population defined by the ROH length). Therefore, these inbreeding coefficients are expressed as probabilities and strictly between 0 and 1.

Modelling of inbreeding depression

Depending on the information source used in the estimation of inbreeding coefficients, three different modelling approaches were used. Models from the first approach were based on the pedigree inbreeding coefficients (F_X , F_{X5} and F_{NEW}). In the second approach, three models were based on inbreeding coefficients derived from the single-step (F_H , $F_{\text{ELA-H}}$, and $F_{\text{ELA-H}}$ with $F_{\text{noELA-H}}$).

Calculations were performed with the GLIMMIX procedure, which fitted logistic linear mixed model for repeated measurements (SAS Inst. Inc.). Here, the dependent variable (y_i) has value of 1 (occurrence of IBH clinical symptoms) with probability π_i or 0 (absence of IBH clinical symptoms) with probability $1 - \pi_i$ for observation i . We modelled this probability as described in Citek *et al.* (2017) with the addition of inbreeding coefficients as covariates:

$$\log\left(\frac{\pi_{ijkl}}{1 - \pi_{ijkl}}\right) = \text{Year}_i + \text{Age}_j + \text{Sex}_k + bF + p_l + a_l,$$

where Year_i was a fixed effect of the i th year of inspection ($i = 1996, \dots, 2009$), Age_j is a fixed effect of the j th year of age of the horse ($j = 1, \dots, 16$), Sex_k is a fixed effect of the k th sex ($k = \text{stallion or mare}$; geldings were not included), p_{li} is a random permanent environment of the l th animal with $\mathbf{p} \sim N(0, I_p \sigma_p^2)$, I_p is the identity matrix of the appropriate dimensions and σ_p^2 is the permanent environmental variance, a_l is a random additive genetic effect of the l th animal with $a \sim N(0, Q \sigma_a^2)$, **Q** is one of the following: pedigree relationship matrix **A** (Henderson 1976) when pedigree inbreeding coefficients were fitted; or single-step relationship **H** matrix (Legarra *et al.* 2009, 2015) when single-step inbreeding coefficients were fitted, while b

presents regression coefficient for the various inbreeding coefficients. Separate models were fitted for each inbreeding coefficient (F_X , F_{X5} , F_{NEW} , F_H , and $F_{\text{ELA-H}}$). We also modelled joint effects of the *ELA class II* regional inbreeding and overall inbreeding (F_H with $F_{\text{noELA-H}}$). The suitability of fixed and random effects included in the model was discussed in Citek *et al.* (2017).

The goodness of fit of each of the analysed models was tested by a second-order bias correction of Akaike information criterion (AICc), which is corrected based on samples size (Burnham & Anderson 2002):

$$\text{AICc} = -2 \log \text{likelihood} + 2k(k+1)/(n-k-1),$$

where k is the number of estimated parameters and n is the number of observations.

Results and discussion

Inbreeding level

Descriptive statistics of various inbreeding coefficients, equivalent complete generations, and inspected horses for IBH are presented in Table 1. The observed pedigree inbreeding level in the Old Kladruber horse (0.139), with increase of inbreeding equal to 0.009 per EQC, was higher than inbreeding level, ranging from 0.103 to 0.125, estimated in other horse breeds with very deep and complete pedigrees such as Thoroughbred (Mahon & Cunningham 1982), Lipizzan (Curik *et al.* 2003), Andalusian (Valera *et al.* 2005), and Lusitano (Vicente *et al.* 2012). Somewhat lower inbreeding, but with higher increase in inbreeding level per EQC (0.0134), was observed for the recent (F_{X5}) as well as for the new inbreeding coefficient (F_{NEW} ; Table 1. Results reveal that the genetic diversity was affected by bottleneck that occurred in the first half of the 20th Century. At that time, there was a substantial reduction in the size of Old Kladruber population. After this period, from 1973, the regeneration process of the Old Kladruber breed began (Vostrá-Vydrová *et al.* 2016).

Genomic ROH inbreeding level ($F_{\text{ROH}>5\text{Mb}}$) was lower than the estimated pedigree inbreeding level (Table 1) but with comparable increase of inbreeding per generations (0.010) assuming that 5 Mb long ROH estimates are expected to descend from ancestors that are 10 generations back in the pedigree. The genomic ROH inbreeding was higher than observed by in Grilz-Seger *et al.* (2019) in Lipizzan (0.035), Austrian Haflinger (0.021), Italian Haflinger (0.051), and Noriker (0.029) horse breeds but similar to the Bosnian Mountain horse (0.101) observed by Druml *et al.* (2018).

A much higher inbreeding estimate, 0.399 (F_H), was obtained from the relationship matrix **H** due to a distant (ancient) base population. There was no notable difference between overall estimates with or without 87 SNPs from the

Table 1 Descriptive statistics for pedigree (F_X , F_{X5} , F_{NEW}), genomic (F_G , $F_{ROH>5Mb}$, F_H) and ELA class II regional (F_{ELA-G} , $F_{ELA-ROH>5Mb}$, F_{ELA-H}) inbreeding coefficients, equivalent complete generations (EQC), and horse age at inspection in Old Kladruber Horse.

Variable	Horses	Mean	Standard deviation	Minimum	Maximum
F_X	558	0.139	0.040	0.017	0.252
F_{X5}	558	0.067	0.030	0.000	0.150
F_{NEW}	558	0.029	0.011	0.003	0.080
F_H	558	0.399	0.032	0.287	0.485
F_{ELA-H}	558	0.343	0.104	0.060	1.087
$F_{ROH>5Mb}$	187	0.099	0.037	0.010	0.205
$F_{ELA-ROH>5Mb}$	187	0.051	0.182	0.000	1.000
EQC	558	15.580	0.761	13.010	17.070
Age (years)	558	5.307	4.110	1.000	16.000

F_X and F_{X5} - pedigree inbreeding coefficients based on all and five-generations; F_{NEW} - Kalinowski's new inbreeding coefficient; F_H and F_{ELA-H} genomic inbreeding coefficients representing whole genome and the ELA class II region; $F_{ROH>5Mb}$ and $F_{ELA-ROH>5Mb}$ - runs of homozygosity (ROH) inbreeding coefficients calculated from the sum of all ROH segments greater than 5 Mb representing whole genome and the ELA class II region; EQC - equivalent complete generations; Age - year of age of the horse.

ELA class II region as the region is only a small proportion of the genome (results not shown). All estimates of the ELA class II regional inbreeding were lower than overall inbreeding estimates ($F_{ELA-H} = 0.343$ vs. $F_H = 0.399$, and $F_{ELA-ROH>5Mb} = 0.051$ vs. $F_{ROH>5Mb} = 0.099$), which might be a consequence of the heterozygote advantage (overdominance) in the ELA class II region.

Relationship among inbreeding coefficients

Correlations between the various inbreeding coefficients are presented in Fig. 2. High correlations, ranging from 0.505 to 0.955, were observed among all overall inbreeding coefficients (F_X , F_{X5} , F_{NEW} , F_H , and $F_{ROH>5Mb}$). Low correlations were observed between all overall and regional ELA class II inbreeding coefficients, while relatively high correlation was observed between two regional estimates F_{H-ELA} and $F_{ELA-ROH>5Mb}$ (0.651).

Inbreeding depression

The results of the inbreeding depression analyses for the prevalence of IBH in Old Kladruber Horse are presented in Table 2. Significant overall inbreeding depression, that is the increase of overall inbreeding increased the prevalence of IBH, was estimated in models with F_X , F_{X5} , and F_H , while the estimate with F_{NEW} was in the same direction and close to significant ($P = 0.0516$).

When fitted alone in the model, ELA class II inbreeding was significantly associated with increasing the prevalence

of IBH, corroborating the negative impact of inbreeding on IBH. Interestingly, according to AICc values, the model with F_{ELA-H} as a single covariate was the best model in this study. All models based on the single-step inbreeding fitted the data better than models based on pedigree inbreeding (Table 2) pointing to the advantage of utilising genomic information in addition to pedigree information.

The presence of a polygenic effect on IBH has been observed in several studies – narrow sense heritability (h^2) on an IBH status (four categories) was estimated at 0.33 in Swedish-born Icelandic horses (Eriksson *et al.* 2008) and at 0.16 on a liability scale in Dutch Friesian Broodmares (Schurink *et al.* 2011), between 0.08 and 0.24 in the Dutch Shetland breeding mares (Schurink *et al.* 2009) and between 0.36 and 0.63 in the Old Kladruber horse (Citek *et al.* 2017). The wide range in the last two studies was due to different model definitions. The polygenic nature of IBH was also manifested in several GWAS studies, while several candidate genes located on *ECA7*, *9*, *11* and *20* were also reported (Schurink *et al.* 2012, 2013; Shrestha *et al.* 2015; Velie *et al.* 2016). The most consistent association with IBH across various breeds was observed for the major histocompatibility complex located on *ECA20*, or more precisely for the ELA class I and II regions (Marti *et al.* 1992; Andersson *et al.* 2012; Schurink *et al.* 2012).

In contrast to other genetic analyses of IBH, this is the first study where significant overall inbreeding depression is reported for IBH. Up to now, the effect of inbreeding on IBH has been estimated only in Dutch Friesian horse (Schurink *et al.* 2011), but no significant estimates were observed. Our results support the positive association of the ELA class II region homozygosity with IBH previously reported by Andersson *et al.* (2012). In the models where the overall and regional ELA class II inbreeding were analysed simultaneously, regional inbreeding was not significant while overall inbreeding was significant. Still, we think that in the Old Kladruber horse, both overall and ELA class II inbreeding are associated with increased prevalence of IBH. We have five arguments this. First, in almost all models overall inbreeding depression was significant, while regional inbreeding depression was significant when fitted independently and gave the best model fit according to AICc. Second, estimates obtained with the single inbreeding covariate models were similar to estimates obtained with two inbreeding covariate models. Third, correlations between overall and regional ELA class II inbreeding coefficients were very small, which should exclude potential confounding. Fourth, the regional inbreeding depression was close to significant ($P = 0.062$) in the two inbreeding covariate model. Fifth, significant regional inbreeding depression was observed in other studies (Andersson *et al.* 2012).

Overall, models based on the single-step relationship matrix combining genomic and pedigree information outperformed models based only on pedigree information




Figure 2 Pearson correlations among pedigree (F_X , F_{X5} , and F_{NEW}), genomic ($F_{ROH>5Mb}$ and F_H) and ELA class II regional ($F_{ELA-ROH>5Mb}$ and F_{ELA-H}) inbreeding coefficients in Old Kladruber Horse. Estimates obtained on 187 (558) genotyped horses are presented in the upper (lower) diagonal. NA marks values that could not be calculated.

Table 2 Inbreeding depression for the insect bite hypersensitivity prevalence in Old Kladruber Horse estimated with the pedigree and single-step models fitted to the 558 phenotyped horses.

Inbreeding coefficient	Estimate (SE)	Odds ratio	P-value	AICc (Δ AICc)	EEM ¹	Matrix ²
F_{X5}	0.071 (0.025)	1.074	0.0049	5541 (222)	Table S1	A
F_X	0.042 (0.019)	1.043	0.0277	5546 (227)	Table S2	A
F_{NEW}	0.138 (0.071)	1.148	0.0516	5551 (232)	Table S3	A
F_{ELA-H}	0.018 (0.008)	1.018	0.0270	5319 (0)	Table S4	H
F_{ELA-H}	0.016 (0.008)	1.016	0.0620	5361 (42)	Table S5	H
$F_{noELA-H}$	0.061 (0.029)	1.063	0.0365			
F_H^3	0.068 (0.029)	1.071	0.0190	5368 (49)	Table S6	H

AICc, corrected Akaike information criterion.

¹EEM gives a detailed presentation of models with all estimated environmental effects in the Supported Information (from Tables S1–S6).

²Relationship matrix used; **A** is a pedigree relationship matrix (Henderson 1976) and **H** is a single-step relationship matrix that combines all the available genomic and pedigree information (Legarra *et al.* 2009, 2015).

³ F_H is inbreeding coefficient for the whole genome.

F_X and F_{X5} - pedigree inbreeding coefficients based on all and five-generations; F_{NEW} - Kalinowski's new inbreeding coefficient; F_H , F_{ELA-H} and $F_{noELA-H}$ genomic inbreeding coefficients representing whole genome, the ELA class II and whole genome without the ELA class II region.

(Table 2). The utilisation of all available information in the estimation of inbreeding depression seems intuitively logical, although computer simulations are required to better explore all properties and assumptions of the applied models. With a recent estimate of inbreeding depression for semen traits in the Basco-Béarnaise dairy sheep breed (Antonios *et al.* 2021), this is the first study to use the single-step approach to estimate total and regional inbreeding depression. This option might be important for analysing traits where genes with large detrimental impact are participating in the inbreeding depression, particularly

as the regional inbreeding might substantially differ from the expected inbreeding estimated by the pedigree inbreeding coefficients (Kardos *et al.* 2016; Curik *et al.* 2017).

Conclusions

The overall pedigree or genomic and regional *ELA class II* inbreeding were positively associated with the prevalence of insect bite hypersensitivity in Old Kladruber horses. The utilisation of single-step inbreeding coefficients that combines all the available genomic and pedigree information is a

useful approach in the estimation of detrimental inbreeding effects.

Acknowledgements

L.V. and H.V.V. acknowledge support from the Ministry of Agriculture of the Czech Republic (Project Nos. QK1910156 and MZE-RO0719). J.C. acknowledges support from the Ministry of Agriculture of the Czech Republic (Project No. QK1910320). I.C. acknowledges support from the Croatian Science Foundation (Project ANAGRAMS-IP-2018-01-8708) while G.G. acknowledges support from the BBSRC to The Roslin Institute (BBS/E/D/30002275) and The University Of Edinburgh's Data-Driven Innovation Chancellor's Fellowship.

Data availability statement

Phenotypic and genotypic data were obtained from National Stud in Kladruby nad Labem. These datasets were used under a material transfer agreement, and hence, are not publicly available. However, data are available upon request to LV and with the permission of National Stud in Kladruby nad Labem.

References

- Andersson L.S., Swinburne J.E., Meadows JRS *et al.* (2012) The same ELA class II risk factors confer equine insect bite hypersensitivity in two distinct populations. *Immunogenetics* **64**, 201–8.
- Antonios S., Rodríguez-Ramilo S.T., Aquilar I., Astruc J.M., Legarra A. & Vitezica Z.G. (2021) Genomic and pedigree estimation of inbreeding depression for semen traits in the Basco-Béarnaise dairy sheep breed. *Journal of Dairy Science* **104**, 3221–30.
- Baker G.A., Mead S.W. & Regan W.M. (1945) Effect of inbreeding on the growth curves of height at withers, weight, and heart girth of Holstein females. *Journal of Dairy Science* **28**, 607–10.
- Baumung R., Farkas J., Boichard D., Mészáros G., Sölkner J. & Curik I. (2015) GRAIN: a computer program to calculate ancestral and partial inbreeding coefficients using a gene dropping approach. *Journal of Animal Breeding and Genetics* **132**, 100–8.
- Bittles A.H. & Black M.L. (2010) Consanguinity, human evolution, and complex diseases. *Proceedings of the National Academy of Sciences* **107**, 1779–86.
- Boichard D. (2002) PEDIG: a Fortran package for pedigree analysis suited for large populations. 7th World Congress on Genetics Applied to Livestock Production. 28.
- Brinkmeyer-Langford C.L., Cai J.J., Gill C.A. & Skow L.C. (2013) Microsatellite variation in the equine MHC. *Animal Genetics* **44**, 267–75.
- Burnham K.P. & Anderson D.R. (2002) *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*, 2nd edn. New York: Springer.
- Chang C.C., Chow C.C., Tellier L.C., Vattikuti S., Purcell S.M. & Lee J.J. (2015) Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience* **4**, 7.
- Christensen O.F. & Lund M.S. (2010) Genomic prediction when some animals are not genotyped. *Genetics Selection Evolution* **42**, 2.
- Citek J., Vostry L., Vostra-Vydrova H., Brzakova M. & Prantlova V. (2017) Genetic parameters of insect bite hypersensitivity in the Old Grey Kladruber horse. *Journal of Animal Science* **95**, 53–8.
- Crnokrak P. & Roff D.A. (1999) Inbreeding depression in the wild. *Heredity* **83**, 260–70.
- Curik I., Ferenčaković M. & Sölkner J. (2014) Inbreeding and runs of homozygosity: a possible solution to an old problem. *Livestock Science* **166**, 26–34.
- Curik I., Ferenčaković M. & Sölkner J. (2017) Genomic dissection of inbreeding depression: a gate to new opportunities. *Revista Brasileira de Zootecnia* **46**, 773–82.
- Curik I., Kövér G., Farkas J., Szendrő Z., Romvári R., Sölkner J. & Nagy I. (2020) Inbreeding depression for kit survival at birth in a rabbit population under long-term selection. *Genetics Selection Evolution* **52**, 39.
- Curik I., Zechner P., Sölkner J., Achmann R., Bodo I., Dovc P., Kavár T., Marti E. & Brem G. (2003) Inbreeding, microsatellite heterozygosity, and morphological traits in lipizzan horses. *Journal of Heredity* **94**, 125–32.
- Doekes H.P., Curik I., Nagy I., Farkas J., Kövér G.Y. & Windig J.J. (2020) Revised calculation of Kalinowski's ancestral and new inbreeding coefficients. *Diversity* **12**, 155.
- Doekes H.P., Veerkamp R.F., Bijma P., de Jong G., Hiemstra S.J. & Windig J.J. (2019) Inbreeding depression due to recent and ancient inbreeding in Dutch Holstein-Friesian dairy cattle. *Genetics Selection Evolution* **51**, 54.
- Druml T., Neuditschko M., Grilz-Seger G., Horna M., Ricard A., Mesarič M., Cotman M., Pausch H. & Brem G. (2018) Population networks associated with runs of homozygosity reveal new insights into the breeding history of the haflinger horse. *Journal of Heredity* **109**, 384–92.
- Eriksson S., Grandinson K., Fikse W.F., Lindberg L., Mikko S., Broström H., Frey R., Sundquist M. & Lindgren G. (2008) Genetic analysis of insect bite hypersensitivity (summer eczema) in icelandic horses. *Animal* **2**, 360–5.
- Gómez M.D., Valera M., Molina A., Gutiérrez J.P. & Goyache F. (2009) Assessment of inbreeding depression for body measurements in Spanish Purebred (Andalusian) horses. *Livestock Science* **122**, 149–55.
- Grevenhof E.M., Ducro B., Heuven H.C.M. & Bijma P. (2007) Identification of environmental factors affecting the prevalence of insect bite hypersensitivity in Shetland ponies and Friesian horses in the Netherlands. *Equine Veterinary Journal* **39**, 69–73.
- Grilz-Seger G., Druml T., Neuditschko M., Dobretsberger M., Horna M. & Brem G. (2019) High-resolution population structure and runs of homozygosity reveal the genetic architecture of complex traits in the Lipizzan horse. *BMC Genomics* **20**, 174.
- Henderson C.R. (1976) A simple method for computing the inverse of a numerator relationship matrix used in prediction of breeding values. *Biometrics* **32**, 69–83.
- Hofmanová B., Vostrá-Vydrová H., Majzlík I. & Vostrý L. (2016) The effect of inbreeding on melanoma and vitiligo occurrence in Old Kladruber grey horses. *Acta Agriculturae Slovenica* **5**, 179–82.
- Kalinowski S.T., Hedrick P.W. & Miller P.S. (2000) Inbreeding depression in the Speke's gazelle captive breeding program. *Conservation Biology* **14**, 1375–84.

- Kardos M., Taylor H.R., Ellegren H., Luikart G. & Allendorf F.W. (2016) Genomics advances the study of inbreeding depression in the wild. *Evolutionary Applications* **9**, 1205–18.
- Klemetsdal G. (1998) The effect of inbreeding on racing performance in Norwegian cold-blooded trotters. *Genetics Selection Evolution* **30**, 351.
- Legarra A., Aguilar I. & Colleau J.J. (2020) Short communication: Methods to compute genomic inbreeding for ungenotyped individuals. *Journal of Dairy Science* **103**, 3363–7.
- Legarra A., Aguilar I. & Misztal I. (2009) A relationship matrix including full pedigree and genomic information. *Journal of Dairy Science* **92**, 4656–63.
- Legarra A., Christensen O.F., Vitezica Z.G., Aguilar I. & Misztal I. (2015) Ancestral relationships using metafounders: finite ancestral populations and across population relationships. *Genetics* **200**, 455–68.
- Leroy G. (2014) Inbreeding depression in livestock species: review and meta-analysis. *Animal Genetics* **45**, 618–28.
- Mahon GAT & Cunningham E.P. (1982) Inbreeding and the inheritance of fertility in the thoroughbred mare. *Livestock Production Science* **9**, 743–54.
- Maignel L., Boichard D. & Verrier E. (1996) Genetic variability of French dairy breeds estimated from pedigree information. *Interbull Bulletin* **14**, 49–54.
- Marti E., Gerber H. & Lazary S. (1992) On the genetic basis of equine allergic diseases: II. Insect bite dermal hypersensitivity. *Equine Veterinary Journal* **24**, 113–7.
- Mc Parland S., Kearney F. & Berry D.P. (2009) Purging of inbreeding depression within the Irish Holstein-Friesian population. *Genetics Selection Evolution* **41**, 16.
- McQuillan R., Leutenegger A.L., Abdel-Rahman R. *et al.* (2008) Runs of homozygosity in European populations. *The American Journal of Human Genetics* **83**, 359–72.
- Meyermans R., Grossen W., Buys N. & Janssens S. (2020) How to study runs of homozygosity using PLINK? A guide for analysing medium density SNP data in livestock and pet species. *BMC Genomics* **21**, 94.
- Müller-Unterberg M., Wallmann S. & Distl O. (2017) Effects of inbreeding and other systematic effects on fertility of Black Forest Draught horses in Germany. *Acta Veterinaria Scandinavica* **59**, 70.
- Peeters L.M., Janssens S., Brebels M. & Buys N. (2015) Genetic parameters and estimated breeding values of insect bite hypersensitivity in Belgian Warmblood horses. *The Veterinary Journal* **206**, 420–2.
- Sánchez-Guerrero M.J., Solé M., Azor P.J., Sölkner J. & Valera M. (2019) Genetic and environmental risk factors for vitiligo and melanoma in Pura Raza Español horses. *Equine Veterinary Journal* **51**, 606–11.
- Santure A.W., Stapley J., Ball A.D., Birkhead T.R., Burke T. & Slate J. (2010) On the use of large marker panels to estimate inbreeding and relatedness: empirical and simulation studies of a pedigreed zebra finch population typed at 771 SNPs. *Molecular Ecology* **19**, 1439–51.
- Schurink A., Ducro B.J., Bastiaansen J.W.M., Frankena K. & van Arendonk J.A.M. (2013) Genome-wide association study of insect bite hypersensitivity in Dutch Shetland pony mares. *Animal Genetics* **44**, 44–52.
- Schurink A., Ducro B.J., Heuven H.C.M. & van Arendonk J.A.M. (2011) Genetic parameters of insect bite hypersensitivity in Dutch Friesian broodmares. *Journal of Animal Science* **89**, 1286–93.
- Schurink A., van Grevenhof E.M., Ducro B.J. & van Arendonk J.A.M. (2009) Heritability and repeatability of insect bite hypersensitivity in Dutch Shetland breeding mares. *Journal of Animal Science* **87**, 484–90.
- Schurink A., Wolc A., Ducro B.J., Frankena K., Garrick D.J., Dekkers J.C. & van Arendonk J.A. (2012) Genome-wide association study of insect bite hypersensitivity in two horse populations in the Netherlands. *Genetics Selection Evolution* **44**, 31.
- Shrestha M., Eriksson S., Schurink A., Andersson L.S., Sundquist M., Frey R., Broström H., Bergström T., Ducro B. & Lindgren G. (2015) Genome-wide association study of insect bite hypersensitivity in Swedish-born Icelandic horses. *Journal of Heredity* **106**, 366–74.
- Šichtař J., Nehasilová A., Šimoník O. & Bubeníčková F. (2017) Effect of two freezing extenders on characteristic of fresh and frozen-thawed semen in endangered Old Kladruber stallions – a pilot study. *Czech Journal of Animal Science* **62**, 227–33.
- Todd E.T., Ho SYW, Thomson P.C., Ang R.A., Velie B.D. & Hamilton N.A. (2018) Founder-specific inbreeding depression affects racing performance in thoroughbred horses. *Scientific Reports* **8**, 6167.
- Unkel M., Simon D., Mayer M. & Sommer H. (1987) Zur genetischen Fundierung des Sommerekzems beim Islandpferd. *Journal of Animal Breeding and Genetics* **104**, 217–30.
- Valera M., Molina A., Gutiérrez J.P., Gómez J. & Goyache F. (2005) Pedigree analysis in the Andalusian horse: population structure, genetic variability and influence of the Carthusian strain. *Livestock Production Science* **95**, 57–66.
- van der Rijt R., Van der Boom R., De Jongema Y. & Sloet van Oldruitenborgh-Oosterbaan M.M. (2008) Culicoides species attracted to horses with and without insect hypersensitivity. *The Veterinary Journal* **178**, 91–7.
- van Eldik P., van der Waaij E.H., Ducro B., Kooper A.W., Stout T.A.E. & Colenbrander B. (2006) Possible negative effects of inbreeding on semen quality in Shetland pony stallions. *Theriogenology* **65**, 1159–70.
- VanRaden P. (2008) Efficient methods to compute genomic predictions. *Journal of Dairy Science* **91**, 4414–23.
- Velie B.D., Shrestha M., Francois L. *et al.* (2016) Using an inbred horse breed in a high density genome-wide scan for genetic risk factors of insect bite hypersensitivity (IBH). *PLoS One* **11**, e0152966.
- Vicente A.A., Carolino N. & Gama L.T. (2012) Genetic diversity in the Lusitano horse breed assessed by pedigree analysis. *Livestock Science* **148**, 16–25.
- Vostrá-Vydrová H., Vostrý L., Hofmanová B., Krupa E. & Zavadilová L. (2016) Pedigree analysis of the endangered Old Kladruber horse population. *Livestock Science* **185**, 17–23.
- Vostrý L., Čapková Z., Přibyl J. & Mach K. (2011) Analysis of czech cold-blooded horses: genetic parameters, breeding value and the influence of inbreeding depression on linear description of conformation and type characters. *Czech Journal of Animal Science* **56**, 217–30.

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Table S1. Estimated environmental and inbreeding effects with significant impact on insect bite hypersensitivity prevalence in Old Kladruber Horse: model is based on pedigree relationship matrix and five generations inbreeding coefficient (F_{X5}); model AICc value was equal to 5541.

Table S2. Estimated environmental and inbreeding effects with significant impact on insect bite hypersensitivity prevalence in Old Kladruber Horse: model is based on pedigree relationship matrix and classical inbreeding coefficient (F_X); model AICc value was equal to 5546.

Table S3. Estimated environmental and inbreeding effects with significant impact on insect bite hypersensitivity prevalence in Old Kladruber Horse: model is based on pedigree relationship matrix and Kalinowski's new inbreeding coefficient (F_{NEW}); model AICc value was equal to 5551.

Table S4. Estimated environmental and inbreeding effects with significant impact on insect bite hypersensitivity

prevalence in Old Kladruber Horse: model is based on single-step relationship matrix and inbreeding coefficient representing the *ELA class II* region (F_{ELA-H}); model AICc value was equal to 5319.

Table S5. Estimated environmental and inbreeding effects with significant impact on insect bite hypersensitivity prevalence in Old Kladruber Horse: model is based on single-step relationship matrix and inbreeding coefficients representing the *ELA class II* region ($F_{noELA-H}$) and the whole genome without the *ELA class II* region ($F_{noELA-H}$); model AICc value was equal to 5361.

Table S6. Estimated environmental and inbreeding effects with significant impact on insect bite hypersensitivity prevalence in Old Kladruber Horse: model is based on single-step relationship matrix and inbreeding coefficient representing the whole genome (F_{ELA-H}); model AICc value was equal to 5368.