



UNIVERSITI PUTRA MALAYSIA

**OPTIMISATION OF LIGNIN PEROXIDASE PRODUCTION FROM
PYCNOPORUS SP.**

ZURAIDAH BINTI ZANIRUN

FBSB 2008 16



OPTIMISATION OF LIGNIN PEROXIDASE PRODUCTION FROM
PYCNOPORUS SP.

By

ZURAIDAH BINTI ZANIRUN

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of Requirement for the Degree of Master of Science

SEPTEMBER 2008



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**OPTIMISATION OF LIGNIN PEROXIDASE PRODUCTION FROM
*PYCNOPORUS SP.***

By

ZURAIDAH BT ZANIRUN

SEPTEMBER 2008

Chairman: Suraini Abd. Aziz, PHD

Faculty: Biotechnology and Biomolecular Sciences

Ligninase or lignin peroxidase is gaining importance for their biotechnology application due to its integral role in the biodegradation of lignin, lignin related aromatic compounds and the potential use in industrial processes such as biopulping, biobleaching and bioremediation. Lignin peroxidase has been extensively studied and has been reported produced by white rot fungus. In this study, a preliminary study was done to screen for the highest lignin peroxidase producer from five locally isolated white rot fungus using agitated and non-agitated culture condition. The highest lignin peroxidase producer, identified as *Pycnoporus sp.* was selected for the optimisation study. Factorial design approach was chosen to determine the optimum conditions which significantly influence the production of lignin peroxidase by *Pycnoporus sp.* Optimum condition for the highest lignin peroxidase activity of 51.1 U/L was successfully achieved at 24 mM of



nitrogen concentration, agitation speed at 110 rpm, pH 3.5, inoculum concentration of 6×10^6 spores/ml and addition of 1 mM inducer (veratryl alcohol) in carbon limited media.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGOPTIMUMAN PENGHASILAN LIGNIN PEROKSIDASE DARIPADA
*PYCNOPORUS SP.***

Oleh

ZURAIDAH BT ZANIRUN

SEPTEMBER 2008

Chairman: Suraini Abd. Aziz, PHD

Faculty: Bioteknologi dan Sains Biomolekul

Ligninase atau lignin peroksidase semakin mendapat perhatian di dalam aplikasi bioteknologi di mana ia memainkan peranan penting di dalam bio-penguraian lignin dan juga sebatian aromatik berlignin serta berpotensi di dalam proses-proses industri seperti bio-pulpa, bio-pemutihan dan bioremediasi . Lignin peroksidase, telah dikaji dengan meluas dan dilaporkan dihasilkan oleh kulat pereput putih. Dalam kajian ini, penyaringan awal telah dilakukan untuk memilih kulat pereput putih tempatan yang menghasilkan lignin peroksidase tertinggi melalui kaedah fermentasi kelalang goncang dan statik. Kulat yang menghasilkan aktiviti lignin peroksidase tertinggi dikenalpasti sebagai *Pycnoporus* sp. telah dipilih untuk kajian pengoptimuman. Pendekatan 'Factorial Design' dipilih untuk menentukan keadaan optimum yang mempengaruhi penghasilan lignin peroksidase oleh kulat *Pycnoporus* sp. Keadaan optimum yang menghasilkan aktiviti lignin peroksidase tertinggi 51.1 U/L diperolehi pada kepekatan nitrogen sebanyak 24 mM,



kelajuan goncangan pada 110 psm, pH 3.5, kepekatan inokulum pada 6×10^6 spora/ml dan penambahan penggalak iaitu alkohol veratril di dalam media yang mengandungi sumber karbon yang terhad.

ACKNOWLEDGEMENTS

Bismillahirrahmanirrahim

First of all syukur Alhamdulillah to the Almighty ALLAH for giving me the strength to endure all challenges and completing this study. I also would like to take this opportunity to give a special thanks to my supervisor Assoc. Prof. Dr Suraini Abdul Aziz, my co-supervisor Assoc. Prof. Dr Foo Hooi Ling and Prof. Dr Mohd Ali Hassan for their advice, patience, concern and invaluable guidance throughout this project.

I would like to thanks all my labmates especially Farah, Fadly, Sauvaphap, Lay Sin and Christine whose help, motivate and enlightened me during the difficult moment.

Sincere thanks to my parents and parents-in-law for their comprehensive support and understanding during this study.

Last but not least, I would like to thank my lovely husband, Zulkarami bin Berahim and daughter Zainatul Insyirah for always being with me and loving me.

Thank you.



I certify that an Examination Committee met on [15 September 2008] to conduct the final examination of Zuraidah binti Zanirun on her Master of Science thesis entitled “Optimisation of Lignin peroxidase Production from *Pycnoporus* sp.” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the examination Committee are follows:

Chairman, PhD

Professor
Faculty of
Universiti Putra Malaysia
(Chairman)

Examiner 1, PhD

Professor
Faculty of
Universiti Putra Malaysia
(Member)

Examiner 2, PhD

Professor
Faculty of
Universiti Putra Malaysia
(Member)

Independent Examiner 1, PhD

Professor
Faculty of
Universiti Putra Malaysia
(Independent Examiner)

Professor/Deputy Dean of Graduate School
Universiti Putra Malaysia

Date:



The thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as the fulfillment of requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

SURAINI ABD AZIZ, Ph.D.

Associate Professor
Department of Bioprocess Technology
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

MOHD. ALI HASSAN, Ph.D.

Professor
Department of Bioprocess Technology
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

FOO HOOI LING, Ph.D.

Associate Professor
Department of Bioprocess Technology
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

HASANAH MOHD. GHAZALI, PhD
Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 15 JANUARI 2009



DECLARATION

I declare that the thesis is my original work except for the quotation and citation which have been duly acknowledged. I also declare that it has not been previously and is not concurrently submitted for any other degree at UPM or other institution.

Name: Zuraidah bt Zanirun

Date:



TABLE OF CONTENTS

ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL SHEETS	vii
DECLARATION FORM	ix
LIST OF TABLES	x
LIST OF FIGURES	xi
LIST OF ABBREVIATIONS	xiii
TABLE OF CONTENT	xiv
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 Biosynthesis and Structure of Lignin	
2.1.1 Structure of Lignin	4
2.1.2 Biosynthesis of Lignin	7
2.2 Lignin Biodegradation	7
2.2.1 Lignin Degrading Fungi	8
a) Soft Rot Fungi	9
b) Brown Rot Fungi	9
c) White Rot Fungi	10
d) Other	11
2.2.2 Lignin Degrading Enzymes	11
a) Lignin Peroxidase (LiP)	12
b) Manganese Peroxidase (MnP)	14
c) Laccase	16
d) Hydrogen Peroxide Producing Enzymes	17
2.3 Physiological Conditions for Lignin Degradation	17
2.3.1 Lignin Degradation as a Metabolic Stage	17
2.3.2 Lignin as a Growth Substrate	19
2.3.3 Role of Nitrogen and Carbon	19
2.3.4 Other Culture Parameters	20



2.4	Applications of Enzyme and White Rot Fungi	21
2.4.1	Bioremediation	21
2.4.2	Xenobiotics Degradation	23
2.4.3	Biopulping and Biobleaching	24
2.4.4	Bioconversion of Agro-Wastes	24
2.5	Optimisation of Lignin Peroxidase Production Through Factorial Design	25
3	MATERIALS AND METHODS	27
3.1	Materials	
3.1.1	Microorganisms	27
3.1.2	Source of Chemicals and Reagents	27
3.2	Production of Lignin peroxidase	28
3.2.1	Inoculum Preparation	28
3.2.2	Culture Media	28
3.2.3	Growth Condition of Spore Inoculum	29
3.2.4	Sample Assays	30
	a) Lignin peroxidase Activity	30
	b) Soluble Protein Concentration	31
	c) Nitrogen concentration	32
	d) Glucose Oxidase Activity	32
	e) Reducing Sugar Concentration	34
3.3	Experimental Procedure	35
3.3.1	Overview of Experimental Procedure	35
3.3.2	Preliminary Screening for The Highest Lignin Peroxidase Producer	35
3.3.3	Preliminary Optimisation Prior to Factorial Design	36
3.3.4	Optimisation of Lignin Peroxidase Production by Factorial Design	37
4	RESULTS AND DISCUSSION	41
4.1	Preliminary Screening For The Highest Lignin Peroxidase Producer	41
4.1.1	Growth on Solid Medium	41
4.1.2	Growth in Liquid Medium	43
	a) Agitated Culture	43
	b) Non-Agitated Culture	45
4.1.3	Comparison of Lignin Peroxidase Production in Agitated and Non-Agitated Cultures	47
4.1.4	Effect of Carbon Limited Medium on Growth of S.16	54
4.1.5	Microbial Identification	57



4.2	Optimisation of Lignin Peroxidase Production by Factorial Design Using <i>Pycnoporus</i> sp.	58
4.2.1	Preliminary Determination of Optimum Temperature	58
4.2.2	Determination on the Effect of Lignin as a Growth Substrate	60
4.2.3	Statistical Design Approach	62
	a) Lignin Peroxidase Production: Effects of Variables With and Without the Presence of Inducer	67
	b) Lignin Peroxidase Production by <i>Pycnoporus</i> sp. Using Optimised Condition Obtained	74
5	CONCLUSIONS AND RECOMMENDATION FOR FUTURE WORKS	
5.1	Conclusion	79
5.2	Recommendation for Future Works	80
	REFERENCES	81
	APPENDICES	96
	BIODATA OF STUDENT	107

Deleted: ¶



LIST OF TABLES

Table		Page
3.1	Factors and levels analysed in factorial experiment	38
3.2	Fractional factorial design of 5 variables in coded units	39
4.1	Fractional factorial design of 5 variables in coded units along with the observed response (Lignin peroxidase Activities)	63
4.2	Analysis of variance (ANOVA) for the production of lignin peroxidase enzyme	65
4.3	Coefficient estimate and P-value calculated from the model	67



LIST OF FIGURES

Figure		Page
2.1	Phenylpropanoid precursors of lignin	5
2.2	Schematic formula for a section of spruce lignin	6
2.3	Shikimic acid pathway for biosynthesis of lignin	7
2.4	Catalytic cycle of lignin peroxidase	14
2.5	Catalytic cycle of manganese peroxidase	15
2.6	Cycle of veratryl alcohol as a secondary metabolite	18
3.1	Overview of the experimental procedure	35
4.1	Fungal growth on malt extract agar plate	42
4.2	Mycelial pellet formed in agitated cultures	44
4.3	Mycelial mat formed in non-agitated cultures	46
4.4	Lignin peroxidase activity profiles in agitated culture	48
4.5	Lignin peroxidase activity profiles in non-agitated culture	49
4.6	Profiles of soluble protein concentration in agitated culture	52
4.7	Profiles of soluble protein concentration in non-agitated culture	53
4.8	Correlation of glucose consumption and lignin peroxidase in agitated culture condition	54
4.9	The effect of different concentration of glucose on the culture morphology	55
4.10	The effects of different glucose concentration on the production of lignin peroxidase activity	57
4.11	Lignin peroxidase production in different temperature	59



4.12	Profiles of lignin peroxidase activity with and without the present of lignin compound	61
4.13	Response surface of lignin peroxidase activity as a function of nitrogen concentration and agitation speed while keeping initial pH and inoculum concentration at centre point.	69
4.14	Response surface of lignin peroxidase activity as a function of inoculum concentration and initial pH while keeping nitrogen concentration and agitation speed at centre point.	69
4.15	Mycelial pellet formed at different agitation speed	71
4.16	Predicted optimum condition of Lignin peroxidase production by <i>Pycnoporus</i> sp.	74
4.17	Profile of lignin peroxidase activity by <i>Pycnoporus</i> sp. using optimised condition	75
4.18	Profile of glucose oxidase activity by <i>Pycnoporus</i> sp.	76
4.19	Profile of soluble protein concentration by <i>Pycnoporus</i> sp.	77
4.20	Profile of nitrogen concentration by <i>Pycnoporus</i> sp.	78



LIST OF ABBREVIATIONS

DHP	Dehydrogenation polymerizate
LiP	Lignin Peroxidase
%	Percentage
H ₂ O ₂	Hydrogen peroxide
MnP	Manganese peroxidases
CuSO ₄	Copper Sulphate



CHAPTER 1

INTRODUCTIONS

The increasing expansion of agro-industrial activity over the last 40 years has led to the accumulation of a large quantity of lignocellulosic residues all over the world (Villas-Boas *et al.*, 2002). Agriculture has played and will continue to play an important economic role in Malaysia. Malaysian agriculture is traditionally based on crop productions, particularly rubber, palm oil, cocoa, pepper, rice and pineapple. By-products derived from the mentioned agricultural sub-sectors have and will continue to provide large quantities of valuable nutrients to sustain livestock, particularly ruminant production in Malaysia. However, they are known mostly containing lignocellulosic materials which is a complex structure of lignin. Removal of lignin is important in order for enzymatic hydrolysis to occur as it acts as a barrier to most of agricultural wastes.

Lignin is a three-dimensional heterogeneous polymer stored in the plant cell wall of all vascular plants. It is the most abundant renewable aromatic biopolymer in the biosphere. Owing to its recalcitrant nature, lignin is remarkably resistant to degradation by most microorganisms, an important factor limiting the rate of degradation of lignocellulosic materials. Filamentous fungi, primarily the white-rot fungi and related ~~basidiomycetous~~ fungi are known to be ~~the~~ most efficient terrestrial ~~micoorganisms~~, capable of catalyzing lignin biodegradation. Studies ~~on~~ lignin biodegradation are important for possible biotechnology application, since lignin polymers are major obstacles to the efficient

Deleted: basidiomycete

Deleted: microorganism

Deleted: of



utilization of lignocellulosic materials in a wide range of industrial processes (Eriksson *et al.*, 1990).

Although removal of lignin can be done using chemical and physical pre-treatment, it was not a natural process as lignin compound and chemical reagent itself will cause environmental pollution. Therefore, biotechnology methods that are environmental friendly are preferred as a tool in any processes.

Various white-rot fungi isolated have the ability to degrade lignin by producing extracellular oxidative enzymes known as ligninase, a generic name for a group of isoenzymic peroxidases that catalyze the oxidative depolymerization of lignin. Another two extracellular ligninolytic peroxidases (lignin peroxidase, LiP, EC 1.11.1.14 and manganese peroxidase, MnP, EC 1.11.1.13) and one phenoloxidase of laccase type (benzenediol:oxydoreductase, EC 1.10.3.2) also have been intensively studied in various white-rot fungi for lignin biodegradation and dye decolourisation (Maganhotto de Souza Silva *et al.*, 2005) . In this study, attempt ~~had~~ been made to focus on lignin peroxidase enzyme production due to it capability to catalyze the oxidation of variety of compounds with high reduction potential (Viral *et al.*, 2005). This enzyme was produced by the fungi during secondary metabolism in response to environmental stress and the level of nutrient condition. Thus, several factors ~~were believed~~ can optimize the capability and potential of ~~a~~ white rot fungus producing lignin peroxidase.

Deleted: has

Deleted: are widely believed

Prior to achieving the objectives listed below, locally isolated fungus was screened to obtain the best lignin peroxidase producer. The potential lignin peroxidase producer was used for further study. Thus, the objectives of this study were:

- 1) To screen for the best lignin peroxidase producer from locally isolated white rot fungus
- 2) To optimise the parameters involved in lignin peroxidase production by locally isolated white rot fungus



CHAPTER 2

LITERATURE REVIEW

2.1 Structure and Biosynthesis of Lignin

2.1.1 Structure of Lignin

Lignin is a complex, three-dimensional, phenolic, natural polymer that is responsible for providing structural support to woody plants (Higuchi, 1990; Lewis and Sarkanen, 1998). The most abundant source of carbon is plant biomass, comprised primarily of cellulose, hemicellulose and lignin. Many microorganisms are capable of degrading and utilising cellulose and hemicellulose as carbon and energy sources, however a much smaller group of filamentous fungi has evolved with the ability to breakdown lignin, the most recalcitrant component of plant cell walls. Formed through oxidation and free radical coupling of phenyl alcohol precursors, the insoluble polymer lacks stereoregularity. In contrast to hydrolysable bonds between subunit of other wood polymers (e.g. cellulose and hemicellulose), lignin degradation requires oxidative attack on the carbon-carbon and ether interunit bonds. The lignin polymer encrusts cellulose microfibrils, particularly within the secondary walls. No microbe, including white rot fungi is known to be capable of utilising lignin as a sole carbon or energy sources, and it is believed that lignin depolymerization is necessary to gain access to cellulose and hemicellulose. Extracellular peroxidases and oxidases are thought to play an important role in the initial depolymerisation of lignin and small molecular weight fragments are subsequently

metabolised intracellularly ultimately to water and carbon dioxide (Cullen and Kersten, 2004).

Lignin is biosynthesized by the polymerization of phenylpropanoid precursors initiated by the enzymes peroxidase (Higuchi, 1985) or laccase (Sterjiades *et al.*, 1992, 1993). There are three of these precursors, differing in the number of methoxyl groups on the aromatic ring (Figure 2.1). Softwood lignin contains mostly guaiacyl (monomethoxy) units. Hardwood lignin contains roughly equal amounts of guaiacyl and syringyl (dimethoxy) units. Grass lignins contain p-hydroxyphenyl (no methoxyl) units as well as the other two types (Reid, 1995).




Figure 2.1: Phenylpropanoid precursors of lignin (Grabber, 2005)

Lignin polymerization occurs in the cell walls after the polysaccharides have been deposited and is initiated by a peroxidase- or laccase-mediated one-electron oxidation of the phenylpropanoid precursors to phenoxy radicals. The radicals couple with each other

and with the radicals in the growing lignin polymer in a random fashion to form a complex, cross-linked network (Figure 2.2) of these linkages, six to eight are more important. The single most predominant is the β -O-4 type between units 1 and 2, 2 and 3, 4 and 5, 6 and 7, 7 and 8, and 13 and 14 and constitutes 40 - 60% of the total linkages in different lignins (Adler, 1977). The other linkages are relatively less numerous and include phenylcoumaran (between units 3 and 4; 6 - 12%), biphenyl (units 5 and 6; 5 - 10%), diarylpropane (units 8 and 9; 5 - 10%), diphenyl ether (units 8 and 10; 4 - 6%), α -aryl ether (units 3 and 13, 15 and 16; 6 - 8%) and pinoresinol (units 10 and 11; < 5%).




Figure 2.2: Schematic formula for a section of spruce lignin (Adler, 1977)

2.1.2 Biosynthesis of Lignin

Biosynthesis of lignin starts with the fixation of carbon dioxide in plants to form carbon dioxide in plants to form carbohydrates. These are further metabolized via the shikimic acid pathway (Figure 2.3) to tyrosine and phenylalanine (Eriksson *et al.*, 1990).




Figure 2.3: Shikimic acid pathway for biosynthesis of lignin

2.2 Lignin biodegradation

Lignin is an insoluble, high molecular weight polymer, so the initial steps in its biodegradation by white rot fungi must be extracellular (Feijoo *et al.*, 1995). The final steps in lignin mineralization, ending with the release of CO₂, are likely to occur inside the fungal hyphae. The extracellular reactions must attack lignin in such a way that

fragments were produced and able to diffuse to the hyphae and cross the cell membranes. Previous work in the 1970s unravelled some of the complicated reactions which occur during the extracellular metabolism of the lignin polymer. The extracellular reactions were found to be predominantly oxidative, since the oxygen content of white rotted lignin is generally higher than the control lignin from sound wood (Kirk and Chang, 1975).

There is also a decrease in methoxyl groups and an increase in carbonyl and carboxyl groups. The chemical changes produced by white rot fungi in the lignin polymer include oxidative cleavage of the propanoid side chains between the α - and β -carbons and between the β - and γ -carbons, and demethylation and oxidative cleavage of aromatic rings (Chen and Chang, 1985). A progressive depolymerization occurs and releases a wide array of low molecular weight lignin fragments (Chen *et al.*, 1982;1983). Lignin biodegradation does not only occur by an orderly removal of the peripheral subunits as single ring compounds; it also involves oxidation of the aromatic rings and side chains in the interior of the polymer, increasing the hydrophilicity and solubility of the polymer core at the same time as fragments of varying sizes are released (Reid, 1995).

2.2.1 Lignin degrading fungi

The only organisms convincingly shown to efficiently degrade lignin are the wood rotting fungi (Eriksson *et al.*, 1990; Blanchette, 1991; Daniel, 1994). They can be divided into three groups according to the morphology of wood decay.