



UNIVERSITI PUTRA MALAYSIA

**BIOLOGY AND ECOLOGY OF DOG CONCH (*Strombus canarium*
Linnaeus, 1758) (GASTROPODA: STROMBIDAE) FROM MERAMBONG
SHOAL, JOHOR STRAITS, MALAYSIA**

ZAIDI CHE COB

FS 2008 41

BIOLOGY AND ECOLOGY OF DOG CONCH
(Strombus canarium Linnaeus, 1758)
(GASTROPODA: STROMBIDAE) FROM
MERAMBONG SHOAL, JOHOR STRAITS,
MALAYSIA

ZAIDI CHE COB

DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA

2008



**BIOLOGY AND ECOLOGY OF DOG CONCH (*Strombus canarium*
Linnaeus, 1758) (GASTROPODA: STROMBIDAE) FROM MERAMBONG
SHOAL, JOHOR STRAITS, MALAYSIA**

By

ZAIDI CHE COB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of
Doctor of Philosophy**

August 2008



DEDICATION

To my parents who always kept praying for me to achieve my goal, especially to my late father who passed away on January 20th 2008. May Allah bless his soul.

Al-Fatihah.

To my wife and my children who have sacrificed so much and be patient with me throughout the course of my study, and for me to be whom I am today.

To all my friend and colleagues who support me all those past years, and to technical staffs of Universiti Putra Malaysia, who provides technical support during my study.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Doctor of Philosophy

**BIOLOGY AND ECOLOGY OF DOG CONCH (*Strombus canarium*
Linnaeus, 1758) (GASTROPODA: STROMBIDAE) FROM MERAMBONG
SHOAL, JOHOR STRAITS, MALAYSIA**

By

ZAIDI CHE COB

August 2008

Chairman: Associate Professor Aziz Arshad, PhD

Faculty: Faculty of Science

Strombus canarium Linnaeus, 1758, locally known as ‘siput gonggong’ is a prosobranch gastropod from the family Strombidae that was highly associated with seagrass bed ecosystem. The main objectives of this study were to investigate and document the life history characteristics, biology, and ecology of the species, which are currently not well established.

This study has been conducted at Merambong Shoal, Johor Straits, Malaysia. Sample collections were conducted using a belt transect method where all individuals within the transect area were collected. The population dynamic was analyzed using both the Electronic Length Frequency Analysis (ELEFAN) and Length-based Fish Stock Assessment (LFSA) approaches, incorporated in the FiSAT software package. Apart from field observation, field experimentation and



laboratory analyses, laboratory culture experiment (from egg mass to juvenile) was also conducted, using a standard larval culture protocol.

Strombus canarium produced a long gelatinous tube of egg strand in the form of an egg mass. It took between 4.5 to 5.5 days of incubation before hatching, and the planktotrophic larvae spent between 18 to 24 days in the planktonic stage before metamorphosed. Metamorphic cue associated with bio-active materials from sediment taken from its nursery habitat showed the strongest effects compared to sediment conditioned seawater, seagrass detritus, seagrass detritus leachate, Potassium Chloride (KCL) and γ -aminobutyric acid (GABA). Changes of behavioral patterns related to metamorphosis were described, which can be divided into 3 phases: initiation, actual metamorphosis, and completion, resulted in completely camouflaged juveniles.

In their natural habitat, juveniles can be found at specific localities (nurseries), characterized by low to medium density *Halophila* bed; high sediment organic content; high sorting coefficient, and low mean grain size. The adult population was widely distributed, but present in local colonies, and was highly seasonal in abundance. Based on the gonadosomatic index (GSI), combined with other biological, ecological and behavioral observations, it could be concluded that the species was highly seasonal in reproduction, which starts from November to late March.

The population showed sexual polymorphisms where 3 different sexual morphs present i.e., males, normal females, and imposex females. Imposex was characterized by superimposition of male sexual characters on female. The imposex females showed significantly larger and heavier shell, and higher degree of outer-lip flaring compared to males and normal females. Imposex severity was defined as Stage 0 without the male genital system; Stage 1 with the appearance of rudimentary penis, but without penis duct; Stage 2 with simple prong of penis and penis duct, but without accessory; and Stage 3 with male penis complete with accessories. Development of other male characters as outlined in the general scheme of imposex evolution in prosobranchs was never observed. The percentage incidence of imposex (%I) was lower compared to other gastropod within the study areas, and there was no case of female sterility observed.

In population dynamic studies, a total of 766 males and 1322 females have been sampled from January to December 2005. Growth parameter estimation using both the ELEFAN-I and LFSA methods showed higher asymptotic length (L_{∞}) value in females compared to the males, but there was no significant difference in growth constant (K) and age at zero-length (t_0) of both sexes. The mean growth rate during the first year growth was estimated at 2.66 ± 0.16 mm/month for male and 4.59 ± 0.59 mm/month for female, which was comparable with the actual growth recorded using mark-recapture technique. The growth performance index (ϕ') was slightly lower in male ($\phi'=3.48$) compared to the female ($\phi'=3.81$), but both was well within the values reported for other marine gastropods. Recruitment



patterns, together with observations on spawning and copulation suggest that the species was highly synchronized (seasonal) in reproduction. The production and biomass was also seasonal, significantly higher during the wet monsoon season compared to the dry season.

In conclusions, many aspects of life history, biology, and ecology of *S. canarium* had been established. However more studies are urgently needed for better understanding and better management of the species, and for advancement of the species into mariculture.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

BIOLOGI DAN EKOLOGI SIPUT GONGGONG (*Strombus canarium* Linnaeus, 1758) (GASTROPODA: STROMBIDAE) DARI BETING MERAMBONG, SELAT JOHOR, MALAYSIA

Oleh

ZAIDI CHE COB

Ogos 2008

Pengerusi: Profesor Madya Aziz Arshad, PhD

Fakulti: Fakulti Sains

Strombus canarium Linnaeus, 1758, yang dikenali oleh penduduk tempatan sebagai ‘siput gonggong’ adalah gastropoda daripada Famili Strombidae yang mempunyai hubungan sangat rapat dengan ekosistem rumput laut. Objektif utama kajian ini adalah untuk menyiasat dan mendokumentasikan sejarah hidup, aspek biologi, dan ekologi spesis tersebut, yang sekarang ini maklumatnya agak terhad.

Kajian ini telah dijalankan di Beting Merambong, Selat Johor, Malaysia. Pengumpulan sampel dilakukan menggunakan kaedah jalur transek di mana semua individu di dalam kawasan transek dikumpulkan. Dinamik populasi dianalisa menggunakan kaedah ‘Electronic Length Frequency Analysis’ (ELEFAN) dan juga ‘Length-based Fish Stock Assessment’ (LFSA), yang terkandung dalam pakej perisian FiSAT. Selain daripada pemerhatian dan



eksperimentasi di lapangan serta analisis makmal, kultur makmal larva (dari telur hingga juvenil) juga dilakukan, menggunakan kaedah piawai bagi kultur larva.

Strombus canarium menghasilkan telur yang tersusun dalam satu tiub bergelatin yang panjang, membentuk satu gumpalan jasad telur. Ia memerlukan tempoh pengeraman dari 4.5 hingga 5.5 hari untuk menetas, menghasilkan larva planktotrofik yang hidup sebagai plankton selama 18 hingga 24 hari, sebelum menjalani proses metamorfosis. Induksi metamorfosis oleh bahan bio-aktif yang terdapat dalam sedimen dari habitat asuhan mempunyai kesan paling tinggi berbanding air laut yang diaktifkan dengan sedimen, detritus rumput laut, air peraman detritus rumput laut, Potassium Chloride (KCL) dan γ -aminobutyric acid (GABA). Perubahan kelakuan yang berkait dengan metamorfosis juga telah dikenalpasti, dan boleh dibahagikan kepada tiga fasa iaitu: inisiasi (permulaan), metamorfosis, dan fasa pengakhiran, menghasilkan juvenil yang mempunyai kesan penyamaran yang baik dengan kawasan sekelilingnya.

Di habitat semulajadinya, juvenil boleh dijumpai di lokaliti yang spesifik, yang dicirikan oleh: kawasan rumput laut *Halophila* yang berkepadatan rendah ke kepadatan sederhana; mempunyai nilai 'sorting coefficient' sedimen yang tinggi, dan purata saiz butiran sedimen yang rendah. Populasi dewasa didapati bertaburan secara meluas, tapi hadir dalam kelompok-kelompok koloni setempat, dan kelimpahannya sangat dipengaruhi oleh musim. Berdasar kepada indeks gonadosomatik (*GSI*), disamping ciri-ciri biologi, ekologi, dan aspek kelakuan,

spesis kajian boleh dirumuskan mempunyai corak pembiakan yang juga sangat di pengaruhi oleh musim, bermula dari November hingga Mac.

Populasi *S. canarium* menunjukkan fenomena polimorfisma seksual di mana terdapat 3 kumpulan morfologi berbeza iaitu jantan, betina normal dan betina imposex. Imposex mempunyai ciri kehadiran organ seksual jantan secara abnormal pada individu betina. Betina imposex mempunyai cangkerang yang secara signifikannya lebih besar dan lebih berat, dan darjah pengembangan bibir cangkerang yang lebih tinggi berbanding jantan dan betina normal. Tahap imposeks terhadap spesies dikelaskan kepada: Tahap 0 tanpa sebarang organ seksual jantan; Tahap 1 mempunyai penis rudimen, tanpa duktus; Tahap 2 mempunyai penis ringkas, berserta duktus, tapi tanpa aksesori; dan Tahap 3 mempunyai penis lengkap, berserta dengan duktus dan aksesori. Selain daripada itu peratus insiden imposex (%I) adalah rendah berbanding insiden ke atas gastropod lain di kawasan kajian, dan tidak ada kes betina steril yang di jumpai.

Dalam kajian dinamik populasi, sejumlah 766 jantan dan 1322 betina telah di sampel dari bulan Januari hingga Disember 2005. Penentuan parameter tumbesaran menggunakan ELEFAN-I dan LFSA menunjukkan nilai panjang asimptot (L_{∞}) yang lebih tinggi bagi betina berbanding jantan, namun tiada perbezaan signifikan bagi nilai pemalar tumbesaran (K) dan umur pada panjang sifar (t_0). Purata kadar tumbesaran bagi tahun pertama dianggarkan pada 2.66 ± 0.16 mm/bulan untuk jantan, dan 4.59 ± 0.59 mm/bulan untuk betina, yang agak

sama berbanding dengan nilai sebenar diperolehi secara kaedah tanda-tangkap semula. Indeks prestasi tumbesaran (ϕ') sedikit rendah bagi jantan ($\phi'=3.48$) berbanding betina ($\phi'=3.81$), namun kedua-duanya masih berada di dalam julat nilai yang dilaporkan untuk gastropoda marin yang lain. Corak rekrutmen, bersama-sama dengan pemerhatian terhadap jumlah telur dan kopulasi mencadangkan yang spesies adalah sangat bermusim dalam pembiakan. Kadar penghasilan dan biomas juga menunjukkan sifat bermusim, di mana nilainya lebih tinggi pada musim hujan berbanding musim kering.

Sebagai kesimpulan, berbagai aspek riwayat hidup, biologi, dan ekologi *S. canarium* telah berjaya dikenalpasti. Walau bagaimanapun kajian lanjut amat diperlukan untuk lebih memahami spesis tersebut, disamping pengurusan mapan sumber tersebut dapat dilakukan, dan untuk membangunkan spesies ini dalam industri marikultur.

ACKNOWLEDGEMENTS

Praise to Allah the Almighty for the strength, health, and peace of mind He provided me throughout this study.

I would like to express my sincere gratitude to my supervisor, Assoc. Prof. Dr. Aziz Arshad for the support, continued guidance and valuable collaboration during the entire process of this thesis at Universiti Putra Malaysia. Without those quality and constructive supervision and guidance, this work may not have come to completion.

I would like to thank my thesis committee members for their advice and critics on this thesis, and for continuous constructive discussion and suggestions: Assoc. Prof. Dr. Japar Sidik Bujang and Assoc. Prof. Dr. Mazlan Abd. Ghaffar.

I am also grateful to several people who were instrumental to the development of this thesis: Perumal Kuppan for his advice during laboratory culture work, Azmi for his technical assistant, and last but not least Pak Long who provide the boat service to the study site.



I certify that an Examination Committee has met on 28 August 2008 to conduct the final examination of Zaidi Che Cob on his Doctor of Philosophy thesis entitled “Aspects of Biology and Ecology of *Strombus canarium* Linnaeus, 1758 (Gastropoda; Strombidae) From Merambong Shoal, Johore, Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the Doctor of Philosophy degree.

Members of the Examination Committee are as follows:

Ahmad Ismail, PhD

Associate Professor
Faculty of Science,
Universiti Putra Malaysia
(Chairman)

Jambari Hj. Ali, PhD

Professor
Faculty of Science,
Universiti Putra Malaysia
(Internal Examiner)

Siti Khalijah Daud, PhD

Associate Professor
Faculty of Science,
Universiti Putra Malaysia
(Internal Examiner)

Zulfigar Yasin, PhD

Professor
School of Biological Sciences
Universiti Sains Malaysia
(External Examiner)

HASANAH MOHD. GHAZALI, PhD
Professor and Deputy Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date: 23 October 2008



This thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirements for the degree of Doctor of Philosophy. The members of Supervisory Committee were as follows:

Aziz Arshad, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Japar Sidik Bujang, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Mazlan Abd Ghaffar, PhD

Associate Professor
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(Member)

AINI IDERIS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 13 November 2008



DECLARATION

I declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declared that it has not been previously and is not concurrently submitted for any other degree at Universiti Putra Malaysia or at any other institutions.

ZAIDI CHE COB

Date: 19 September 2008



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLES	xvii
LIST OF FIGURES	xxi
CHAPTER	
1 GENERAL INTRODUCTION	1
1.1 Background	1
1.2 Statement of the Problems	5
1.3 Objectives	8
2 LITERATURE REVIEW	9
3 GENERAL METHODOLOGY	59
3.1 Study Site	59
3.2 Climatic Condition	60
3.3 Sample Collection and Measurements	61
3.4 Population Dynamic Analysis	62
3.5 Laboratory Culture Method	63
3.6 Habitat Characteristics and Physico-chemical Parameters	64
3.7 Statistical Analysis	65
3.8 Results of Physico-chemical Parameters	66
4 EARLY LIFE HISTORY CHARACTERISTICS: EMBRYONIC, LARVAL DEVELOPMENT AND METAMORPHOSIS OF <i>Strombus canarium</i> LARVAE	68
4.1 Background	68
4.2 Materials and Methods	71
4.3 Results	82
4.3.1 Spawning and Egg Mass Description	82
4.3.2 Embryonic and Pre-Hatch Larval Development	85
4.3.3 Post-Hatched Larval Development	89
4.3.4 Metamorphic Competence and Metamorphosis	92
4.3.5 Larval Development Stages and Description	94
4.3.6 Behavioral Observations	101
4.3.7 Larval Development, Growth and Survivals	105
4.3.8 Settlement Cues Experiment	111



4.3.9 Juvenile Behavior and Growth	115
4.4 Discussion	121
4.5 Conclusions	141
5 BIOLOGY OF <i>Strombus canarium</i>	144
5.1 Background	144
5.2 Materials and Methods	149
5.3 Results	156
5.3.1 Allometry Analysis	156
5.3.2 Reproductive Biology	158
5.3.4 Imposex, Morphometry and Sexual Polymorphism	184
5.4 Discussion	194
5.5 Conclusions	203
6 ECOLOGY OF <i>Strombus canarium</i>	205
6.1 Background	205
6.2 Materials and Methods	210
6.3 Results	215
6.3.1 Monthly Density	215
6.3.2 Seasonal Variations in Abundance	218
6.3.3 Spatial Distribution	219
6.3.4 Feeding Ecology	222
6.3.5 Habitat Preferences	227
6.3.6 Nursery Habitats	233
6.4 Discussion	237
6.5 Conclusions	249
7 POPULATION DYNAMICS OF <i>Strombus canarium</i>	251
7.1 Background	251
7.2 Materials and Methods	253
7.3 Results	264
7.3.1 Growth Parameters (ELEFAN-I Method)	264
7.3.2 Growth Parameters (LFSA Method)	282
7.3.3 Growth Parameters: Mark-Recapture and Length-At-Age data	292
7.3.4 Production and Biomass	295
7.3.5 <i>Strombus canarium</i> Fishery	299
7.4 Discussion	303
7.5 Conclusions	311
8 GENERAL DISCUSSION AND CONCLUSION	313
REFERENCES	323
APPENDICES	358
BIODATA OF STUDENT	375
LIST OF PUBLICATIONS	376



LIST OF TABLES

Table		Page
2.1	Taxonomic classification of <i>Strombus canarium</i>	16
2.2	Nutrient content of <i>Strombus canarium</i> and comparisons with other mollusks	51
4.1	Natural and artificial inducers type and method of preparation.	78
4.2	Summary of the major events in embryonic developments of <i>Strombus canarium</i>	89
4.3	Day to appearance of development characteristics of <i>S. canarium</i> larvae cultured during the wet and dry season	91
4.4	<i>Strombus canarium</i> larval growth (shell length): comparison between culture experiments conducted during wet and dry season	107
4.5	Model coefficients for Gompertz regression to describe <i>Strombus canarium</i> larval growth using length-at-age data	108
4.6	Metamorphosis response of <i>Strombus canarium</i> larvae exposed to different metamorphosis inducers	112
4.7	Settlement cue experiment on larvae: Post-hoc analysis using Tukey's pairwise comparison	114
4.8	Mean shell length and growth rates of <i>Strombus canarium</i> juveniles reared in laboratory	120
4.9	Time (h) of embryonic development of <i>Strombus canarium</i> and other gastropod mollusk	124
4.10	Day to appearance of development characteristics of the larvae of <i>Strombus canarium</i> (current study) in comparisons with <i>Strombus gigas</i> and <i>Strombus pugilis</i>	128
5.1	Correlations between log-transformed values of ash-free dry-weight (AFDW) against shell-length (SL), and shell-width (SW) for <i>Strombus canarium</i>	158
5.2	Tukey's pairwise comparisons between monthly gonadosomatic index (<i>GSI</i>) values for male <i>Strombus canarium</i>	170



5.3	Comparisons of male and female <i>Strombus canarium</i> gonado-somatic index (<i>GSI</i>) within months	171
5.4	Correlation between characters associated with reproduction in <i>Strombus canarium</i>	174
5.5	Correlation between <i>Strombus canarium</i> gonado-somatic index (<i>GSI</i>) and environmental parameters	174
5.6	Tukey's pairwise comparisons on monthly condition factor (<i>K</i>) values of female <i>Strombus canarium</i>	177
5.7	Tukey's pairwise comparisons on monthly condition factor (<i>K</i>) values of male <i>Strombus canarium</i>	178
5.8	Comparison between male and female <i>Strombus canarium</i> condition factor (<i>K</i>) within months	178
5.9	Tukey's pairwise comparisons on monthly condition index (<i>CI</i>) values of female <i>Strombus canarium</i>	181
5.10	Comparisons between male and female <i>Strombus canarium</i> condition index (<i>CI</i>) within month	183
5.11	Pearson correlations on monthly biological indices of <i>Strombus canarium</i> (January to December 2005)	184
5.12	Percent incidence (%I) of imposex in <i>Strombus canarium</i>	185
5.13	Relative penis size index (<i>RPSI</i>) and relative penis length index (<i>RPLI</i>) of <i>Strombus canarium</i>	187
5.14	Canonical variates analysis (<i>CVA</i>) on <i>Strombus canarium</i> shell linear characters of all the groups pooled (males, females and imposex females)	191
5.15	Comparisons between morphometric characters of different morphs of matured <i>Strombus canarium</i>	193
6.1	Classes of dominance used to record macrophyte coverage	213
6.2	Tukey's pairwise comparisons for <i>Strombus canarium</i> monthly density	218
6.3	Pearson correlation between <i>Strombus canarium</i> density and environmental parameters	218



6.4	Field observations on <i>Strombus canarium</i> feeding preference	224
6.5	Percentages of food items found in the stomach of adult <i>Strombus canarium</i>	227
6.6	<i>Strombus canarium</i> density at each sampling station (dependent variable)	228
6.7	Benthic macrophyte characteristic of the sampling stations (independent variables, biotic)	229
6.8	Benthic habitat characteristics of the sampling stations (independent variables, abiotic)	230
6.9	Eigenvalues and percentage of variation described by each axes	231
6.10	Eigen vectors for variables used in PCA of environmental (seagrass habitat) data, indicating the strengths of the correlations between variables and the principle components (axes)	232
6.11	<i>Strombus canarium</i> nursery habitat: conch density and habitat characteristics	235
6.12	<i>Strombus canarium</i> nursery habitat: Eigenvalues and percentage of variance described by each axes	236
6.13	<i>Strombus canarium</i> nursery habitat: Eigen vectors for variables used in PCA of environmental data, indicating the strengths of the correlations between variables and the principle components (axes)	236
7.1	Monthly length-frequency distribution of male <i>Strombus canarium</i>	265
7.2	Monthly length-frequency distribution of female <i>Strombus canarium</i>	266
7.3	Identified age groups from length-frequency analysis of male <i>Strombus canarium</i>	283
7.4	Identified age groups from length-frequency analysis of female <i>Strombus canarium</i>	284



7.5	Specific cohort growth parameters for male <i>Strombus canarium</i> determined using ELEFAN-1	289
7.6	Specific cohort growth parameters for female <i>Strombus canarium</i> determined using ELEFAN-1	289
7.7	Mark and recapture data of <i>Strombus canarium</i>	293
7.8	Length and age of newly metamorphosed juvenile <i>Strombus canarium</i> reared in laboratory	294
7.9	Annual production and standing biomass of male <i>Strombus canarium</i> cohorts	296
7.10	Annual production and standing biomass of female <i>Strombus canarium</i> cohorts	297
7.11	Production (P , gAFDW/m ² /yr), biomass (\bar{B} , gAFDW/m ²) and P/\bar{B} ratio for <i>Strombus canarium</i> , and comparison with other gastropods	299
7.12	<i>Strombus canarium</i> monthly catch data for 2005/2006 collecting season	300
7.13	<i>Strombus canarium</i> fishery: Biological parameters of catch	302



LIST OF FIGURES

Figure		Page
1.1	<i>Strombus canarium</i> Linnaeus, 1758 adult and juvenile	4
2.1	Geographical distribution of <i>Strombus canarium</i>	19
2.2	Schematic drawing of a generalized <i>Strombus</i> anatomy	22
2.3	Gonad of an immature male <i>Strombus canarium</i>	24
2.4	Schematic drawing of a generalized female <i>Strombus</i> reproductive tract	27
3.1	Location of the study site, the Merambong Shoal, Johor Straits, Malaysia	60
3.2	Monthly rainfall data for southern Johor area	61
3.3	Salinity and temperature (January to December 2005)	66
3.4	Conductivity (bottom), January to December, 2005	67
3.5	Dissolved oxygen (bottom) January to December, 2005	67
4.1	Sediment taken from <i>Strombus canarium</i> nursery ground habitat that was used for juvenile culture, and close-up of the surface sediment	80
4.2	Juvenile culture vessel, with sediment taken from <i>Strombus canarium</i> nursery habitat used as bottom sediment	80
4.3	Juvenile <i>Strombus canarium</i> behavior: direct observation under the dissecting microscope	81
4.4	Spawning female with egg mass still attached	82
4.5	Fragment of egg strand with parts of outer membrane peeled	84
4.6	<i>Strombus urceus</i> egg mass	84
4.7	<i>Strombus canarium</i> embryo stages	86
4.8	<i>Strombus canarium</i> veliger stages	87

4.9	<i>Strombus canarium</i> egg stand during the hatching process	88
4.10	A highly competence <i>Strombus canarium</i> larvae	93
4.11	<i>Strombus canarium</i> larval operculum	94
4.12	<i>Strombus canarium</i> veliger Stage I	95
4.13	<i>Strombus canarium</i> veliger Stage II	96
4.14	<i>Strombus canarium</i> veliger Stage III	98
4.15	<i>Strombus canarium</i> veliger Stage IV	99
4.16	Newly metamorphosed juvenile <i>Strombus canarium</i>	101
4.17	<i>Strombus canarium</i> in the midst of metamorphosis	103
4.18	Juvenile <i>Strombus</i> moments after underwent metamorphosis	104
4.19	Juvenile <i>Strombus canarium</i> day-1, completely camouflaged with sediment particle, sand and debris	105
4.20	Shell length at age relationship for <i>Strombus canarium</i> larvae cultured during wet season, plotted using Gompertz model	108
4.21	Shell length at age relationship for <i>Strombus canarium</i> larvae cultured during dry season, plotted using Gompertz model	109
4.22	Growth rate at specific age intervals for <i>Strombus canarium</i> larvae cultured during wet season and dry season	109
4.23	Percent survivals of <i>Strombus canarium</i> larvae cultured during wet season and dry season	110
4.24	Percentage of metamorphosed <i>Strombus canarium</i> larvae after 24 h	114
4.25	<i>Strombus canarium</i> larvae feeding underneath the surface film	115
4.26	<i>Strombus canarium</i> with shell partly buried into sediment	116
4.27	<i>Strombus canarium</i> juvenile covered with sediment as a form of camouflaged	118
4.28	Development of <i>Strombus canarium</i> operculum	119



4.29	Power function plot of post-metamorphic <i>Strombus canarium</i> juvenile growth in laboratory culture condition	120
4.30	Growth rate at age intervals of <i>Strombus canarium</i> reared in laboratory.	121
5.1	The different measurements used for comparing shell morphology in <i>Strombus canarium</i> .	150
5.2	Shell length to weight (AFDW) relationship for male and female <i>Strombus canarium</i>	157
5.3	Shell width to weight (AFDW) relationship for male and female <i>Strombus canarium</i> .	158
5.4	<i>Strombus canarium</i> gonad	159
5.5	Male <i>Strombus canarium</i> reproductive organ	160
5.6	Schematic line drawing of <i>Strombus</i> verge, drawn from fresh specimen taken from Merambong Shoal	160
5.7	Female <i>Strombus canarium</i> reproductive organ (egg groove, vulva, capsule gland)	161
5.8	Probability of sex determination in female <i>Strombus canarium</i>	162
5.9	Probability of sex determination in male <i>Strombus canarium</i>	162
5.10	Early sexual development in <i>Strombus canarium</i>	163
5.11	Schematic line drawing of a matured <i>Strombus canarium</i> verge	164
5.12	Logit curve plot of percentage of mature <i>Strombus canarium</i> males for the respective shell length class	165
5.13	Logit curve plot of percentage of mature females <i>Strombus canarium</i> for the respective shell length class	165
5.14	Probit curve plot of probability of mature males <i>Strombus canarium</i> for the respective lip thick class	166
5.15	Probit curve plot of probability of mature females <i>Strombus canarium</i> for the respective lip thick class	166

