



**UNIVERSITI PUTRA MALAYSIA**

**CLONING AND EXPRESSION OF RECOMBINANT HUMAN  
EPIDERMAL GROWTH FACTOR IN *Escherichia coli***

**AHMAD FAIZAL ABDULL RAZIS**

**IB 2007 5**

**CLONING AND EXPRESSION OF RECOMBINANT HUMAN EPIDERMAL  
GROWTH FACTOR IN *Escherichia coli***

**By**

**AHMAD FAIZAL ABDULL RAZIS**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Master of Science**

**September 2007**



With the name of ALLAH,  
To my parents ABDULL RAZIS and SITI RUKIAH and my siblings for their  
encouragement and support.....



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**CLONING AND EXPRESSION OF RECOMBINANT HUMAN EPIDERMAL GROWTH FACTOR IN *Escherichia coli***

By

**AHMAD FAIZAL ABDULL RAZIS**

**September 2007**

**Chairman: Associate Professor Zarida Hambali, PhD**

**Institute: Institute of Bioscience**

The expression of recombinant hEGF (human epidermal growth factor) in *Escherichia coli* was conducted to produce hEGF with free of inclusion bodies and biologically active. The recombinant hEGF was constructed using sticky ends ligation and resulted in successful insertion of the hEGF gene into the multiple cloning sites of the pFLAG-ATS. This insertion was confirmed by restriction enzyme analysis, PCR (polymerase chain reaction) and DNA (deoxyribonucleic acid) sequencing with 100% homology. The recombinant hEGF was expressed in *E. coli* at 2, 4 and 6 hours with induction of 0.5 mM and 1.0 mM IPTG (isopropylthiogalactopyranoside). This recombinant was found to be expressed as periplasmic fraction and whole cell insoluble fraction as confirmed with SDS-PAGE (sodium dodecyl sulphate-polyacrylamide gel electrophoresis) and western blotting analyses. These analyses showed the molecular weight of hEGF was approximately 6.8 kDa. There was no significant difference in the production of hEGF when the expression was induced with different IPTG concentration; however, for each IPTG concentration, there was significant difference between 0 hour and all the post-induction hours. In addition, hEGF was found to be significantly higher in



periplasmic fraction as compared to the whole cell insoluble fraction (196.5 ng/ml as compared with 167 ng/ml at 2 hour, 175.7 ng/ml as compared with 115.3 ng/ml at 4 hour and 168.3 ng/ml as compared with 140 ng/ml at 6 hour). Growth-stimulating activity of periplasmic hEGF was studied using BrdU (bromodeoxyuridine) cell proliferation assay and MTT (3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide) cell proliferation assay. The periplasmic hEGF as compared to the standard hEGF was found to be biologically active and showed similar activity. There were significant findings of periplasmic hEGF in stimulating the growth of HEK (human epidermal keratinocytes) at 24, 48 and 72 hours of incubation. Even at the highest concentration of periplasmic hEGF (10 ng/ml), the growth-stimulating activity still occurred and incubation at 48 hour resulted with the highest stimulation (85.4% at 10 ng/ml, 77.7% at 1 ng/ml, 70.7% at 0.1 ng/ml, 55.9% at 0.01 ng/ml and 37.6% at 0.001 ng/ml). Besides, MTT cell proliferation assay of periplasmic hEGF on HDF (human dermal fibroblasts) showed significant increased in the growth-stimulating activity when the duration of incubation increased. Highest percentage of HDF growth was found at 72 hour incubation as compared with 24 and 48 hours incubation. In conclusion, the findings showed recombinant hEGF was successfully expressed in *E. coli* and the growth-stimulating activity of hEGF was determined.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**PENGLONAN DAN PENGEKSPRESAN REKOMBINAN FAKTOR  
PERTUMBUHAN MANUSIA DALAM *Escherichia coli***

Oleh

**AHMAD FAIZAL ABDULL RAZIS**

**September 2007**

**Pengerusi: Profesor Madya Zarida Hambali, PhD**

**Institut: Institut Biosains**

Pengekspresan rekombinan faktor pertumbuhan manusia dalam *Escherichia coli* telah dijalankan untuk menghasilkan hEGF (human epidermal growth factor) bebas daripada badan inklusi dan aktif biologi. hEGF rekombinan telah dibentuk menggunakan kaedah percantuman hujung lekit dan hasilnya menunjukkan kemasukan gene hEGF ke pelbagai tapak klon pFLAG-ATS. Kemasukan gene ini telah dibuktikan melalui analisa enzim pembatas, PCR (polymerase chain reaction) dan penjujukan DNA (deoxyribonucleic acid) menunjukkan homologi 100%. hEGF rekombinan telah diekspres dalam *E. coli* pada jam 2, 4 dan 6 dengan diinduksi oleh 0.5 mM dan 1.0 mM IPTG (isopropylthiogalactopyranoside). Rekombinan ini didapati telah diekspres sebagai fraksi periplasmik dan fraksi keseluruhan sel tak larut yang telah dibuktikan dengan analisa SDS-PAGE (sodium dodecyl sulphate-polyacrylamide gel electrophoresis) dan kaedah 'western blotting'. Keputusan ini menunjukkan berat molekul hEGF lebih kurang 6.8 kDa. Tiada perbezaan penghasilan hEGF apabila diinduksi dengan berbeza kepekatan IPTG. Walau bagaimanapun, untuk setiap kepekatan IPTG terdapat perbezaan ketara penghasilan hEGF diantara induksi pada jam 0 dan semua jam pasca induksi. Tambahan lagi,



kepekatan hEGF didapati lebih tinggi dalam fraksi periplasmik berbanding dengan fraksi keseluruhan sel tak larut (196.5 ng/ml berbanding 167 ng/ml pada jam 2, 175.7 ng/ml berbanding 115.3 ng/ml pada jam 4 dan 168.3 ng/ml berbanding 140 ng/ml pada jam 6. Aktiviti rangsangan pertumbuhan oleh hEGF periplasmik telah dikaji menggunakan kaedah proliferasi sel BrdU (bromodeoxyuridine) dan MTT (3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide). hEGF periplasmik didapati aktif secara biologi dan mempunyai aktiviti yang sama apabila dibandingkan dengan Standard hEGF. Terdapat keputusan ketara menunjukkan hEGF periplasmik merangsang pertumbuhan HEK (human epidermal keratinocytes) pada jam 24, 48 dan 72 inkubasi. Aktiviti rangsangan pertumbuhan masih berlaku meskipun pada kepekatan tertinggi hEGF periplasmik digunakan (10 ng/ml) dan inkubasi pada jam 48 menunjukkan rangsangan yang tertinggi (85.4% pada kepekatan 10 ng/ml, 77.7% pada kepekatan 1 ng/ml, 70.7% pada kepekatan 0.1 ng/ml, 55.9% pada kepekatan 0.01 ng/ml dan 37.6% pada kepekatan 0.001 ng/ml). Disamping itu, kaedah proliferasi sel MTT pada hEGF ke atas HDF (human dermal fibroblasts) menunjukkan pertambahan ketara dalam aktiviti rangsangan pertumbuhan dengan pertambahan tempoh inkubasi. Peratusan pertumbuhan HDF adalah tertinggi pada jam 72 inkubasi dalam perbandingan dengan inkubasi pada jam 24 dan jam 48. Kesimpulannya, keputusan menunjukkan bahawa hEGF rekombinan telah berjaya diekspres dalam *E. coli* dan aktiviti rangsangan pertumbuhannya telah ditentukan.

## ACKNOWLEDGEMENTS

First of all, a lot of thanks to Utmost Allah the Almighty by His grace and bless, I have made this valuable accomplishment.

Secondly, I would like to express my extremely grateful and deepest gratitude to my supervisor Assoc. Prof Dr. Zarida Hambali for her guidance, suggestion, attention, criticism and patience throughout the course of this study.

My appreciation also goes to my co-supervisors Assoc. Prof Dr. Muhammad Nazrul Hakim Abdullah, Prof. Dr. Abdul Manaf Ali and Prof. Dr. Mohd Azmi Mohd Lila for their critical readings, discussion, comments and suggestions.

Additionally, I would like to acknowledge Assoc. Prof Dr. Saidi Moin for his assistance in statistical analyses, Mr. Kamaruddin, Dr. Lai Kit Yee and Dr. Zeenathul Nazariah for their help and guidance in laboratory works and generously sharing their knowledge.

I am grateful to all my friends at the Virology Laboratory, Faculty of Veterinary Medicine and Laboratory of Immunotherapeutics and Vaccines, Institute of Bioscience such as Mrs. Elysha, Suria, Hayati, Dr. Phong, Dr. Sandy, Kim Seng, Pong, Caryn, Lo, Tam, Kelvin, Nik Mohd Afizan, Mohd Razif, Fatema, Ram, Fadzlie Wong, and my housemates Radin Mohd Arif, Mohd Fazli, Azman, Huzairi, Termizi and Hafizuddin for their friendship.





Finally, I would like to acknowledge my financial sponsor, the Malaysian Ministry of Science, Technology and Innovation (MOSTI) for National Science Fellowship (NSF) and IRPA grant, without the assistance, this research would not have made possible.



I certify that an Examination Committee has met on 10<sup>th</sup> September 2007 to conduct the final examination of Ahmad Faizal Abdull Razis on his Master of Science thesis entitled "Cloning and expression of recombinant human epidermal growth factor in *Escherichia coli*" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Master of Science.

Members of the Examination Committee were as follows:

**Abdul Rahman Omar, PhD**

Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

**Raha Abd Rahim, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Latifah Saiful Yazan, PhD**

Lecturer  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Internal Examiner)

**Sahidan Senafi, PhD**

Associate Professor  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(External Examiner)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 22 November 2007



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

**Zarida Hambali, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Chairman)

**Muhammad Nazrul Hakim Abdullah, PhD**

Associate Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Member)

**Abdul Manaf Ali, PhD**

Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)

**Mohd Azmi Mohd Lila, PhD**

Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

---

**AINI IDERIS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 13 December 2007



## **DECLARATION**

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

**AHMAD FAIZAL ABDULL RAZIS**

Date: 6 November 2007



## TABLE OF CONTENTS

	<b>Pag e</b>
<b>DEDICATION</b>	ii
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	v
<b>ACKNOWLEDGEMENTS</b>	vii
<b>APPROVAL</b>	ix
<b>DECLARATION</b>	xi
<b>LIST OF TABLES</b>	xv
<b>LIST OF FIGURES</b>	xvi
<b>LIST OF ABBREVIATIONS</b>	xviii
<b>CHAPTER</b>	
1 <b>INTRODUCTION</b>	1
2 <b>LITERATURE REVIEW</b>	
2.1    Discovery and Characterization of Epidermal Growth Factor	4
2.2    Biosynthesis and Tissue Distribution of EGF	7
2.3    Binding and metabolism of hEGF	8
2.4    Biological Effects of hEGF	11
2.5    Genes and Structure of hEGF	13
2.6    Potential Clinical and Industrial Uses of hEGF	15
2.7 <i>E. coli</i> Expression System	17
2.8    Secretory production of recombinant proteins using <i>E. coli</i>	19
2.9    Enhancement of secretion efficiency by manipulating periplasmic chaperones	24
2.10   Expression of Recombinant hEGF in <i>E. coli</i>	26
2.11   Fusion proteins application	28
2.12   FLAG fusion proteins	29
2.13   Studies on the biological activities of hEGF	31
3 <b>CLONING AND SEQUENCING OF hEGF GENE</b>	
3.1    Introduction	33
3.2    Materials and Methods	35
3.2.1    Templates and Primers for Polymerase Chain Reaction (PCR)	35
3.2.2    Preparation of hEGF DNA Template	36
3.2.3    PCR Amplification of hEGF gene	36
3.2.4    Agarose Gel Electrophoresis	37
3.2.5    Purification of hEGF gene	37
3.2.6    Double Digestion of pFLAG-ATS and hEGF gene	38
3.2.7    Cloning of hEGF gene into a pFLAG-ATS	38
3.2.8    Bacterial Cell Transformation	41



3.2.9	Identification and Verification of Positive Clones	41
3.2.10	PCR Amplification of Positive Clones	42
3.2.11	Miniprep Plasmid Extraction	42
3.2.12	Restriction Enzyme Analysis	43
3.2.13	DNA sequencing of the hEGF Gene	44
3.2.14	Computer Analysis	44
3.3	Results	45
3.3.1	Amplification of hEGF Gene by the Polymerase Chain Reaction	45
3.3.2	Screening of Recombinant hEGF	46
3.3.3	The DNA Sequencing of the hEGF Gene	47
3.4	Discussion	49
4	<b>EXPRESSION, ISOLATION AND PURIFICATION OF hEGF</b>	
4.1	Introduction	54
4.2	Materials and Methods	55
4.2.1	Expression of Recombinant hEGF	55
4.2.2	Optimization of Expression of Recombinant hEGF	56
4.2.3	Isolation of FLAG Fusion Proteins	56
4.2.3.1	Periplasmic Fraction by Osmotic Shock	56
4.2.3.2	Whole Cell Extract: Soluble and Insoluble Fractions	57
4.2.4	Immuno-Affinity Purification of FLAG Fusion Proteins	57
4.2.4.1	Purification of FLAG Fusion Proteins with the ANTI-FLAG M2 Affinity Gel	57
4.2.4.2	Elution of the FLAG-fusion protein	58
4.2.4.3	Precipitation of FLAG-fusion protein	59
4.2.4.4	Removal of the FLAG peptide	59
4.2.5	SDS-PAGE analysis	60
4.2.5.1	Gel Preparation	60
4.2.5.2	Sample Preparation	61
4.2.5.3	Staining of SDS-PAGE	61
4.2.5.4	Western Blotting analysis	61
4.2.5.4.1	Preparation for Blotting of SDS-PAGE	61
4.2.5.4.2	Preparation of Trans-Blot Transfer System	62
4.2.5.4.3	Immunodetection of Transferred Protein	62
4.2.5.5	Dot-blot and Western blot analyses of purified recombinant hEGF protein	63
4.2.5.6	Quantification of recombinant hEGF protein	63
4.3	Results	64
4.3.1	Expression of recombinant hEGF	64
4.3.2	Optimization of Expression of recombinant hEGF	66

4.3.2.1	Periplasmic Fraction by Osmotic Shock Procedure	66
4.3.2.2	Whole Cell Extract: Insoluble Fraction	68
4.3.2.3	Effect of IPTG induction on the growth of recombinant hEGF and non-recombinant hEGF	71
4.3.2.4	Quantification of recombinant hEGF protein by Sandwich ELISA	73
4.3.3	Immuno-Affinity Purification of FLAG Fusion Protein	78
4.4	Discussion	81
5	<b>GROWTH-STIMULATING ACTIVITY OF hEGF ON CULTURES OF HUMAN EPIDERMAL KERATINOCYTES AND HUMAN DERMAL FIBROBLASTS</b>	
5.1	Introduction	94
5.2	Materials and Methods	96
5.2.1	Cell Culture	96
5.2.1.1	Human Epidermal Keratinocytes	96
5.2.1.2	Human Dermal Fibroblasts	96
5.2.2	Cell Counting by Hemocytometer	97
5.2.2.1	Cells preparation	97
5.2.2.2	Slide Preparation	97
5.2.2.3	Analysis	98
5.2.3	Growth-stimulating activities of hEGF	98
5.2.3.1	BrdU Cell Proliferation Assay on HEK	98
5.2.3.2	MTT Cell Proliferation Assay on HDF	99
5.2.4	Percentage of growth-stimulating activity	100
5.2.5	Statistical Analysis	100
5.3	Results	101
5.3.1	BrdU Cell Proliferation Assay on HEK	101
5.3.2	MTT Cell Proliferation Assay on HDF	107
5.4	Discussion	113
5.4.1	BrdU Cell Proliferation Assay on HEK	113
5.4.2	MTT Cell Proliferation Assay on HDF	116
6	<b>SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	120
	<b>REFERENCES</b>	131
	<b>APPENDICES</b>	144
	<b>BIODATA OF STUDENT</b>	175
	<b>LIST OF PUBLICATIONS</b>	176

## LIST OF TABLES

Table		Page
1	Amino acid sequence of Epidermal Growth Factor.	5
2	Signal sequences for the secretory production of recombinant proteins in <i>Escherichia coli</i> .	20
3	Quantification of recombinant hEGF protein as periplasmic fraction at different induction with 1.0 mM and 0.5 mM IPTG concentration.	74
4	Quantification of periplasmic fraction and whole cell insoluble fraction with induction of 1.0 mM IPTG.	77
5	Percentage of growth-stimulating activity of periplasmic hEGF on HEK after incubation at 24, 48 and 72 hours.	102
6	Percentage of growth-stimulating activity of periplasmic hEGF on HDF after incubation at 24, 48 and 72 hours.	108
7	Solutions for Preparing Resolving Gels for Tris-glycine SDS-Polyacrylamide Gel Electrophoresis.	146
8	Solutions for Preparing 5% Stacking Gels for Tris-glycine SDS- Polyacrylamide Gel Electrophoresis.	146



## LIST OF FIGURES

Figure		Page
1	Amino acid sequence of hEGF.	6
2	Diagram of F-hEGF interaction with A-431 cells.	11
3	A-C Strategies for the secretory production of recombinant proteins in <i>Escherichia coli</i> .	21
4	Physical map of the pFLAG-ATS vector.	40
5	PCR amplification of hEGF gene.	45
6	PCR screening on the colonies of transformant.	46
7	Restriction enzyme (RE) analysis on recombinant hEGF.	47
8	The DNA sequence of the hEGF gene as compared to the established hEGF sequence (hEGF).	48
9	SDS-PAGE analysis of recombinant hEGF protein after 2 hours induction with 1.0 mM IPTG.	65
10	Western-blot analysis of recombinant hEGF protein after 2 hours induction with 1.0 mM IPTG using monoclonal anti-hEGF produced in mouse.	65
11	SDS-PAGE analysis for periplasmic fraction of recombinant hEGF protein at 0, 2, 4 and 6 hours post-induction.	67
12	Western-blot analysis for periplasmic fraction of recombinant hEGF protein at 0, 2, 4 and 6 hours post-induction using monoclonal anti-hEGF produced in mouse (Sigma Aldrich, U.S.A.).	67
13	Western-blot analysis for periplasmic fraction of recombinant hEGF protein at 0, 2, 4 and 6 hours post-induction using ANTI-FLAG <sup>®</sup> mouse monoclonal antibodies M1 (IgG2b) (Sigma Aldrich, U.S.A.).	68
14	SDS-PAGE analysis for whole cell insoluble fraction of recombinant hEGF protein at 0, 2, 4 and 6 hours post-induction.	70
15	Western-blot analysis for whole cell insoluble fraction of recombinant hEGF protein at 0, 2, 4 and 6 hours pos-induction using monoclonal anti-hEGF produced in mouse (Abcam, U.S.A.).	70



<b>16</b>	Effect of IPTG induction on the growth of recombinant hEGF (pFLAG-EGF) and non-recombinant hEGF (pFLAG-ATS).	72
<b>17</b>	Effect of IPTG induction at different concentration on the growth of recombinant hEGF.	72
<b>18</b>	Interaction analysis between time of post-induction and different IPTG concentration used in the production of recombinant hEGF protein.	75
<b>19</b>	Comparison of periplasmic fraction with induction of 0.5 mM and 1.0 mM IPTG concentration.	76
<b>20</b>	The comparison between whole cell insoluble fraction and periplasmic fraction with induction of 1.0 mM IPTG concentration.	79
<b>21</b>	Dot-blot assay by chemiluminescent detection kit of purified recombinant hEGF protein for periplasmic fraction and whole cell insoluble fraction using monoclonal anti-hEGF produced in mouse (Abcam, U.S.A.).	79
<b>22</b>	SDS-PAGE analysis of purified recombinant hEGF protein for periplasmic fraction and whole cell insoluble fraction.	80
<b>23</b>	Western-blot analysis of purified recombinant hEGF protein for periplasmic fraction and whole cell insoluble fraction using monoclonal anti-hEGF produced in mouse (Abcam, U.S.A.).	80
<b>24</b>	Comparison of growth-stimulating activity between standard hEGF and periplasmic hEGF on HEK at 24 hour incubation.	104
<b>25</b>	Comparison of growth-stimulating activity between standard hEGF and periplasmic hEGF on HEK at 48 hour incubation.	105
<b>26</b>	Comparison of growth-stimulating activity between standard hEGF and periplasmic hEGF on HEK at 72 hour incubation.	106
<b>27</b>	Comparison of growth-stimulating activity between standard hEGF and periplasmic hEGF on HDF at 24 hour incubation.	110
<b>28</b>	Comparison of growth-stimulating activity between standard hEGF and periplasmic hEGF on HDF at 48 hour incubation.	111
<b>29</b>	Comparison of growth-stimulating activity between standard hEGF and periplasmic hEGF on HDF at 72 hour incubation.	112

## LIST OF ABBREVIATIONS

<sup>51</sup> Cr-labeled	<sup>51</sup> Chromium-labeled
Amp	Ampicillin
ARF	Acute Renal Failure
ATCC	American Type Culture Collection
BrdU	Bromodeoxyuridine
BSA	Bovine Serum Albumin
BUN	Blood Urea Nitrogen
C-terminal	Carboxyl-terminal
DAB	3, 3'-diaminobenzidine
<i>E. coli</i>	<i>Escherichia coli</i>
EBA	Expanded Bed Adsorption
EGF	Epidermal Growth Factor
EGF-R	Epidermal Growth Factor Receptor
fMet	<i>N</i> -formyl methionine
FTNF	Feline Tumor Necrosis Factor
HB-EGF	Heparin binding EGF-like growth factor
HDF	Human Dermal Fibroblasts
hEGF	Human Epidermal Growth Factor
HEK	Human Epidermal Keratinocytes
hOB	Human obese
K1 (SK)	Kringle 1 (Streptokinase)
MCS	Multiple cloning site
mEGF	Mouse Epidermal Growth Factor
MRC-5	Normal Human Fetal Lung Fibroblast



NHEK	Normal Human Epidermal Keratinocytes
NMWL	Nominal Molecular Weight Limit
N-terminal	Amino-terminal
OmpA	Outer membrane protein A
ORF	Open Reading Frame
Prepro	Protein precursor
rEGF	Rat Epidermal Growth Factor
T3	Triiodothyronine
TGF $\alpha$	Transforming Growth Factor alpha
TNF	Tumor Necrosis Factor

## CHAPTER 1

### INTRODUCTION

Human epidermal growth factor (hEGF), consisting of 53 amino acid residues, is a single chain polypeptide with a molecular weight of about 6,200 Da. hEGF is considered to be identical with human urogastrone (Gregory, 1975), which is produced by the human duodenum and salivary glands (Cohen and Carpenter, 1975). hEGF is widely used in clinical and cosmetic fields because it stimulates the growth of a variety of cell types in cultures as well as the growth and differentiation of certain tissues *in vivo* (Carpenter and Cohen, 1979). It also promotes protein synthesis, RNA synthesis and metabolic uptake in epidermal cells (Kim and Muller, 1999), repairs corneal wounds and gastric ulcers and inhibits the proliferation of human gastric cancer cells (Jiang *et al.*, 1999; Reim *et al.*, 1988). Besides, hEGF has been observed to have many biological actions both *in vitro* and *in vivo* including proliferative effects on fibroblasts, keratinocytes and epithelial cells.

Since the finding of hEGF properties as strong anti-ulcer (Gregory, 1975) and growth-stimulating factors (Hashimoto *et al.*, 1994), the sufficient production for medical application has been in great demand. Conventional method such as recovery of hEGF from human urine via complicated steps is established (Starkey *et al.*, 1975). However, it failed to provide sufficient amount of homogenous hEGF with well-defined quality for clinical application and inappropriate for human consumption.



Many studies have reported using recombinant DNA technique to produce high hEGF productivity. However, the cytoplasmic expression of heterologous protein in *E. coli*, particularly small peptides, has encountered many obstacles. It is commonly found that proteins expressed in this form are rapidly degraded by proteases within the cell (Skipper *et al.*, 1985). It was reported that the hEGF expressed in high density precipitated intracellularly form aggregates called inclusion bodies (Shimizu *et al.*, 1991). This causes the expressed proteins to be severely misfolded. Although there are many protocols available for refolding this protein, the production is inefficient and the procedure is costly.

Thus, to obtain hEGF in sufficient quantities for this study, the recombinant hEGF is developed. It is desirable that recombinant hEGF be secreted into the cell growth medium. Purification of this protein would then be simpler for an intracellular protein as the product would not be contaminated with cytoplasmic components. In addition, the formation of inclusion bodies would be avoided and possible toxic effects of the hEGF polypeptide product on the host cell would be reduced.

In this study, a secretory plasmid is used, which contains strong regulatory elements such as *tac* promoter, the consensus ribosome-binding site, and the *ompA* leader sequence for efficient expression and secretion of hEGF in *E. coli*. This plasmid also encodes a *lac* repressor and an ampicillin (Amp) resistance gene. It was shown that secretory recombinant *E. coli* system was able to express and accumulate authentic hEGF in the culture medium (Wong and Sutherland, 1993). Normally, the secretion of high-level expressed protein in *E. coli* will eventually elicit a stress response in the host such as energy drain, membrane perturbation and lethal effect which leads to the

phenomenon known as “overproduction lethality”. Andersson *et al.* (1996) suggested that overexpression of foreign protein was due to the appearance of non-culturable cells which do not lose all metabolic activities and even continued to maintain glucose uptake and respiratory ability. However, only few studies reported the optimization of conditions for the process control in the expression and secretion of hEGF in *E. coli*.

In the present study, recombinant hEGF is expressed as a fusion protein using pFLAG expression system (Sigma, U.S.A.). This system allows recombinant protein to be expressed as a fusion protein containing the FLAG peptide sequence. Monoclonal antibodies recognizing the FLAG peptide can then be used to purify the fusion protein using affinity chromatography. The purification of hEGF protein has been made more convenient and efficient by using this system.

The aim of this study is to express the recombinant hEGF in *E. coli* and study the growth-stimulating activity of hEGF on human epidermal keratinocytes and human dermal fibroblasts cultures. The objectives involved in this study were:-

1. cloning and sequencing of hEGF gene.
2. expression, isolation and purification of hEGF.
3. determination of the growth-stimulating activity of hEGF on human epidermal keratinocytes and human dermal fibroblasts cultures.

## CHAPTER 2

### LITERATURE REVIEW

#### 2.1 Discovery and Characterization of Epidermal Growth Factor

Epidermal growth factor (EGF) was discovered as a contaminant of nerve growth factor in 1960 by Stanley Cohen (Cohen, 1962). EGF was recognized by its ability to accelerate the eruption of mouse teeth and the opening of eyelids of newborn mice. The abundance of EGF in mouse salivary glands facilitated its purification and in 1972 the full amino acid sequence of mouse EGF was determined (Savage *et al.*, 1972). Although no human equivalent of EGF had been purified during that time, it was known that a similar protein was present in concentrates of human urine (Starkey *et al.*, 1975). The human urinary protein responsible for the inhibition of gastric acid secretion ( $\beta$ -urogastrone) was finally purified and analyzed. Harry Gregory (1975) recognized that the amino acid composition of murine EGF, human urinary EGF and  $\beta$ -urogastrone were closely related. The biology of urogastrone indicated that it might also promote the proliferation and epithelialization of gastric mucosa, so he concluded that urogastrone and human epidermal growth factor are one and the same (Gregory, 1975). The amino acid sequence of EGF is as displayed in Table 1.



**Table 1. Amino acid sequence of Epidermal Growth Factor**

---

EGF	AMINO ACID SEQUENCE
rat EGF	NSNTGCPPSYDGYCLNGGVCMYVESVDRYVCNCVIGYIGERCQHRDLR
mouse EGF	NSYPGCPSSYDGYCLNGGVCMHIESLDSYTCNCVIGYSGDRCQTRDLRWWELR
guinea pig EGF	QDAPGCPPSHDGYCLHGGVCMHIESLNTYACNCVIGYVGERCEHQDLDDWE
human EGF	NSDSECPLSHDGYCLHDGVCMYIEALDKYACNCVVGYIGERCQYRDLKWWELR

---

(Source: Burgess, 1989)

