



UNIVERSITI PUTRA MALAYSIA

**DIRECT RECOVERY OF RECOMBINANT HEPATITIS B CORE
ANTIGEN FROM UNCLARIFIED *ESCHERICHIA COLI* FEEDSTOCK
USING EXPANDED BED ADSORPTION CHROMATOGRAPHY**

MICHELLE NG YEEN TAN

IB 2008 5

**DIRECT RECOVERY OF RECOMBINANT HEPATITIS B CORE ANTIGEN
FROM UNCLARIFIED *ESCHERICHIA COLI* FEEDSTOCK USING
EXPANDED BED ADSORPTION CHROMATOGRAPHY**

By

MICHELLE NG YEEN TAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of
Doctor of Philosophy**

August 2008



Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**DIRECT RECOVERY OF RECOMBINANT HEPATITIS B CORE ANTIGEN
FROM UNCLARIFIED *ESCHERICHIA COLI* FEEDSTOCK USING
EXPANDED BED ADSORPTION CHROMATOGRAPHY**

By

MICHELLE NG YEEN TAN

August, 2008

Chairman: Associate Professor Dr. Tey Beng Ti, PhD

Institute: Bioscience

The capsid of hepatitis B virus (HBV), which consists of hepatitis B core antigen (HBcAg) has become one of the most frequently studied viral-like-particle (VLP) for the display of foreign epitopes. Many studies have been carried out to purify this capsid. However, the conventional method of purification requires multiple steps of operation, which could lead to excessive product loss and high production costs. Therefore, it is of importance to develop a fast and cost effective protein recovery method such as expanded bed adsorption chromatography (EBAC) to ensure a better and efficient recovery of protein, especially in large scale downstream process.

In this study, thermal treatment of the *Escherichia coli* cell feedstock at 60°C for 30 min prior to solid removal in the conventional method has resulted in 1.4 times and 18% higher in purity and recovery yield respectively compared to non-heat-treated feedstock. In direct capture of HBcAg from unclarified feedstock using the STREAMLINE DEAE (weak anion-exchangers) in batch adsorption, heat treatment at 60°C for 45 min has increased the recovery yield and purity by 2.3 and 3.8 times

respectively compared with non-heat-treated feedstock. When these conditions were applied in large scale purification of HBcAg via EBAC, the yield and purity have increased by 1.2 and 1.8 folds respectively, compared with that purified from non-heat-treated feedstock. Heating the crude feedstock has resulted in denaturation and precipitation of contaminants in the feedstock, hence reducing non-specific interactions between the cell debris and anion-exchanger. The present study has also demonstrated that purification of HBcAg from heat-treated unclarified feedstock was most efficient when EBAC operation using Fastline™ 20 contactor was operated at constant velocity (127.9 cm/h) in feedstock containing 5% of biomass. Although the current study showed that heat-treatment of unclarified feedstock could increase the purity of HBcAg and reduced non-specific binding of contaminant onto Streamline DEAE, the purity obtained was lower compared with that purified using conventional methods. Therefore, development of an affinity adsorbent using M13 phage bearing a disulfide constrained heptapeptide at the gpIII protein coat with the sequence, C-WSFFSNI-C as the ligand has been carried out in this study. M13 phage immobilised onto Streamline Base Matrix via epoxy activation was used in direct capture of HBcAg from unclarified feedstock via two different modes of EBAC operations; typical single pass operation and modified EBAC operation with recirculation of feedstock.

Higher yield of HBcAg was obtained using modified EBAC operation due to increase in protein residence time in the column, however, the purity was reduced by 15% compared with typical EBAC operation, which could be due to diffusion of contaminants into the internal volume of the macroporous adsorbents. Although the purity of HBcAg recovered using M13 phage ligand adsorbents was higher (70-80%)

but the yield was lower compared with that purified using anion-exchanger. Therefore, this study showed that peptide displayed on M13 phage can be employed as an affinity ligand in direct capture of HBcAg from unclarified feedstock using EBAC. When analysed with ELISA, the antigenicity of HBcAg purified using both adsorbents in EBAC was still preserved.

Abstrak tesis yang dikemukakan kepada Senat Univerisiti Putra Malaysia sebagai memenuhi keperluan ijazah Doktor Falsafah

**PEMULIHANAN SECARA LANGSUNG REKOMBINAN ANTIGEN TERAS
HEPATITIS B DARIPADA SUAPAN *ESCHERICHIA COLI* TANPA
PERJERNIHAN MENGGUNAKAN PENJERAPAN LAPISAN
TERKEMBANG KROMATOGRAFI**

Oleh

Michelle Ng Yeen Tan

Ogos, 2008

Pengerusi: Profesor Madya Dr. Tey Beng Ti, PhD

Institut: Biosains

Kapsid virus hepatitis B (HBV) yang terdiri daripada antigen teras (HBcAg) telah menjadi salah satu daripada partikel bak virus (VLP) yang kerap digunakan untuk mempamerkan epitop asing. Pelbagai kajian telah dijalankan untuk penulenan HBcAg, akan tetapi, kaedah konvensional memerlukan beberapa langkah penjernihan yang boleh mengakibatkan kehilangan protein, kapital dan kos buruh yang tinggi serta masa pemprosesan yang panjang. Oleh itu, kaedah penulenan yang berkesan, murah dan pantas seperti penjerapan lapisan terkembang kromatografi (EBAC) harus digunakan, terutamanya dalam pelaksanaan skala besar.

Dalam kajian ini, rawatan haba ke atas suapan *Escherichia coli* dalam penulenan konvensional pada suhu 60°C selama 30 min telah menyebabkan peningkatan ketulenan sebanyak 1.4 kali dan 18% dalam hasil akhir jika dibandingkan dengan suapan tanpa rawatan haba. Di dalam penulenan langsung HBcAg dari suapan tanpa penjernihan menggunakan Streamline DEAE (penjerap penukaran

anion lemah) dalam penjerapan kelompok, rawatan haba pada suhu 60°C selama 45 min telah meningkatkan hasil terakhir sebanyak 2.3 kali dan 3.8 kali dalam ketulenan HBcAg berbanding dengan suapan tanpa rawatan haba. Apabila ciri-ciri rawatan haba ini digunakan dalam penulenan HBcAg pada skala besar menggunakan EBAC, peningkatan sebanyak 1.2 dan 1.8 kali masing-masing dalam hasil terakhir dan ketulenan HBcAg berbanding dengan suapan tanpa rawatan. Rawatan haba ke atas suapan telah menyebabkan penyahhasilan dan pemendakan bendasing lalu mengurangkan interaksi di antara bendasing dengan penjerap penukaran anion. Kajian ini juga telah menunjukkan bahawa penulenan HBcAg daripada suapan tanpa penjernihan yang telah dirawat adalah lebih berkesan jika operasi EBAC menggunakan turus Fastline™ 20 dilaksanakan pada halaju malar (127.9 cm/j) dengan suapan tanpa penjernihan yang mengandungi 5% berat basah biojisim (berat/isipadu). Walaupun ketulenan HBcAg diperolehi daripada suapan yang telah melalui rawatan haba telah meningkat jika dibandingkan dengan tanpa rawatan, ketulennanya masih tidak setaraf dengan apa yang diperolehi dengan kaedah konvensional. Oleh itu, penjerapan affiniti menggunakan faj M13 yang membawa peptida C-WSFFSNI-C telah digunakan dalam pemulihan secara langsung HBcAg melalui EBAC. Faj M13 telah dipegunkan pada Streamline matrik basal melalui pengaktifan epoksi dan telah digunakan dalam dua operasi EBAC yang berlainan iaitu operasi biasa EBAC dan operasi EBAC yang telah diubahsuaikan dengan pengitaran semula suapan secara berterusan melalui lapisan terkembang dalam turus.

Walaupun hasil terakhir HBcAg melalui penulenan operasi EBAC yang telah diubahsuai adalah lebih tinggi berbanding dengan operasi biasa, tahap ketulenan yang diperolehi telah menurun sebanyak 15% akibat daripada resapan bendasing ke

isipadu dalaman penjerap bermakroliang. Ketulenan HBcAg yang diperolehi melalui penjerapan affiniti ini adalah lebih tinggi jika berbanding dengan penjerapan penukaran anion (melebihi 80% ketulenan) walaupun hasil terakhirnya adalah lebih rendah. Dengan itu, kajian ini telah membuktikan bahawa pembawa peptida faj M13 boleh digunakan sebagai penjerap affiniti dalam pemulihan secara langsung HBcAg melalui EBAC. Apabila HBcAg yang diperolehi melalui kedua-dua kaedah penjerapan dianalisis dengan menggunakan kaedah ELISA, tahap keantigenannya masih dikekalkan.

ACKNOWLEDGEMENTS

My deepest and sincere appreciation goes to my main supervisor Assoc. Prof. Dr. Tey Beng Ti, his patience, encouragement and invaluable advice have been my driving force to work the best throughout this project. My sincere appreciations are also extended to Assoc. Prof. Dr. Tan Wen Siang, whom his passion and enthusiasm in research have been an inspiration to me and also to Dr. Ling Tau Chuan and Dr. Norhafizah Abdullah for their invaluable advice and time for discussions.

I would also like to thank all the staff in the Microscopy Unit in Institute of Bioscience for their assistance in my microscopic work. Not forgetting to all my labmates in Lab 134, especially Swee Tin, Wei Boon, Kie Hie, my neighbour, Lip Nam, Senthil, Ho and others who have made my time spent in the lab more meaningful and interesting.

To my husband, Ricky, his understanding, love and my shoulder to lean on, has given me strength, courage and perseverance to go through rain or shine throughout this project, and to my wonderful parents and brothers, a big thank you to all of them for their encouragements, love and supports. I couldn't have done this without all of them.

Last but not least, to Kuok Foundation Berhad for funding my three-year study in UPM. This study was supported by the e-Science Fund (02-01-04-SF0486 and 03-01-04-SF0755) from Ministry of Science, Technology and Innovation of Malaysia (MOSTI).

I certify that an Examination Committee has met on 8 August 2008 to conduct the final examination of Michelle Ng Yeen Tan on her degree thesis entitled “Direct Recovery of Recombinant Hepatitis B Core Antigen from Unclarified *Escherichia coli* Feedstock Using Expanded Bed Adsorption Chromatography” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Doctor of Philosophy.

Members of the Examination Committee were as follows:

Janna Ong Abdullah, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Arbakariya Ariff, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Rosfarizan Mohamad, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Conan Fee, PhD

Professor

College of Engineering

University of Canterbury, New Zealand

(External Examiner)

HASANAH MOHD. GHAZALI, PhD

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 16 October 2008

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Tey Beng Ti, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Tan Wen Siang, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Ling Tau Chuan, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Norhafizah Abdullah, PhD

Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Member)

AINI IDERIS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 16 October 2008

DECLARATION

I declare that the thesis is my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

MICHELLE NG YEEN TAN

Date: 21 August 2008

TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	ii
ACKNOWLEDGEMENTS	v
APPROVAL	viii
DECLARATION	ix
LIST OF TABLES	xi
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xviii
	xxi

CHAPTER

1	INTRODUCTION	1
2	LITERATURE REVIEW	4
	2.1. Hepatitis B virus	4
	2.1.1. Virion structure	6
	2.2. Hepatitis B core antigen (HBcAg)	7
	2.2.1. HBcAg as carrier protein	11
	2.2.2. Production and purification of HBcAg	14
	2.3. Expanded bed adsorption chromatography (EBAC)	17
	2.3.1. Protein adsorption equilibrium	18
	2.3.2. Principles of operations in EBAC	20
	2.4. EBAC matrices	29
	2.5. Types of ligands	30
	2.5.1. Ion-exchanger	30
	2.5.2. Affinity ligand	32
3	HEAT-TREATMENT OF UNCLARIFIED FEEDSTOCK IMPROVED THE RECOVERY EFFICIENCY OF RECOMBINANT HEPATITIS B CORE ANTIGEN.	35
	3.1. INTRODUCTION	35
	3.2. MATERIALS AND METHODS	37
	3.2.1. Cultivation of <i>E. coli</i> cells	37
	3.2.2. Conventional method of HBcAg purification and heat treatment	37
	3.2.3. Preparation of unclarified feedstock	38
	3.2.4. Optimisation of thermal treatment of unclarified feedstock direct capture of HBcAg	39
	3.2.5. Batch adsorption	39
	3.2.6. The Bradford assay	40
	3.2.7. SDS-Polyacrylamide gel electrophoresis (SDS-PAGE)	40
	3.2.8. Quantitation of HBcAg	42
	3.2.9. Enzyme-linked immunosorbent assay (ELISA)	42



3.3.	RESULTS AND DISCUSSION	43
3.3.1.	Application of heat treatment in the conventional method of HBcAg purification	43
3.3.2.	Direct capture of HBcAg from heat-treated unclarified feedstock	44
3.3.3.	Effect of heat treatment on the antigenicity of HBcAg	49
3.4.	CONCLUSION	50
4	DIRECT PURIFICATION OF RECOMBINANT HEPATITIS B CORE ANTIGEN FROM TWO DIFFERENT PRE-CONDITIONED UNCLARIFIED FEEDSTOCKS	51
4.1.	INTRODUCTION	51
4.2.	MATERIALS AND METHODS	53
4.2.1.	Materials	53
4.2.2.	Feedstocks preparation	54
4.2.3.	Method scouting and optimisation	54
4.2.4.	Equilibrium adsorption isotherm	55
4.2.5.	Operation of the EBAC column	56
4.2.6.	Quantitation of total protein and HBcAg	58
4.2.7.	Qualitative analysis of HBcAg	58
4.2.8.	Calculations	59
4.3.	RESULTS AND DISCUSSION	60
4.3.1.	Optimisation of binding pH	60
4.3.2.	Optimisation of elution condition	62
4.3.3.	Equilibrium adsorption isotherm	64
4.3.4.	Operation of the EBAC column	65
4.3.5.	Operation of EBAC	70
4.4.	CONCLUSION	78
5	EFFECT OF DIFFERENT OPERATING MODES AND BIOMASS CONCENTRATIONS ON THE RECOVERY OF RECOMBINANT HEPATITIS B CORE ANTIGEN	79
5.1.	INTRODUCTION	79
5.2.	MATERIALS AND METHODS	81
5.2.1.	Materials	81
5.2.2.	Preparation of heat-treated unclarified feedstock	81
5.2.3.	Viscosity measurement	81
5.2.4.	Operation of EBAC	82
5.2.5.	Total protein and HBcAg quantitation	84
5.3.	RESULTS AND DISCUSSION	85
5.3.1.	Liquid dispersion behaviour in the expanded bed	85
5.3.2.	Effect of different bed heights and biomass concentration on the performance EBAC operation	86
5.4.	CONCLUSION	97

6	IMMOBILISATION OF FUSION M13 BACTERIOPHAGE ON EBAC ADSORBENT IN AFFINITY CAPTURING OF HEPATITIS B CORE ANTIGEN	99
6.1.	INTRODUCTION	99
6.2.	MATERIALS AND METHODS	101
6.2.1.	Materials	101
6.2.2.	Large scale preparation and purification of M13 bacteriophage	101
6.2.3.	Preparation of unclarified <i>E. coli</i> feedstock	103
6.2.4.	Selection of a suitable activation method	103
6.2.5.	Immobilisation of M13 phage	106
6.2.6.	Batch adsorption	108
6.2.7.	Quantitation of total protein and HBcAg	109
6.2.8.	Qualitative evaluations on M13-immobilised adsorbents	109
6.3.	RESULTS AND DISCUSSION	111
6.3.1.	Selection of a suitable activation method	111
6.3.2.	Effect of different ECH concentrations	114
6.3.3.	Effect of different NaOH concentrations	116
6.3.4.	Effect of reaction time and temperature	117
6.3.5.	Optimisation of M13 immobilisation	119
6.3.6.	Effect of ligand density	123
6.3.7.	Blocking of residual groups	124
6.3.8.	Stability and reusability of M13 immobilised adsorbents	126
6.4.	CONCLUSION	128
7	DIRECT PURIFICATION OF RECOMBINANT HEPATITIS B CORE ANTIGEN USING M13 PHAGE-IMMOBILISED ADSORBENTS	129
7.1.	INTRODUCTION	129
7.2.	MATERIALS AND METHODS	130
7.2.1.	Materials	130
7.2.2.	Preparation of M13 phage	130
7.2.3.	Epoxy activation and immobilization of M13 phage onto Streamline Base Matrix	131
7.2.4.	Preparation of unclarified <i>E. coli</i> feedstock	131
7.2.5.	Method scouting and optimization	131
7.2.6.	EBAC operation	133
7.2.7.	Quantitation of total protein and HBcAg	135
7.2.8.	Qualitative analysis of purified HBcAg	135
7.2.9.	Calculations	136
7.3.	RESULTS AND DISCUSSION	136
7.3.1.	Method scouting and optimization	136
7.3.2.	HBcAg uptake rate	142
7.3.3.	Equilibrium adsorption isotherm	144
7.3.4.	Operation of EBAC	145
7.4.	CONCLUSION	157

8	OVERALL SUMMARY AND RECOMMENDATION FOR FUTURE RESEARCH	159
---	---	-----

	REFERENCES	165
	BIODATA OF THE STUDENT	188
	LIST OF PUBLICATIONS	189



LIST OF TABLES

Table		Page
2.1	Summary of products purified from various sources using EBAC.	22
3.1	Comparison of the purity, purification factor and yield of HBcAg of the thermal and non-thermal treated unclarified feedstock.	45
3.2	Comparison between the yield and the purity obtained after batch adsorption with unclarified feedstocks treated at different temperatures.	47
3.3	Comparison between the yield and the purity obtained after batch adsorption with crude feedstocks treated at 60°C of different treatment durations.	47
4.1	Effect of pH on the binding efficiency of HBcAg from unclarified feedstock.	61
4.2	Effect of NaCl concentrations on the elution of HBcAg.	62
4.3	Results of the linear regression of Richardson-Zaki plot.	67
4.4	Purification of HBcAg from heat-treated and non heat-treated unclarified feedstocks.	73
5.1	Velocities employed to maintain constant bed expansion degree of 2.	84
5.2	Liquid dispersion of the expanded bed operating at constant velocity of 127.9 cm/h with degree expansion of 2.	87
5.3	The adsorption performance of Streamline DEAE in different mode of EBAC operations.	89
5.4	Viscosity of equilibration buffer, non-heat treated and heat-treated biomass containing feedstock. The data represent the mean \pm standard deviation of triplicate measurements.	94
6.1	Effect of different activation methods on the immobilization of M13 and binding of HBcAg from unclarified <i>E. coli</i> feedstock with Streamline base matrix.	112
6.2	Effect of pH on the immobilization of M13 onto epoxidated Streamline base matrix and efficiency of HBcAg binding from unclarified <i>E. coli</i> feedstock onto M13-immobilised adsorbents.	120

6.3	Effect of different ligand densities on HBcAg binding.	124
6.4	Effect of different ethanolamine concentrations on residual epoxy groups.	125
6.5	Stability of M13-immobilised adsorbents during storage.	127
6.6	Reusability of M13-immobilised adsorbents.	127
7.1	Effect of different pH on the binding of HBcAg onto M13 phage-immobilised adsorbents.	138
7.2	Elution efficiency of different chaotropic salts as eluting agents.	141
7.3	Effect of different urea concentrations on elution.	142
7.4	Effect of different EBAC operations on direct capture of HBcAg from crude feedstock.	150
7.5	Amount of ligand leakage at every purification step in typical and modified EBAC operations.	155
7.6	Light scattering analysis on typical and modified EBAC purified core particles.	156



LIST OF FIGURES

Figure		Page
2.1	A schematic representative of HBV virion structure.	6
2.2	A schematic diagram of an EBAC column.	21
3.1	SDS-PAGE gel of the heat-treated (60°C, 30 min) and the non-heat-treated feedstock in the conventional method of HBcAg purification.	45
3.2	SDS-PAGE gel of the batch binding assay by using anion-exchange adsorbents and crude feedstock without heat treatment.	48
3.3	SDS-PAGE gel of the batch binding assay by using anion-exchange adsorbents and heat-treated feedstock at 60°C for 45 min.	48
3.4	The antigenicity level exhibited by the core particles after thermal-treated at different temperatures.	49
4.1	Elution of HBcAg using different NaCl concentrations.	63
4.2	Equilibrium adsorption isotherm of HBcAg from non-heat-treated and heat-treated unclarified feedstocks on Streamline DEAE.	65
4.3	Bed expansion characteristics of Streamline DEAE with different liquid phases.	66
4.4	Breakthrough curve of the HBcAg from unclarified feedstock containing 5% of biomass onto Streamline DEAE in EBAC.	69
4.5	EBAC recovery of HBcAg.	71
4.6	SDS-PAGE gels of the pooled fractions collected throughout the EBAC purification of HBcAg from two different pre-conditioned unclarified feedstocks.	74
4.7	Breakthrough curves of HBcAg and total protein from two different pre-conditioned feedstocks in EBAC.	75
4.8	Light scattering analysis of HBcAg and other contaminating proteins present in two different pre-conditioned unclarified feedstocks.	77
4.9	ELISA analysis of purified HBcAg from heat-treated unclarified feedstock.	78

5.1	EBAC operation at constant velocity.	90
5.2	EBAC operation at constant bed expansion.	91
5.3	Breakthrough curves of the HBcAg and total protein onto Streamline DEAE in EBAC operated at constant bed expansion.	96
5.4	Breakthrough curves of the HBcAg and total protein onto Streamline DEAE in EBAC operated at constant superficial velocity.	96
6.1	Optimisation of epoxy activation on Streamline Base Matrix using different concentrations of epichlorohydrin.	115
6.2	Effect of different concentrations of NaOH on epoxy activation of Streamline Base Matrix.	116
6.3	Effect of different temperatures on epoxy activation.	118
6.4	Epoxy activation of Streamline Base Matrix at 25°C.	118
6.5	SEM micrograph of epoxidated Streamline Base Matrix.	119
6.6	Macroporous surface of Streamline Base Matrix.	121
6.7	Immunofluorescence staining with non-epoxidated adsorbents.	122
6.8	Immunofluorescence staining of M13 phage-immobilised adsorbents.	122
6.9	Immunofluorescence staining of epoxidated Streamline Base Matrix.	126
7.1	Optimisation of HBcAg elution using different eluting agents.	140
7.2	Uptake rate of HBcAg using M13 phage-immobilised adsorbents.	143
7.3	Equilibrium adsorption isotherm of HBcAg on M13 phage-immobilised adsorbents.	145
7.4	Dynamic binding capacity of the expanded bed using M13 phage-immobilised adsorbents against HBcAg from unclarified feedstock containing 5% biomass.	146
7.5	The breakthrough curves of total protein and HBcAg from unclarified feedstock containing 5% of biomass in typical EBAC operation.	148
7.6	Recovery of HBcAg using typical single pass EBAC.	150



7.7	Recovery of HBcAg from modified EBAC operation.	152
7.8	Elution profile from typical single pass EBAC operation.	154
7.9	Elution profile from modified EBAC operation.	154
7.10	ELISA analysis of HBcAg purified from different modes of EBAC operations.	157
8.1	Process scheme for the recovery of HBcAg from <i>E. coli</i> unclarified feedstock via EBAC purification using Streamline DEAE.	159
8.2	Process scheme for the recovery of HBcAg from <i>E. coli</i> unclarified feedstock via EBAC purification using M13 phage-immobilised adsorbents.	160



LIST OF ABBREVIATIONS

aa	amino acid
BSA	bovine serum albumin
CIP	clean-in-place
cm	centimetre
cm/h	centimetre per hour
cP	centipoise
C-terminus	carboxy-terminus
DNA	deoxy-ribonucleic acid
DNAse	deoxyribonuclease
EBAC	expanded bed adsorption chromatography
ELISA	enzyme-linked immunosorbent assay
FITC	fluorescein isothiocyanate
g	gram
g/ml	gram per milliliter
h	hour
HBcAg	hepatitis B core antigen
HBV	hepatitis B virus
HCl	hydrochloric acid
HPLC	high performance liquid chromatography
IPTG	isopropyl- β -D-thiogalactopyranoside
KCl	potassium chloride
KH ₂ PO ₄	potassium di-hydrogen phosphate
KSCN	potassium thiocyanate
mg	milligram

min	minute
mL	milliliter
mM	millimolar
Na ₂ HPO ₄	di-sodium hydrogen phosphate
NaCl	sodium chloride
NaI	sodium iodide
NaOH	sodium hydroxide
Ni ²⁺	nickel ion
N-terminus	amino-terminus
OD	optical density
PAGE	polyacrylamide gel electrophoresis
PEG	polyethylene glycol
pfu	plaque forming unit
pH	<i>Puissance hydrogene</i>
p-npp	ρ-nitro-phenyl phosphate
RNA	ribonucleic acid
rpm	revolution per minute
s	second
SDS	sodium dodecyl sulphate
SEM	scanning electron microscope
<i>T</i>	triangulation number
TEMED	tetramethyl ethylenediamine
x-gal	5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside

CHAPTER 1

INTRODUCTION

Capsid of the hepatitis B virus (HBV), which is composed of multiple subunits of core antigens (HBcAg) has become one of the most studied viral-like-particle (VLP) that possess similar morphology and self-assemble ability to the native capsids without involvement of the viral genome (Pumpens and Grens, 2001). HBcAg is a strong immunogen (Milich *et al.*, 1997; Milich and McLachlan, 1986), and its capability in generating both humoral and cell-mediated immunity has made hepatitis B virus (HBV) core particle an excellent tool in delivering antigens for vaccination purpose (Nassal *et al.*, 2008; Chen *et al.*, 2004). This has also benefited in development of various immunoassays that are promising to use as diagnostic reagents (Li *et al.*, 2007; Touze *et al.*, 1999). Therefore, the quality of HBcAg should not be compromised as presence of impurities could lead to various problems such as inaccuracies in detection of specific antibodies in diagnostic kit (Li *et al.*, 2007). Therefore, due to the wide spread application of HBcAg, it is essential to produce this protein at low cost in both upstream and downstream processes (Mechtcheriakova *et al.*, 2006). However, apart from being too tedious to perform as it requires rounds of clarification steps, the available current method of HBcAg purification such as size-exclusion chromatography (Tang *et al.*, 2007; Broos *et al.*, 2007) and sucrose gradient ultracentrifugation (Huang *et al.*, 2006; Tan *et al.*, 2003), also involves higher production cost, longer processing time and low in product yield (Rolland *et al.*, 2001).



Therefore, alternative method such as expanded bed adsorption chromatography (EBAC) can be sought to improve the downstream process of HBcAg. EBAC has long been used in direct recovery of biomolecules from various sources such as *E. coli* and yeast cells homogenate (Chow *et al.*, 2007; Tan *et al.*, 2005; Chase and Draeger, 1992) as well as more complex feedstocks such as plant extracts (Valdès *et al.*, 2003) and chicken egg white (Chang and Chang, 2006) without the need of pre-clarification steps. Unlike the conventional packed-bed chromatography that requires particulate-free feed stream, this method of purification forms a stable expanded bed that allows unhindered passage for cell or cells debris to flow through the column without the risk of clogging up the bed (Chase, 1994). Furthermore, EBAC combines three important steps in purification; clarification, capture and concentration steps into one single operation, hence, resulted in shorter processing time (Anspach *et al.*, 1999; Chase and Draeger, 1992).

At present, the commonly-used EBAC matrices are ion-exchangers (Clemmitt and Chase, 2002) and metal-chelating (Tan *et al.*, 2005) adsorbents. However, EBAC operation using ion-exchangers is greatly influenced by electrostatic interactions between cell or cell debris in the feedstock and the adsorbent (Lin *et al.*, 2003) as well as the viscosity and ionic strength of the applied feedstock (Chang and Chase, 1996a; Anspach *et al.*, 1999). Therefore, affinity ligands are preferred as they are less likely to be affected by the aforementioned factors (Chase and Draeger, 1992). Unlike packed bed chromatography resin, affinity chromatographies in EBAC are usually carried out using commercially-available affinity ligands such as metal ions (Tan *et al.*, 2005) and dye-ligand (Reichert *et al.*, 2001). In view of these, many new solid phases, which can be chemically-linked with various synthetic and biological