



UNIVERSITI PUTRA MALAYSIA

EFFECTS OF FEEDING METABOLITES FROM *LACTOBACILLUS PLANTARUM* STRAINS ON LIPID METABOLISM, GUT MORPHOLOGY AND GROWTH PERFORMANCE OF BROILER CHICKENS

NGUYEN TIEN THANH

FP 2008 6



**EFFECTS OF FEEDING METABOLITES FROM *LACTOBACILLUS*
PLANTARUM STRAINS ON LIPID METABOLISM, GUT MORPHOLOGY
AND GROWTH PERFORMANCE OF BROILER CHICKENS**

By

NGUYEN TIEN THANH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

July 2008



DEDICATION

Dedicated to my Dearest Mother, my beloved family and my brother and sisters



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree of Doctor of Philosophy

EFFECTS OF FEEDING METABOLITES FROM *LACTOBACILLUS PLANTARUM* STRAINS ON LIPID METABOLISM, GUT MORPHOLOGY AND GROWTH PERFORMANCE OF BROILER CHICKENS

By

NGUYEN TIEN THANH

July 2008

Chairman: Associate Professor Loh Teck Chwen, PhD

Faculty: Agriculture

Four experiments were conducted to study the effects of feeding metabolite combinations produced from six strains of *L. plantarum* on the performance of broiler chickens. The inhibitory activity of different combinations of metabolites produced by locally isolated *L. plantarum* against various pathogens was studied in the first experiment. Sixty-three combinations of metabolites obtained from 6 strains of *L. plantarum*: UL4, TL1, RS5, RI11, RG14 and RG11 were equally and homogenously mixed. The inhibitory activity was then determined against 5 selected indicators, which are *E. coli*, *L. monocytogenes*, *S. typhimurium*, Vancomycin resistant enterococci (VRE) and *Pediococcus acidilactici*. The inhibitory activity was measured based on the diameter of inhibitory zone. Four combinations with the highest inhibitory scores were identified. The combination of four strains RS5, RI11, RG14 and RG11 has given the highest score, followed by the combinations of TL1, RG14 and RG11 strains, combinations of TL1, RI11 and RG11 strains, and combinations of TL1, RS5, RI11 and RG14 strains. These results indicate that

different combinations of metabolites had different antibacterial activity, which could be used in food and feed industries. Combinations of different metabolites further enhance the antimicrobial activity.

Four combinations of metabolites with the highest inhibitory activity were used in the following experiment to study the performance of broiler chickens. A total of 432 day-old male Ross broiler chicks were raised to 42 days of age in deep litter system. Each pen consisted of 12 chicks and was randomly allocated to the open house with wood shavings litter. The birds were vaccinated (IB-ND live vaccine, Fort Dodge, USA) against infectious bronchitis (IB) and Newcastle disease (ND). The birds were also vaccinated with IBD vaccine (MyVac UPM93, Malaysia) against infectious bursal disease (IBD) on day 14. Wing band was applied to all of the birds for identification. Water and feed were provided *ad libitum*. The feed intake and body weight were recorded weekly. The starter and finisher diets were offered to the birds from 0 - 21 and 22 - 42 days of age, respectively. The dietary treatments consisted of: (i) corn-soybean based diet without antibiotic (-ve control) diet; (ii) basal diet with neomycin and oxytetracyclin (+ve control); (iii) basal diet supplemented with 0.3% metabolite combinations of *L. plantarum* RS5, RI11, RG14 and RG11 (Com3456); (iv) basal diet supplemented with 0.3% metabolite combinations of *L. plantarum* TL1, RG14 and RG11 (Com256); (v) basal diet supplemented with 0.3% metabolite combinations of *L. plantarum* TL1, RI11 and RG11 (Com246); (vi) basal diet supplemented with 0.3% metabolite combinations of *L. plantarum* TL1, RS5, RG14 and RG11 (Com2456). Greater final body weight (BW), weight gain (WG), average daily gain (ADG), volatile fatty acids (VFA) and lower feed conversion ratio (FCR) were significantly ($p < 0.05$) found in the birds fed with the 4 metabolite combinations.

Supplementation of metabolites combination also increased faecal lactic acid bacteria (LAB) population, villi height of small intestine and lowered plasma and meat cholesterol and faecal *Enterobacteriaceae* (ENT) population.

In the third experiment, the effect of feeding different dosages of Com3456 obtained from the second experiment on the performance of broiler chickens was studied. A total of 504 day-old male Ross broiler chicks were grouped into 7 treatments and offered with different diets: (i) standard corn-soybean based diet (negative control); (ii) standard corn-soybean based diet + neomycin and oxytetracycline (positive control); (iii) standard corn-soybean based diet + 0.1% metabolite combination of *L. plantarum* RS5, RI11, RG14 and RG11 strains (Com3456); (iv) standard corn-soybean based diet + 0.2% of Com3456; (v) standard corn-soybean based diet + 0.3% of Com3456 (vi) standard corn-soybean based diet + 0.4% of Com3456 and (vii) standard corn-soybean based diet + 0.5% of Com3456. Supplementation of Com3456 with different dosages improved growth performance, reduced ENT and increased LAB count, lowered plasma and meat cholesterol, and increased villi height of small intestine and volatile fatty acids (VFA). Nevertheless, abdominal fat deposition was not affected by Com3456 metabolites. Only slight increase of antibody titers against Newcastle disease, infectious bronchitis and infectious bursal disease was found in those birds supplemented with the metabolites. Treatment with 0.2% Com3456 had the best results with minimal cost, especially in terms of growth performance, FCR and plasma and meat cholesterol reduction among other dosages. These results indicate that 0.2% is optimum dosages to be included in the diets of broiler chickens in order to replace antimicrobial growth promoters (AGP).

In the last experiment, the very low-density lipoprotein (VLDL) lipid profiles, intestinal LAB count and bile salts deconjugation of LAB were studied. The results showed that metabolite combinations supplemented in broilers feed reduced free cholesterol and cholesterol esters in VLDL particles and increased LAB count in digesta of small intestine. The results also suggest that LAB are able to deconjugate bile salts and lead to the increase of cholesterol utilization for synthesizing new conjugated bile salts. These properties of metabolite combinations in the diets of broiler chickens contribute to the reduction of cholesterol in plasma and meat.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KESAN PEMBERIAN METABOLIT DARIPADA STRAIN
LACTOBACILLUS PLANTARUM TERHADAP METABOLISME LIPID,
MORFOLOGI SALUR GASTRO-USUS DAN PRESTASI
PERTUNBAHAN AYAM PEDAGING**

Oleh

NGUYEN TIEN THANH

July 2008

Pengerusi: Profesor Madya Loh Teck Chwen, PhD

Fakulti: Pertanian

Empat eksperimen telah dijalankan untuk mengkaji kesan kombinasi metabolit yang dihasilkan daripada 6 strain *L. plantarum* ke atas prestasi ayam pedaging. Aktiviti perencat daripada kombinasi metabolit berlainan yang dihasilkan melalui pemencilan *L. plantarum* terhadap pelbagai patogen telah dikaji dalam eksperimen pertama. Enam puluh tiga kombinasi metabolit diperolehi daripada enam strain *L. plantarum*: UL4, TL1, RS5, RI11, RG14 dan RG11 yang dicampur secara sekata dan homogenus. Aktiviti perencat telah dijalankan ke atas 5 petunjuk terpilih, iaitu *E. coli*, *L. monocytogenes*, *S. typhimurium*, Vancomycin resistan enterococci (VRE) dan *Pediococcus acidilactici*. Aktiviti perencat ditindakan berdasarkan kepada diameter zon perencat. Empat kombinasi metabolit dengan skor perencat tertinggi telah dikenal pasti. Kombinasi bagi empat strain RS5, RI11, RG14 dan RG11 menunjukkan skor tertinggi, diikuti dengan kombinasi strain TL1, RG14 dan RG11, kombinasi strain TL1, RG11 dan RG11, dan kombinasi strain TL1, RS5, RI11 dan RG14. Keputusan ini menunjukkan perbezaan kombinasi metabolit yang berbeza

menyebabkan berbezaan aktiviti antibakteria, maklumat maklumat ini boleh diguna pakai dalam industri makanan manusia dan makanan haiwan. Pada umumnya kombinasi metabolit ini akan mempertingkatkan aktiviti antimikrob.

Empat kombinasi metabolit dengan aktiviti perencat tertinggi telah digunakan dalam eksperimen kedua untuk mengkaji prestasi ayam pedaging. Sejumlah 432 ekor ayam pedaging Ross jantan yang diperoleh dari sebuah syarikat tempatan telah ditenak dari umur satu hari hingga 42 hari menggunakan sistem sarap. Setiap reban terdiri daripada 12 anak ayam yang ditempatkan secara rawak dalam sistem rumah terbuka dengan menggunakan sarap tebal berasaskan habuk kayu. Ayam disuntikkan dengan vaksin (IB-ND vaksin hidup, Fort Dodge, USA) IBD dan ND. Ayam pedaging ini juga diberikan vaksin IBD (MyVac UPM93, Malaysia) pada hari ke-14. 'Wing banding' juga dikenakan kepada semua ayam bagi memudahkan pengecaman. Air dan makanan dibekalkan secara *ad libitum*. Makanan dan berat badan ditimbang setiap minggu. Makanan permulaan dan penghabisan masing-masing diberikan kepada ayam dari 0-21 dan 22-42 hari. Rawatan terdiri daripada: (i) diet berasaskan jagung-kacang soya tanpa antibiotik (kawalan -ve); (ii) diet ditambahkan antibiotik (kawalan +ve); (iii) diet asas ditambahkan dengan 0.3% kombinasi metabolit *L. plantarum* RS5, RI11, RG14 dan RG11 (Com3456); (iv) diet asas ditambahkan dengan 0.3% kombinasi metabolit *L. plantarum* TL1, RG14 dan RG11 (Com 256); (v) diet asas ditambahkan dengan 0.3% kombinasi metabolit *L. plantarum* TL1, RI11 dan RG11 (Com 246) dan (vi) diet asas ditambahkan dengan 0.3% kombinasi metabolit *L. plantarum* TL1, RI11, RG14 dan RG11 (Com 2456). Kelebihan berat badan akhir, pertambahan berat badan, purata pertambahan berat harian, VFA dan kadar pertukaran makanan yang lebih rendah diperoleh daripada kumpulan ayam

yang diberikan metabolit. Pemberian kombinasi metabolit juga meningkatkan populasi bakteri asid laktik tinja, ketinggian vilus usus kecil di samping merendahkan kolesterol plasma dan populasi ENT tinja.

Dalam eksperimen ketiga, kesan dos Com3456 yang berlainan yang diperolehi daripada eksperimen kedua telah dikaji ke atas prestasi ayam pedaging. Sejumlah 504 ekor ayam pedaging Ross jantan dibahagikan kepada 7 kumpulan melalui dan diberikan rawatan pemberian diet yang berbeza seperti berikan: (i) diet kawalan berasaskan jagung-kacang soya (kawalan negatif); (ii) diet kawalan berasaskan jagung-kacang soya + neomycin dan oxytetracyclin (kawalan positif); (iii) diet kawalan berasaskan jagung-kacang soya + 0.1% kombinasi metabolit *L. plantarum* strain RS5, RI11, RG14 dan RG11 (Com3456); (iv) diet kawalan berasaskan jagung-kacang soya + 0.2% kombinasi metabolit Com3456; (v) diet kawalan berasaskan jagung-kacang soya + 0.3% kombinasi metabolit Com3456; (vi) diet kawalan berasaskan jagung-kacang soya + 0.4% kombinasi metabolit Com3456 dan (vii) diet kawalan berasaskan jagung-kacang soya + 0.5% kombinasi metabolit Com3456. Pemberian Com3456 yang berlainan dos meningkatkan prestasi pertumbuhan, merendahkan ENT dan meningkatkan bilangan LAB, merendahkan kolesterol plasma dan daging, serta meningkatkan ketinggian vilus usus kecil dan VFA. Walau bagaimana pun, pertambahan lemak abdomine tidak dipengaruhi oleh metabolit. Hanya sedikit peningkatan pada antibodi titer "Newcastle disease", "infection bronchitis" dan "infectious bursal disease" dijumpai pada ayam yang diberikan metabolit. Rawatan dengan 0.2% Com3456 memberi keputusan yang terbaik, terutamanya dari segi prestasi pertumbuhan, kadar pertukaran makanan dan penurunan kolesterol plasma dan daging dibandingkan dengan dos lain. Keputusan

menunjukkan bahwa 0.2% merupakan optimum untuk menggantikan AGP di dalam diet ayam pedaging.

Dalam eksperimen terakhir, profil lemak VLDL, bilangan LAB dalam usus dan "bile salt deconjugation" oleh LAB telah dikaji. Keputusan menunjukkan kombinasi metabolit dalam makanan ayam pedaging dapat merendahkan "free cholesterol" dan "cholesterol ester" di dalam VLDL serta meningkatkan LAB dalam "digesta" usus kecil. Keputusan ini juga mencadangkan LAB berupaya untuk "menyahkonjugat bile salts" serta menggalakkan penggunaan kolesterol untuk mensitesiskan "bile salts" konjugat baru. Kandungan daripada kombinasi metabolit dalam diet ayam pedaging menyumbang kepada penurunan kolesterol dalam plasma dan daging.

ACKNOWLEDGEMENTS

I would like to express my profound gratitude to Associate Professor Dr. Loh Teck Chwen, chairman of the supervisory committee for granting Graduate Research Assistantship and his beneficial advice, guidance, scientific criticism and invaluable suggestion during the study.

I also wish to express my deep gratitude and appreciation to members of the supervisory committee: Professor Dr. Mohd Hair Bejo, Associate Professor Dr. Azhar Kasim and Associate Professor Dr. Foo Hooi Ling for their guidance, concern, patience, inspiring and constructive comments throughout the course of the study.

Deep appreciation is also extended to School of Graduate Studies, Universiti Putra Malaysia for granting Graduate Research Fellowship and providing excellent services and support. Special thanks to Prof. Raha, Head of Department, Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, Dr. Goh Yong Meng from Physiology Laboratory, Faculty of Veterinary Medicine for the supports. I wish to thank all lecturers and staffs from the Department of Animal Science, Faculty of Agriculture, UPM for their support. Thanks are also due to staffs of Fermentation Unit, Bacteriology Laboratory, Faculty of Biotechnology and Biomolecular Sciences, Microbiology and Pathology Laboratory 1, Faculty of Veterinary Medicine, Food Engineering Laboratory, Faculty of Food Science, UPM for the enthusiastic supports. Thanks to Hamidah Binti Ali Kamarulzaman for translating the abstract. Thanks to Dr. Le Ngoc Thach for the support.

Sincere thanks and appreciation is also extended to my institution, Faculty of Animal Science and Veterinary Medicine, Nong Lam University, Ho Chi Minh City for the supports. Finally, the deepest gratitude goes to my dearest Mom, who dedicated her whole life to her children and passed away in 1992, my beloved wife for taking care of our children during the study leave, my brother Nguyen Tien Dai and sisters for the support during my absence.



I certify that an Examination Committee met on 28 July, 2008 to conduct the final examination of Nguyen Tien Thanh on his PhD of Agriculture Science thesis entitled “Effects of Feeding Metabolites from *Lactobacillus Plantarum* Strains on Lipid Metabolism, Gut Morphology and Growth Performance of Broiler Chickens” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 a Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Halimatun Yaakub, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Abdul Razak Alimon, PhD

Professor
Faculty of Agriculture
Univerisiti Putra Malaysia
(Internal Examiner)

Awis Qurni Sazili, PhD

Faculty of Agriculture
Univerisiti Putra Malaysia
(Internal Examiner)

Wan Zahari Mohamed, PhD

Malaysia Agricultural Research Development Institute
(Independent Examiner)

HASANAH MOHD. GHAZALI, PhD

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: _____ 2008

This thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirements for the degree of PhD of Agriculture Science. The members of the Supervisory Committee were as follows:

Loh Teck Chwen, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Mohd Hair Bejo, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Azhar Bin Kasim, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

AINI IDERIS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11 September 2008

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

NGUYEN TIEN THANH

Date:

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xi
DECLARATION	xiv
LIST OF TABLES	xix
LIST OF FIGURES	xxii
LIST OF PLATES	xxiii
ABBREVIATIONS	xxiv
I INTRODUCTION	1
II LITERATURE REVIEW	4
2.1 Antimicrobial Growth Promoters in Animal Production	
2.1.1 Antibiotic Resistance of Bacteria	4
2.1.2 Mode of Action and Benefits of Antimicrobial Growth Promoters	5
2.2 Background of Lactic Acid Bacteria	6
2.2.1 Basic Characteristics	6
2.2.2 Classification	7
2.3 Lactic Acid Bacteria Fermentation	9
2.3.1 Fermentation	9
2.3.2 Pathways of Lactic Acid Bacteria Fermentation	10
2.3.3 Benefits of Basic Fermentative Metabolites	12
2.4 Lactic Acid Bacteria Roles in The Gut	20
2.4.1 Benefits of Lactic Acid Bacteria in Lowering Cholesterol	20
2.4.2 Lactic Acid Bacteria in Enhancing Immunity	23
2.4.3 Lactic Acid Bacteria in Reducing <i>Enterobacteriaceae</i> Population	25
2.5 Chicken Intestinal Morphology	26
2.5.1 General Overview	26
2.5.2 Crypts and Villi	28
2.5.3 Factors Affecting Villi Height	29
2.6 Bile Metabolism	30
2.6.1 Bile Synthesis, Storage and Secretion	30
2.6.2 Bile Acids Conjugation and Its Functions	32
2.6.3 Enterohepatic Circulation	34
2.6.4 Bacterial Alteration of Bile Acids	34
2.7 Lipid Metabolism	36
2.7.1 General Background	36
2.7.2 Intestinal Uptake of Lipids	36
2.7.3 Cholesterol and Its Metabolism	38
2.7.4 Lipoprotein Complexes	42



2.7.5 Very Low-density Lipoprotein Metabolism	43
2.7.6 Fat Tissue in Broiler Chickens	47
2.8 Microbial Community in Gastrointestinal Tract	48
2.8.1 General Information	48
2.8.2 pH Value in Chicken Gastrointestinal Tract	50
2.8.3 Microflora in Different Parts of Chicken Gastrointestinal Tract	52
2.9 Chicken Maternal Antibody and Antibody Response	54
III INHIBITORY ACTIVITY OF DIFFERENT METABOLITE COMBINATIONS FROM STRAINS OF <i>LACTOBACILLUS PLANTARUM</i> AGAINST PATHOGENS	55
3.1 Introduction	55
3.2 Materials and Methods	57
3.2.1 Treatments and Experimental Design	57
3.2.2 Strains, Media and Metabolite Preparation	57
3.2.3 Agar-well Diffusion Method	59
3.2.4 Large-scale Fermentation Profile	59
3.2.5 Assessment of Inhibitory Zone Diameter	62
3.2.6 Data Analysis	63
3.3 Results	63
3.3.1 Inhibitory Test Results of Metabolites against Bacterial Strains	63
3.3.2 Characteristics of Metabolites from <i>Lactobacillus plantarum</i> Strains	69
3.4 Discussion	74
3.4.1 Antimicrobial Activity of Metabolite Combinations	74
3.4.2 Characteristics of Metabolites from <i>Lactobacillus plantarum</i> Strains	77
3.5 Conclusions	78
IV EFFECTS OF FEEDING DIFFERENT METABOLITE COMBINATIONS PRODUCED BY <i>LACTOBACILLUS PLANTARUM</i> ON THE PERFORMANCE OF BROILER CHICKENS	79
4.1 Introduction	79
4.2 Materials and Methods	80
4.2.1 Broiler chickens and Experimental Design	80
4.2.2 Data and Sample Collection	81
4.2.3 Faecal Lactic Acid Bacteria and <i>Enterobacteriaceae</i> Count	83
4.2.4 Small Intestine Histology	84
4.2.5 Total Plasma Cholesterol Level	85
4.2.6 Meat Cholesterol Determination	85
4.2.7 Volatile Fatty Acid Determination	87
4.2.8 pH Determination	88
4.2.9 Serological Assay	88
4.2.10 Data Analysis	89
4.3 Results	89
4.3.1 Growth Performance	89

4.3.2 Faecal Lactic Acid Bacteria and <i>Enterobacteriaceae</i> Count	90
4.3.3 Small Intestinal Villi Height and Crypt Depth	92
4.3.4 Plasma and Meat Cholesterol	97
4.3.5 Relationship of Plasma and Meat Cholesterol and Faecal Microflora Count	98
4.3.6 Faecal Volatile Fatty Acids	101
4.3.7 Faecal pH	103
4.3.8 Abdominal Fat Pad Weight	103
4.3.9 Antibody Titers against IBD, ND and IB	105
4.4 Discussion	107
4.4.1 Growth Performance	107
4.4.2 Lactic Acid Bacteria and <i>Enterobacteriaceae</i> Population	108
4.4.3 Villi Height and Crypt Depth	109
4.4.4 Plasma and Meat Cholesterol	111
4.4.5 Faecal pH and Volatile Fatty Acids	112
4.4.6 Abdominal Fat Pad Weight	114
4.4.7 Antibody Titers against IBD, ND and IB	115
4.5 Conclusions	116
V EFFECTS OF FEEDING DIFFERENT DOSAGES OF COM3456 METABOLITE COMBINATION ON THE PERFORMANCE OF BROILER CHICKENS	117
5.1 Introduction	117
5.2 Materials and Methods	118
5.2.1 Animal and Experimental Design	118
5.2.2 Data and Sample Collection	120
5.2.3 Data Analysis	121
5.3 Results	121
5.3.1 Growth Performance	121
5.3.2 Faecal Lactic Acid Bacteria and <i>Enterobacteriaceae</i> Count	122
5.3.3 Intestinal Villi Height and Crypt Depth	123
5.3.4 Plasma and Meat Cholesterol	128
5.3.5 Relationship of Plasma and Meat Cholesterol and Faecal Microflora Count	130
5.3.6 Faecal Volatile Fatty Acids	132
5.3.7 Faecal pH	134
5.3.8 Abdominal Fat Pad Weight	134
5.3.9 Antibody Titer against IBD, ND and IB	134
5.4 Discussion	138
5.4.1 Growth Performance	138
5.4.2 Lactic Acid Bacteria and <i>Enterobacteriaceae</i> Count	139
5.4.3 Villi Height and Crypt Depth	140
5.4.4 Plasma and Meat Cholesterol	142
5.4.5 Faecal pH and Volatile Fatty Acids	143
5.4.6 Abdominal Fat Pad Weight	144
5.4.7 Antibody Titers against IBD, ND and IB	145
5.5 Conclusions	145

VI	EFFECTS OF FEEDING METABOLITE COMBINATIONS ON THE PROFILES OF VERY LOW-DENSITY LIPOPROTEIN LIPIDS IN PLASMA AND BILE SALT DECONJUGATION ABILITY OF LACTIC ACID BACTERIA IN THE SMALL INTESTINE OF BROILER CHICKENS	146
	6.1 Introduction	146
	6.2 Materials and Methods	147
	6.2.1 Sample Collection	147
	6.2.2 Very Low-density Lipoprotein Preparation	148
	6.2.3 Protein Determination: Peterson's Modified Micro Lowry Method	149
	6.2.4 Triacylglycerol Assay	149
	6.2.5 Total Cholesterol Assay	150
	6.2.6 Free Cholesterol Assay	151
	6.2.7 Phospholipids Assay	152
	6.2.8 Calculation of the Ratios of Lipids in Very Low-density Lipoprotein	152
	6.2.9 Lactic Acid Bacteria Population in Small Intestine Digesta	153
	6.2.10 Deconjugation of Bile Salts by Lactic Acid Bacteria	153
	6.2.11 Data Analysis	154
	6.3 Results	154
	6.3.1 Very Low-density Lipoprotein Profile	154
	6.3.2 Relationship of Very Low-density Lipoprotein Lipids, Cholesterol and Abdominal Fat Deposition	158
	6.3.3 Lactic Acid Bacteria Count	158
	6.3.4 Deconjugated Bile Acid Released from Deconjugation	161
	6.4 Discussion	163
	6.4.1 Very Low-density Lipoprotein Profile	163
	6.4.2. Lactic Acid Bacteria Count and Bile Salts Deconjugation	165
	6.5 Conclusions	166
VII	GENERAL DISCUSSION AND CONCLUSIONS	167
	REFERENCES	179
	APPENDICES	203
	BIODATA OF STUDENT	212
	LIST OF PUBLICATIONS	213

LIST OF TABLES

Table	Page
2.1 Major lactic acid bacteria in fermented products	9
2.2 Characteristics and functions of the major lipoproteins in chicken	44
2.3 pH of the digestive tract contents of avian species	51
3.1 Various metabolites and their combinations produced from <i>Lactobacillus plantarum</i>	58
3.2a Inhibitory score rank of 63 combinations of metabolites produced by 6 strains of <i>Lactobacillus plantarum</i> against pathogens	66
3.2b Inhibitory score rank of 63 combinations of metabolites produced by 6 strains of <i>Lactobacillus plantarum</i> against pathogens (cont.)	67
3.2c Inhibitory score rank of 63 combinations of metabolites produced by 6 strains of <i>Lactobacillus plantarum</i> against pathogens (cont.)	68
3.3 Average inhibitory zone of 63 combinations of metabolites produced by 6 strains of <i>Lactobacillus plantarum</i> against different pathogens	69
3.4 Average optical density of different <i>Lactobacillus plantarum</i> strains culture during 24 hours of fermentation	71
3.5 Average pH of different <i>Lactobacillus plantarum</i> strains culture during 24 hours of fermentation	72
4.1 Percentage composition of starter diet	82
4.2 Percentage composition of finisher diet	83
4.3 Faecal lactic acid bacteria and <i>Enterobacteriaceae</i> count at weeks 3 and 6 of treatments supplemented with different metabolite combinations	91
4.4 Villi height and crypt depth in small intestine at weeks 3 and 6 of treatments supplemented with different metabolite combinations	95
4.5 The correlation coefficient and p-value between body weight and villi height at weeks 3 and 6	97
4.6 Plasma and meat cholesterol at weeks 3 and 6 of treatments supplemented with different metabolite combinations	99

4.7	Pearson correlation coefficients (r) and p-value of cholesterol, lactic acid bacteria and <i>Enterobacteriaceae</i> count	100
4.8	Faecal volatile fatty acid at weeks 3 and 6 of treatments supplemented with different metabolite combinations	102
4.9	Faecal pH at weeks 3 and 6 of treatments supplemented with different metabolite combinations	103
4.10	Abdominal fat pad weight and relative percentage compared to body weight at weeks 3 and 6 of treatments supplemented with different metabolite combinations	104
4.11	Antibody titers of IBD, ND and IB at weeks 4 and 5 treatments supplemented with different metabolite combinations	106
5.1	Percentage composition of starter diet	119
5.2	Percentage composition of finisher diet	120
5.3	Faecal lactic acid bacteria and <i>Enterobacteriaceae</i> count at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	124
5.4	Villi height and crypt depth in small intestine at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	127
5.5	Correlation coefficient between body weight and villi height of duodenum, jejunum and ileum	128
5.6	Plasma and meat cholesterol at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	129
5.7	Pearson correlation coefficients (r) and p-value of cholesterol, lactic acid bacteria and <i>Enterobacteriaceae</i> count	131
5.8	Faecal volatile fatty acid at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	133
5.9	Faecal pH at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	135
5.10	Abdominal fat weight and percentage compared to body weight at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	135

5.11	Antibody titers of IBD, ND and IB at weeks 3 and 6 of treatments supplemented with different dosages of metabolites from Com3456	137
6.1a	Very low-density lipoprotein lipid profile of treatments supplemented with different metabolite combinations	156
6.1b	Very low-density lipoprotein lipid profile of treatments supplemented with different dosages of metabolites from Com3456	157
6.2	Pearson correlation coefficients (r) and significance value between Very low-density lipoprotein lipids, cholesterol and abdominal fat deposition	159
6.3	Lactic acid bacteria count in small intestine digesta of treatments supplemented with different dosages of metabolites from Com3456	160
6.4	Deconjugated bile acid released by bile salts deconjugation by lactic acid bacteria in treatments supplemented with different dosages of metabolites from Com3456	162

LIST OF FIGURES

Figure		Page
2.1	Genus of lactic acid bacteria	8
2.2	Fermentative pathways	11
2.3	Mechanism of organic acid to pass through the cell membrane	15
2.4	The effect of antimicrobial peptides with different levels of their concentration	19
2.5	The gastrointestinal tract of chickens	26
2.6	The biosynthetic pathway of bile acids	32
2.7	Circulation of bile acids	33
2.8	Lipid metabolism in birds	37
2.9	Chemical structure of cholesterol	39
2.10	The biosynthesis pathway of cholesterol, triacylglycerol and phospholipid	41
2.11	The transport of triacylglycerol and cholesterol by lipoprotein in chickens	43
3.1	Bacteriocin activity of metabolites from <i>Lactobacillus plantarum</i> strains culture	72
3.2	The content of acetic acid in metabolites of the 6 strains	73
4.1	Growth performance of broiler chickens	90
5.1	Growth performance of broiler chickens	122
6.1	The principle of the reaction in triacylglycerol assay	150
6.2	The principle of reaction in total cholesterol assay	151
6.3	The principle of reaction in free cholesterol assay	151
6.4	The principle of reaction in phospholipid assay	152
7.1	Possible mechanism of cholesterol reduction in plasma and meat	173

LIST OF PLATES

Plate		Page
3.1	Inhibitory activity of metabolite combinations against pathogens	65
4.1	Measurement of ileal villi height and crypt depth	96

LIST OF ABBREVIATIONS

3 β HSD	3 β -hydroxysteroid dehydrogenase/isomerase
4MV	4-methyl-valeric acid
ADG	Average daily gain
ADP	Adenosine diphosphate
AGP	Antimicrobial growth promoters
AKR1C4	3 α -hydroxysteroid dehydrogenase
AKR1D1	Δ^{4-3} -oxosteroid-5 β -reductase
Apo	Apolipoprotein
ATP	Adenosine triphosphate
AU	Arbitrary unit
BSH	Bile salt hydrolase
BW	Body weight
CA	Cholic acid
CDCA	Chenodeoxycholic acid
CE	Cholesterol esters
CFS	Cell-free supernatant
CFU	Colony-forming unit
Com3456	Metabolite combination from <i>L. plantarum</i> RS5, RI11, RG14 and RG11
Com246	Metabolite combination from <i>L. plantarum</i> TL1, RI11 and RG11
Com2456	Metabolite combination from <i>L. plantarum</i> TL1, RI11, RG14 and RG11
Com256	Metabolite combination from <i>L. plantarum</i> TL1, RG14 and RG11
CYP27A1	Sterol 27-hydroxylase