



UNIVERSITI PUTRA MALAYSIA

**CHEMICAL COMPOSITION, ANTIOXIDATIVE AND CHOLESTEROL
LOWERING PROPERTIES OF SELECTED MALAYSIA SEaweeds**

PATRICIA MATANJUN

FSTM 2008 15

**CHEMICAL COMPOSITION, ANTIOXIDATIVE AND
CHOLESTEROL LOWERING
PROPERTIES OF SELECTED MALAYSIA SEAWEEDS**

By

PATRICIA MATANJUN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

December 2008



**To my husband Andre,
daughter Amanda, sons Eric and Isaac**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of
the requirement for the degree of Doctor of Philosophy

**CHEMICAL COMPOSITION, ANTIOXIDATIVE AND CHOLESTEROL
LOWERING PROPERTIES OF SELECTED MALAYSIA SEAWEEDS**

By

PATRICIA MATANJUN

DECEMBER 2008

Chairman : Professor Dr. Suhaila Mohamed, PhD

Faculty : Food Science and Technology

The objectives of this study were to evaluate *in vitro* antioxidant activity and total phenolic screenings of eight species of Malaysia seaweeds (*Kappaphycus alvarezii*, *Eucheuma denticulatum*, *Halymenia durvillaei*, *Caulerpa lentillifera*, *Caulerpa racemosa*, *Dicyota dichotoma*, *Sargassum polycystum* and *Padina* spp.), determine chemical composition of three selected edible seaweeds and investigate effects of these seaweeds on antioxidative, cholesterol-lowering, and their effects on biochemical, morphological and histological characteristics of selected tissues of rats fed on high-cholesterol/high-fat (HCF) diets. *In vitro* antioxidant activities of the eight species of seaweeds were evaluated using TEAC (trolox equivalent antioxidant capacity) and FRAP (ferric reducing antioxidant power) assays. Total phenolic contents of these seaweeds were determined using Folin-Ciocalteu assay. Red seaweed *K. alvarezii*, green seaweed *C. lentillifera* and brown seaweed *S. polycystum* were selected based on their high *in vitro* antioxidant activity, and further evaluated for their chemical composition,



in vivo antioxidant activity and cholesterol-lowering effects in Sprague Dawley rats fed with HCF diet for 16 weeks. Chemical analysis of seaweeds comprised of proximate composition, dietary fiber, vitamin C, vitamin E (α -tocopherol), minerals, carotenoids, chlorophylls, fatty acids and amino acids. Animal experimental diets comprised of eight groups: normal diet (N, control group), HCF diet (HCF group), normal diet supplemented with 5% seaweeds (N+KA, N+CL and N+SP groups), and HCF diet supplemented with 5% seaweed (HCF+KA, HCF+CL and HCF+SP groups). Effects of seaweeds in preventing hypercholesterolaemia and peroxidation in rats were studied via assessing the plasma lipids and, plasma and organs malondialdehyde (MDA) concentrations. Likewise, activities of antioxidant enzymes such as superoxide dismutase (SOD), glutathione peroxidase (GSH-Px) and catalase (CAT) were accessed as indices of oxidative stress. Biochemical markers for liver, heart and kidney damage such as alanine aminotransferase (ALT), aspartate aminotransferase (AST), γ -glutamyltransferase (GGT), creatinine kinase (CK), CK-MB isoenzyme, urea, creatinine and uric acid were measured. Somatic index and descriptive histological changes in the liver, heart, kidney, brain, spleen and eye of the experimental rats were also performed, while quantitative histology was restricted only to necrosis in the liver, kidney and brain. The results showed that administration of *K. alvarezii* and *C. lentillifera* reduced ($P<0.05$) plasma low-density lipoprotein cholesterol and triglyceride, and increased ($P<0.05$) plasma high-density lipoprotein cholesterol thus improving the atherogenic index of rats fed a HCF diet. These seaweeds were shown to reduce body weight gain in rats fed a HCF diet in the following order *S. polycystum*>*C. lentillifera*>*K. alvarezii*. However, *K. alvarezii* and *C. lentillifera* were more effective than *S. polycystum* in

improving the antioxidant status by reducing ($P<0.05$) lipid peroxidation and increasing ($P<0.05$) antioxidant enzymes in liver, heart and kidney of rats fed the HCF diet. Histological examinations demonstrated consumption of all three seaweeds did not exert any damage to the liver, heart, kidney, brain, spleen and eyes in normal rats. In conclusion, *K. alvarezii* and *C. lentillifera* showed hypolipidaemic effects, improve antioxidant status and exert a protective effect in mitigating the cardiac, hepatic, renal and brain abnormalities in rats fed HCF diet. The presence of high dietary fiber especially soluble fiber, omega-3 fatty acids such as eicosapentaenoic acid (C20:5 ω 3), and antioxidant compounds such as polyphenols, vitamin C, α -tocopherol, carotenoids and selenium may probably contributed to the cholesterol-lowering and antioxidant efficacy of these seaweeds.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KOMPOSISI KIMIA, CIRI-CIRI ANTIOKSIDAN DAN PENURUNAN
KOLESTEROL RUMPAI LAUT MALAYSIA TERPILIH**

Oleh

PATRICIA MATANJUN

DISEMBER 2008

Pengerusi : Profesor Dr. Suhaila Mohamed, PhD

Fakulti : Sains dan Teknologi Makanan

Objektif-objektif kajian ini adalah untuk menilai aktiviti antioksidan *in vitro* dan menyaring kandungan fenolik lapan spesies rumpai laut Malaysia (*Kappaphycus alvarezii*, *Eucheuma denticulatum*, *Halymenia durvillaei*, *Caulerpa lentillifera*, *Caulerpa racemosa*, *Dicyota dichotoma*, *Sargassum polycystum* dan *Padina* spp.), menentukan komposisi kimia tiga rumpai laut terpilih yang boleh dimakan dan menyelidik kesan rumpai laut ini ke atas antioksidan, penurunan kolesterol, dan kesan biokimia, morfologi serta ciri-ciri histologi tisu-tisu terpilih tikus-tikus yang diberi makan diet tinggi-kolesterol/tinggi-lemak (HCF). Aktiviti antioksidan *in vitro* lapan spesies rumpai laut dinilai melalui ujian TEAC (kapasiti antioksidan equivalen troloks) dan FRAP (kuasa antioksidan penurunan ferik). Kandungan jumlah fenolik rumpai laut dinilai melalui ujian Folin-Ciocalteu. Rumpai laut merah *K. alvarezii*, rumpai laut hijau *C. lentillifera* dan rumpai laut perang *S. polycystum* telah dipilih berdasarkan aktiviti antioksidan *in vitro* yang tinggi dan dinilai selanjutnya dari segi komposisi kimia,



aktiviti antioksidan *in vivo* serta penurunan kolesterol ke atas tikus Sprague Dawley yang diberi makan diet HCF selama 16 minggu. Komposisi kimia rumpai laut terdiri daripada komposisi proksimat, serabut diet, vitamin C, vitamin E (α -tokoferol), mineral, karotenoid, klorofil, asid lemak dan asid amino. Eksperimen diet haiwan merangkumi lapan kumpulan: diet biasa (kumpulan kawalan, N), diet HCF (kumpulan HCF), diet biasa ditambah 5% rumpai laut (kumpulan N+KA, N+CL dan N+SP) dan diet HCF ditambah 5% rumpai laut (kumpulan HCF+KA, HCF+CL dan HCF+SP). Kesan rumpai laut untuk mencegah hiperkolesterolemia dan peroksidaan dalam tikus dikaji melalui ukuran lipid plasma, dan kepekatan malondialdehid (MDA) dalam plasma serta organ. Begitu juga untuk aktiviti enzim antioksidan seperti superoksid dismutase (SOD), glutathion peroksidase (GSH-Px) dan katalase (CAT) telah dinilai sebagai penunjuk untuk tekanan oksidatif. Penanda biokimia untuk kerosakan hati, jantung dan buah pinggang seperti alanin aminotrasferase (ALT), aspartat aminotransferase (AST), γ -glutamilttransferase (GGT), kreatinin kinase (CK), isoenzim CK-MB, urea, kreatinin dan asid urik telah disukat. Indeks somatik dan perubahan histologi diskriptif dalam hati, jantung, buah pinggang, otak, limpa dan mata tikus kajian juga telah dilakukan, sementara histologi kuantitatif dihadkan hanya kepada nekrosis dalam hati, buah pinggang dan otak. Keputusan menunjukkan *K. alvarezii* dan *C. lentillifera* menurunkan ($P<0.05$) plasma kolesterol lipoprotein ketumpatan rendah dan trigliserid, serta meningkatkan ($P<0.05$) plasma kolesterol lipoprotein ketumpatan tinggi, oleh itu memperbaiki indeks atherogenik tikus yang diberi makan diet HCF. Rumpai laut ini didapati menurunkan penambahan berat badan mengikut susunan *S. polycystum* > *C. lentillifera* > *K. alvarezii*. Walaubagaimanapun, *K. alvarezii* dan *C. lentillifera* didapati

lebih berkesan daripada *S. polycystum* dalam memperbaiki status antioksidan dengan menurunkan ($P<0.05$) peroksidaan lipid dan meningkatkan ($P<0.05$) enzim antioksidan dalam hati, jantung dan buah pinggang tikus yang diberi makan diet HCF. Pemeriksaan histologi mendapati pengambilan *K. alvarezii*, *C. lentillifera* dan *S. polycystum* tidak mengakibatkan kerosakan kepada hati, jantung, buah pinggang, otak, limpa dan mata tikus-tikus normal. Kesimpulannya, *K. alvarezii* dan *C. lentillifera* didapati memberi kesan hipolipidemik, memperbaiki status antioksidan dan kesan perlindungan dengan meringankan ketidaknormalan kardiak, hepatik, renal dan otak tikus yang diberi makan diet HCF. Kehadiran serabut diet yang tinggi khususnya serabut larut, asid lemak omega-3 seperti asid eikosapentaenoik (C20:5 ω 3), dan sebatian antioksidan seperti polifenol, vitamin C, α -tokoferol, karotenoid dan selenium mungkin menyumbang kesan penurunan kolesterol dan keberkesanan antioksidan rumput laut ini.

ACKNOWLEDGEMENTS

First and foremost I would like to thank the Almighty for his blessings and wisdom to enable me to complete this thesis.

I would like to take this opportunity to express my deepest gratitude to my supervisor, Prof. Dr. Suhaila Mohamed for her invaluable suggestions, guidance, motivation and patience throughout the research. I would also like to express my special gratitude to Assoc. Prof. Dr. Noordin Mohamed Mustapha and Assoc. Prof. Dr. Kharidah Muhammad for their constant guidance, support, tremendous encouragement and constructive comments.

Furthermore, I would like to thank the Dean of School of Food Science and Nutrition at Universiti Malaysia Sabah (UMS), Assoc. Prof. Dr. Mohd Ismail Abdullah, for providing additional research funding that assisted in the completion of my research works, as well as for his constant encouragement. I would also like to thank the Ministry of Science, Technology and Innovation of Malaysia (MOSTI) for funding part of this research project, and also to Public Service Department/UMS for the SLAB scholarship.

Special thanks to all the staff of Faculty of Food Science and Technology, and Faculty of Veterinary Medicine at Universiti Putra Malaysia (UPM) and National Institute of Pharmaceutical and Nutraceutical, especially staffs of Animal House, Post Mortem Laboratory, Histopathology Laboratory, Haematology and Clinical Biochemistry Laboratory, and Food Biochemistry Laboratory who had contributed in one way or



another towards the success of this research. Also, I would like to thank Borneo Marine Research Institute at UMS, and Sabah Fisheries Department for supplying the seaweeds and Dr. Charles Vairappan for collection of seaweeds and technical assistance. Also, thank you to Prof. Dr. Cheng Hwee Ming to allow me to use the facilities at the Antioxidant Laboratory at University Malaya.

Special thanks to my friends, Radhiah, Siti Anarita, Imilia, Juliana, Intan, Syida, Yang Hayati, Ooi Kock, Boon Kuan, Zailawati, Sanaz, Dr. Lee and Dr. Yasmin whose help, suggestions, comments and moral supports have helped in the improvement and completion of this thesis. Also thank you to everyone who had contributed in one way or another to the success of this research.

Lastly but not least, my heartfelt gratitude goes to my husband, children and family for their continuous encouragement, patience, understanding, support and unwavering love throughout the years of my study.



I certify that an Examination Committee has met on **16 December 2008** to conduct the final examination of **Patricia Matanjun** on her **Doctor of Philosophy** thesis entitled **“Chemical Composition, Antioxidative and Cholesterol Lowering Properties of Selected Malaysia Seaweeds”** in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the degree of Doctor of Philosophy.

Members of the Examination Committee are as follows:

Mohd. Yazid Abdul Manap, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Azizah Abdul Hamid, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Maznah Ismail, PhD

Professor
Faculty of Medicine and Health Science
Universiti Putra Malaysia
(Internal Examiner)

Ibrahim Jantan, PhD

Professor
Faculty of Pharmacy
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)

BUJANG KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of **Doctor of Philosophy**. The members of the Supervisory Committee were as follows:

Suhaila Mohamed, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Noordin Mohamed Mustapha, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Sharifah Kharidah Syed Muhammad, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

HASANAH MOHD. GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

PATRICIA MATANJUN

Date:



TABLE OF CONTENTS

ABSTRACT	Page
ABSTRAK	iii
ACKNOWLEDGEMENTS	vi
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xiii
LIST OF FIGURES	xviii
LIST OF PLATES	xxi
LIST OF ABBREVIATIONS	xxiii
	xxvi

CHAPTER

1	INTRODUCTION	1
2	LITERATURE REVIEW	
2.1	Seaweed	7
2.2	<i>Kappaphycus alvarezii</i> (Rhodophyta)	8
2.3	<i>Caulerpa lentillifera</i> (Chlorophyta)	12
2.4	<i>Sargassum polycystum</i> (Phaeophyta)	15
2.5	Chemical constituents of seaweeds and potential use as functional food	17
2.5.1	Polysaccharides and dietary fibers	18
2.5.2	Minerals	20
2.5.3	Vitamins	22
2.5.4	Proteins and amino acids	23
2.5.5	Lipids and fatty acids	25
2.5.6	Polyphenols	26
2.5.7	Carotenoids	27
2.5.8	Chlorophylls	27
2.6	Hyperlipidaemia	29
2.7	Atherosclerosis	31
2.8	Free radicals and reactive oxygen species	33
2.9	Lipid peroxidation	35
2.10	Antioxidant defense system	37
2.10.1	Superoxide dismutase	37
2.10.2	Glutathione peroxidase	39
2.10.3	Catalase	40
2.11	Oxidative stress and cardiovascular disease	41



3	ANTIOXIDANT ACTIVITIES AND PHENOLIC CONTENT OF EIGHT SPECIES OF SEAWEEDS FROM SABAH	
3.1	Introduction	43
3.2	Materials and methods	44
3.2.1	Chemicals	44
3.2.2	Raw materials	45
3.2.3	Preparation of extracts	47
3.2.4	Determination of <i>in vitro</i> antioxidant activity	48
3.2.5	Total phenolic content	49
3.2.6	Statistical analysis	50
3.3	Results and discussion	50
3.3.1	Extraction yields	50
3.3.2	TEAC assay	52
3.3.3	FRAP assay	54
3.3.4	Total phenolic content	55
3.3.5	Correlations	56
3.4	Conclusion	58
4	NUTRIENTS CONTENT OF TROPICAL EDIBLE SEAWEEDS, <i>KAPPAPHYCUS ALVAREZII</i>, <i>CAULERPA LENTILLIFERA</i> AND <i>SARGASSUM POLYCYSTUM</i>	
4.1	Introduction	59
4.2	Materials and methods	60
4.2.1	Chemicals	60
4.2.2	Sample preparation	61
4.2.3	Proximate composition	61
4.2.4	Dietary fiber	63
4.2.5	Vitamin C (ascorbic acid)	65
4.2.6	Vitamin E (α -tocopherol)	65
4.2.7	Minerals and trace elements	66
4.2.8	Carotenoids and chlorophylls	67
4.2.9	Fatty acid profile	71
4.2.10	Amino acid profile	71
4.2.11	Statistical analysis	73
4.3	Results and discussion	73
4.3.1	Macro- and micro-nutrient composition	73
4.3.2	Fatty acid profile	87
4.3.3	Amino acid profile	90
4.4	Conclusion	92

5	EFFECT OF SELECTED SEAWEEDS INTAKE ON PLASMA LIPID AND BLOOD ANTIOXIDANT STATUS IN RATS FED NORMAL AND HIGH- CHOLESTEROL/HIGH-FAT DIET	
5.1	Introduction	93
5.2	Materials and methods	95
5.2.1	Chemicals	95
5.2.2	Seaweed material	96
5.2.3	Chemical composition of animal feeds	96
5.2.4	Animal and diets	97
5.2.5	Analytical procedures	101
5.2.6	Statistical analysis	105
5.3	Results and discussion	106
5.3.1	Body weight and food intake	106
5.3.2	Plasma lipids and atherogenic index	112
5.3.3	Lipid peroxidation	121
5.3.4	Enzyme antioxidants activity	124
5.3.5	Seaweed compounds	128
5.4	Conclusion	131
6	EFFECT OF SELECTED SEAWEEDS INTAKE ON THE BIOCHEMISTRY AND MORPHOLOGY OF RATS FED NORMAL AND HIGH-CHOLESTEROL/HIGH-FAT DIET	
6.1	Introduction	133
6.2	Materials and methods	135
6.2.1	Chemicals	135
6.2.2	Seaweed material	135
6.2.3	Chemical composition of animal feeds	135
6.2.4	Animal and diets	135
6.2.5	Analytical procedures	136
6.2.6	Histopathological studies	138
6.2.7	Statistical analysis	138
6.3	Results and discussion	139
6.3.1	Liver	139
6.3.2	Heart	156
6.3.3	Kidney	170
6.3.4	Brain	189
6.3.5	Spleen	196
6.3.6	Lens	199
6.3.7	Seaweed compounds	200
6.4	Conclusion	202

7	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS FOR FUTURE RESEARCH	204
	REFERENCES	209
	BIODATA OF STUDENT	253
	LIST OF PUBLICATIONS	257



LIST OF TABLES

Table		Page
3.1	Extraction yield of seaweeds methanolic and diethyl ether extracts on dry weight basis (DW)	51
3.2	Antioxidant activities of seaweeds methanolic extracts determined by TEAC and FRAP assays	53
3.3	Total phenolic (TP) contents of seaweeds methanolic dry extracts expressed as phloroglucinol equivalents (PGE)	55
4.1	Gradient program used for the separation of seaweed pigments	70
4.2	Nutrient composition of seaweeds <i>K. alvarezii</i> , <i>C. lentillifera</i> and <i>S. polycystum</i> (mean \pm S.E.M., % dry weight of sample)	74
4.3	Calibration relations, method precision and retention times for the different pigment standard analyzed	81
4.4	Carotenoids and chlorophylls content of seaweeds <i>K. alvarezii</i> , <i>C. lentillifera</i> and <i>S. polycystum</i> (% dry weight of sample)	82
4.5	Pigment identified of acetone extract of <i>S. polycystum</i> by LC-DAD-ESI(+) without standard available	83
4.6	Fatty Acid Content (% of total fatty acid content) of <i>K. alvarezii</i> , <i>C. lentillifera</i> and <i>S. polycystum</i>	88
4.7	Amino Acid Content (mg/g dry weight of sample) of <i>K. alvarezii</i> , <i>C. lentillifera</i> and <i>S. polycystum</i>	91
5.1	Ingredients of the experimental diets (1000g feed/batch)	98
5.2	Proximate composition and total energy (kcal/100g feed) of the experimental feeds, and calories intake of rats from eight experimental diets	100
5.3	Body weight gain of rats of all groups after 16 weeks of experimental diet	108
5.4	Total cholesterol (TC) in plasma of rats during experimental period	114
5.5	Low-density cholesterol (LDL-C) in plasma of rats during experimental period	116



5.6	High-density cholesterol (HDL-C) in plasma of rats during experimental period	118
5.7	Triglycerides (TG) in plasma of rats during experimental period	119
5.8	Atherogenic Index (LDL-C/HDL-C) of all groups at the end of the experimental period	121
5.9	Malondialdehyde (MDA) level in plasma of rats during experimental period	122
5.10	SOD activity in plasma of rats during experimental period	125
5.11	Glutathione peroxidase (GSH-Px) activity in plasma of rats during the experimental period	126
5.12	Catalase (CAT) activity in plasma of rats during the experimental period	128
6.1	Liver weight and hepatosomatic index (liver weight/terminal body weight x 100) of rats at the end of the experimental period	141
6.2	Quantitative histology in liver of experimental groups (Mean±S.E.M.)	142
6.3	ALT (U/L) in plasma of rats during the experimental period	148
6.4	AST (U/L) in plasma of rats during the experimental period	150
6.5	GGT (U/L) in plasma of rats during the experimental period	152
6.6	Malondialdehyde level and enzymic antioxidants activities in liver of rats at the end of the experimental period	154
6.7	Heart weight (g) and cardiosomatic index (organ weight/body weight x 100) of rats at the end of the experimental period	157
6.8	Quantitative histology scores of lesion in the heart of experimental groups (Mean±S.E.M.)	158
6.9	Creatinine kinase (CK) of plasma of rats during the experimental period	166
6.10	Creatinine Kinase-MB (CK-MB) isoenzyme of plasma of rats during the experimental period	166

6.11	Malondialdehyde level and enzymic antioxidants activities in heart of rats at the end of the experimental period	167
6.12	Kidney weight and nephrosomatic index (kidney weight/terminal body weight x 100) of rats at the end of the experimental period	170
6.13	Quantitative histology in kidney of experimental groups (Mean±S.E.M.)	171
6.14	Urea (mmol/L) in plasma of rats during the experimental period	179
6.15	Creatinine (µmol/L) in plasma of experimental rats during the experimental period	181
6.16	Uric acid (mmol/L) in plasma of rats during the experimental period	183
6.17	Malondialdehyde level and enzymic antioxidants activities in kidney of rats at the end of the experimental period	186
6.18	Brain weight and encephadosomatic index (organ weight/terminal body weight x 100) of rats at the end of the experimental period	190
6.19	Quantitative histology in brain of experimental groups (Mean±S.E.M.)	192
6.20	Malondialdehyde level and enzymic antioxidants activities in brain of rats at the end of the experimental period	194
6.21	Spleen weight and splenosomatic index (organ weight/terminal body weight x 100) of rats at the end of the experimental period	196
6.22	Malondialdehyde level and enzymic antioxidants activities in spleen of rats at the end of experimental period	199

LIST OF FIGURES

Figures		Page
2.1	Photograph of various colour strains of <i>K. alvarezii</i> and one strain of <i>Eucheuma denticulatum</i> (place middle top) cultivated in Banggi Island, Sabah	9
2.2	Photograph illustrated seaweed cultivation using floating long-line system which works well in deep water areas. Farmers harvesting seaweed that grows on nylon line (located at Semporna waters, Sabah)	11
2.3	Photograph illustrated another seaweed cultivation method using the floating raft method made from square bamboo timber frame and seaweeds attached to the rope (located at Banggi Island, Sabah)	11
2.4	Photograph of <i>Caulerpa lentillifera</i> or also known as “lato” collected from Semporna waters, Sabah	14
2.5	Photograph of <i>Caulerpa racemosa</i> collected from Kota Kinabalu waters, Sabah	14
2.6	Photograph of <i>Sargassum polycystum</i> collected from wild population at Kota Kinabalu waters, Sabah	16
3.1	Correlation of FRAP and TEAC values of seaweed methanolic dry extracts from (1) <i>Kappaphycus alvarezii</i> , (2) <i>Eucheuma denticulatum</i> , (3) <i>Halymenia durvella</i> , (4) <i>Caulerpa lentillifera</i> , (5) <i>Caulerpa racemosa</i> , (6) <i>Dicyota dichotoma</i> , (7) <i>Sargassum polycystum</i> , (8) <i>Padina</i> spp, (9) BHT and (10) Quercetin	57
3.2	Correlation of FRAP and TP content of seaweed methanolic extracts from (1) <i>Kappaphycus alvarezii</i> , (2) <i>Eucheuma denticulatum</i> (3) <i>Halymenia durvella</i> , (4) <i>Caulerpa lentillifera</i> , (5) <i>Caulerpa racemosa</i> , (6) <i>Dicyota dichotoma</i> , (7) <i>Sargassum polycystum</i> and (8) <i>Padina</i> spp.	57
4.1	LC-ESI(+)-MS chromatogram of acetone extract from <i>K. alvarezii</i> (1) lutein (RT = 12.63 min), (2) chlorophyll <i>a</i> (RT = 27.38 min)	79
4.2	LC-ESI(+)-MS chromatogram of acetone extract from <i>C. lentillifera</i> (1) lutein (RT = 12.62 min), (2) chlorophyll <i>b</i> (RT = 22.03 min), (3) chlorophyll <i>a</i> (RT = 27.42 min), (4) phaeophytin <i>a</i> (RT = 42.55 min)	80
4.3	LC-ESI(+)-MS chromatogram of acetone extract from <i>S. polycystum</i> *(1) fucoxanthin (RT = 6.95 min), *(2) dinoxanthin (RT = 8.42 min), (3) zeaxanthin (RT = 12.57 min), (4) chlorophyll <i>b</i> (RT = 22.00 min), (5) chlorophyll <i>a</i> (RT = 27.37 min), * means tentative identification	80

4.4	On-line UV-vis spectra of fucoxanthin as obtained by LC-DAD analysis of acetone extract of <i>S. polycystum</i>	84
4.5	Mass spectra of fucoxanthin via ESI(+) interface	84
4.6	On-line UV-vis spectra of dinoxanthin as obtained by LC-DAD analysis of acetone extract of <i>S. polycystum</i>	85
4.7	Mass spectra of dinoxanthin via ESI(+) interface	85
4.8	Molecular structure of fucoxanthin	86
4.9	Molecular structure of dinoxanthin	86
5.1	Body weight of rats fed with normal (N) and high-cholesterol/high-fat (HCF) diets with and without seaweeds supplementation for 16 weeks	107
5.2	Photograph showing HCF rat (top) had higher amount of adipose tissue as compared to N rat (bottom)	110
5.3	Photographs showing HCF+KA (top) rat, HCF+CL (middle) rat and HCF+ SP (bottom) rat had less amount of adipose tissue compared to HCF rat	111
6.1	Photograph of a liver of rat from HCF group. The liver appeared much paler than normal, soft, mottled and fatty	139
6.2	Photograph of liver of rats from HCF+CL (left) and HCF+KA (right) groups respectively. Both livers although were slightly pale but were less fatty	140

LIST OF PLATES

Plates		Page
6.1	Photomicrograph, liver of rat from N group at necropsy showed normal hepatic cells with well-preserved hepatic cords and devoid of evidence of degenerative cells and inflammation [H&E, x200]	144
6.2	Photomicrograph, liver of rat from HCF group at necropsy showed severe and extensive fatty changes and foci of inflammation [H&E, x200]	144
6.3	Photomicrograph, liver of rat from N group at necropsy. Necrotic and Kupffer cells are within normal limits [H&E, x400]	145
6.4	Photomicrograph, liver of rat from HCF group at necropsy. Vacuolations occur and loss of normal hepatic architecture. Note the presence of foci of inflammation within the portal triad [H&E, x400]	145
6.5	Photomicrograph, liver of rat from HCF+KA group at necropsy. Normal parenchymal structure with occasional intracytoplasmic vacuolation. Histological features of the hepatocytes almost bear resemblance to that of the N group [H&E, x400]	146
6.6	Photomicrograph, liver of rat from HCF+CL group at necropsy. Histological features of the hepatocytes bear resemblance to that of the N group [H&E, x400]	146
6.7	Photomicrograph, liver of rat from HCF+SP group at necropsy. Note the presence of intracytoplasmic vacuoles within the hepatocytes but less extensive compared to that seen in the HCF group [H&E, x400]	147
6.8	Photomicrograph, heart of rat from N group at necropsy. Note that the size and orientation of the myocardial fibers are within normal limits [H&E, x100]	159
6.9	Photomicrograph, heart of rat from HCF group at necropsy. Note that most of the myocardial fibers are homogenously pinkish, lacks nuclei and hypertrophied. Note the presence of fibroblastic infiltration between muscle fibers (arrows) [H&E, x100]	159
6.10	Photomicrograph, heart of rat from HCF group at necropsy depicting numerous intracytoplasmic “lipid” vacuolation (L) interspersed within myocardial fibers [H&E, x400]	160

6.11	Photomicrograph, heart of rat from HCF group at necropsy. In some areas, fibrosis (F), inflammation (I), intracytoplasmic vacuolation (V) and necrosis (N) are also evident [H&E, x400]	160
6.12	Photomicrograph, heart of rat from HCF+KA group at necropsy. Note myocardial fibers are within normal appearance with histological features bearing resemblance to that of the N group [H&E, x100]	161
6.13	Photomicrograph, heart of rat from HCF+CL group at necropsy. Note myocardial fibers are within normal appearance with histological features bearing resemblance to that of the N group [H&E, x100]	161
6.14	Photomicrograph, heart of rat from HCF+SP group at necropsy. Note that although the orientation of muscle fibers are that of normal appearance but the individual cardiac muscle appeared slightly enlarged [H&E, X 100]	162
6.15	Photomicrograph, heart of rat from HCF+SP group at necropsy. Note marked loss of cardiocytic mass and the heavy infiltration of the loss areas by fibrous tissue (F). Furthermore, the remaining cardiomyocytes are either enlarged or depicting degenerative or necrotic changes [H&E, x400]	162
6.16	Photomicrograph, kidney of rat from N group at necropsy. Note the glomeruli and tubules of kidney are within normal appearance [H&E, x100]	172
6.17	Photomicrograph, kidney of rat from HCF group at necropsy. Note the presence of mononuclear inflammatory cell infiltration (I) and thickened glomerular basement membranes (arrow) indicating glomerulopathy [H&E, x100].	172
6.18	Photomicrograph, kidney of rat from N group at necropsy at higher magnification shows a normal structure of glomerulus and tubules [H&E, x200]	173
6.19	Photomicrograph, kidney of rat from HCF group at necropsy. The loss of whole glomerus (G) and tubular dilation (T) were observed [H&E, x400]	173
6.20	Photomicrograph, kidney of rat from HCF group at necropsy. Note the expansion of mesangium (A) [H&E, x400]	174
6.21	Photomicrograph, kidney of rat from HCF group at necropsy. Note the foamy appearance (arrows) in the tubular epithelial cells [H&E, x400]	174