



**UNIVERSITI PUTRA MALAYSIA**

**CHARACTERIZATION OF *Bacillus cereus* ISOLATED FROM  
READYTO-EAT CEREALS**

**LEE HAI YEN**

**FSTM 2009 2**

**CHARACTERIZATION OF *Bacillus cereus* ISOLATED  
FROM READY-TO-EAT CEREALS**

**LEE HAI YEN**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**2009**



**CHARACTERIZATION OF *Bacillus cereus* ISOLATED FROM READY-  
TO-EAT CEREALS**

**By**

**LEE HAI YEN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia in Fulfillment of the Requirements for the Degree of Master of  
Science**

**2009**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science

**CHARACTERIZATION OF *Bacillus cereus* ISOLATED FROM READY-TO-EAT CEREALS**

**By**

**LEE HAI YEN**

**APRIL 2009**

**Chairman : Farinazleen Mohamad Ghazali, PhD**

**Faculty : Faculty of Food Science and Technology**

Since availability of data on the contamination of spore formers such as *B. cereus* in ready-to-eat foods is scarce particularly from developing countries, this surveillance was conducted to address the issue. The preliminary findings from this study revealed a high prevalence of *B. cereus s.l.* being detected using MPN-PCR in 76% of 111 samples of ready-to-eat cereals tested. The range of concentration was from 30 MPN/g to more than 24,000 MPN/g. Results indicated the differences in the level of contamination for *B. cereus s.l.* in various products based on factors such as product types, ingredients added and location of manufacturer. The highest concentration of *B. cereus s.l.* was found in samples with ingredients from weaning products which are, product made from vegetable origin.



The alarming findings on the high prevalence of *B. cereus s.l.* in ready-to-eat cereals prompted further studies on the isolates from these samples. Isolation of colonies from these samples were characterized based on toxin gene screening, plasmid profiles, antibiotic resistance and fingerprinted using RAPD-PCR analysis. In addition to the high prevalence in RTE cereals, the toxin screening profile indicated 58% of the isolate carry the *Bacillus* enterotoxin T (*bceT* gene) and 34% carries the tri-component non-hemolytic enterotoxin (*nhe* gene). This shows that majority of the isolates from ready-to-eat cereals are diarrheagenic. The plasmid profile revealed one isolate carrying the plasmid size similar to cereulide protein which is responsible for the emetic disease therefore showing the less common prevalence of emetic isolates from RTE cereals. Isolates of *B. cereus s.l.* were found to be resistant to ampicillin, metronidazole but highly susceptible to antibiotics with mechanism of action that inhibits the protein synthesis such as erythromycin, oxytetracycline, spectinomycin, neomycin, furozolidone, quinopristin/dalfopristin. Even though *B. cereus* usually manifests a self limiting disease, the emergence of antibiotic resistance bacteria is a major concern worldwide and due to ease of horizontal gene transfer between the



subspecies of *B. cereus s.l.*, the risk of other more lethal subspecies such as *B. anthracis* obtaining the antibiotic resistance gene may also occur.

This preliminary finding revealed an interesting risk profile for *B. cereus s.l.* as there are no known data available for the assessment on the microbiological quality of RTE cereals especially for spore formers. Based on the risk assessment study conducted, an estimate of on worst case scenario showed 48 diarrhea cases in 27 million populations are known to occur per annum from consumption of RTE cereals.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk Ijazah Sarjana

**PENCIRIAN *Bacillus cereus* YANG DIPENCILKAN DARI BIJIRIN SEDIA  
DIMAKAN**

Oleh

**Pengerusi : Farinazleen Mohd. Ghazali, PhD**

**Fakulti : Fakulti Sains dan Teknologi Makanan**

**ABSTRAK**

Sejak kesediaan data mengenai kontaminasi pembentuk spora seperti *B. cereus* dalam bijirin sedia dimakan adalah terhad terutamanya dari negara yang sedang membangun, pengawasan ini telah dijalankan untuk mengalamatkan isu tersebut. Hasil penyelidikan preliminari daripada kajian ini menunjukkan kekerapan *B. cereus s.l.* yang dikesan menggunakan MPN-PCR dalam 76% daripada 111 sample bijirin sedia dimakan yang telah diuji. Jajaran konsentrasi didapati adalah dari 30 MPN/g sehingga melebihi 24,000 MPN/g. Keputusan ini menunjukkan perbezaan dalam tahap kontaminasi bagi *B. cereus s.l.* dalam pelbagai produk berdasarkan faktor-faktor seperti jenis produk, ingredients tambahan serta lokasi manufacturer. Konsentrasi *B. cereus s.l.* yang tertinggi didapati dalam sampel yang mengandungi produk 'weaning' iaitu produk yang dihasilkan dari sayuran.



Keputusan yang menggelisahkan mengenai kekerapan *B. cereus s.l.* yang tinggi daripada bijirin sedia dimakan telah menyaran ujikaji lanjutan ke atas pencilan koloni dari sampel-sampel makanan. Pemencilan koloni-koloni dicari berdasarkan pemeriksaan gen toxin, profail plasmid, kerintangan antibiotic serta analisis RAPD-PCR fingerprint. Profail yang didapati dari kajian ini menunjukkan 58% daripada pencilan mengandungi *Bacillus enterotoxin T* gen (*bceT* gen) dan 34% daripada pencilan mengandungi tiga komponen 'non-hemolytic enterotoxin (*nhe* gen). Ini menunjukkan bahawa majoriti daripada pencilan dari bijirin sedia dimakan adalah jenis cirit. Profail plasmid memperlihatkan satu pencilan yang mengandungi plasmid saiz yang serupa kepada protein 'cereulide' yang menyebabkan muntah-muntah oleh itu menunjukkan kekerapan yang kurang lazim untuk pencilan jenis muntah dari bijirin sedia dimakan. Pencilan *B. cereus s.l.* ditemui menjadi rintangan kepada ampisilin dan metronidazole tetapi mudah dipengaruhi oleh antibiotik dengan mekanisme tindakan yang mencegah sintesis protein seperti erythromycin, oxytetracycline, spectinomycin, neomycin, furozolidone dan quinopristin/dalfopristin. Walaupun *B. cereus* menyebabkan penyakit batas sendiri, kemunculan bakteria rintangan antibiotic adalah satu perhatian utama di dunia. Oleh kerana kemudahan pemindahan horisontal gen antara subspesies *B. cereus s.l.*, risiko subspesies





yang lebih maut seperti *B. anthracis* memperoleh gene kerintangan antibiotik mungkin juga akan berlaku.

Hasil penemuan preliminari menunjukkan satu profil risiko yang menarik bagi *B. cereus s.l.* kerana ketiadaan data untuk penilaian kualiti mikrobiologi untuk bijirin sedia dimakan terutamanya bagi pembentuk spora. Berdasarkan kajian penilaian risiko yang dijalankan, taksiran bagi senario paling buruk menunjukkan 48 kes cirit dalam 27 juta populasi berlaku setiap tahun daripada pemakanan bijirin sedia dimakan.

## ACKNOWLEDGMENT

I have to thank God for everything, for being able to come this far; I thank God for giving me strength and courage when I falter; thank God for my family because they taught me to be me and believe in myself even if I have to defy them; thank God for the fate that brought me to my supervisor (Dr. Farin) and co-supervisor (Prof Son), which opened up this road and bring me a step closer to my dreams; thank God for my friends (especially Lay Ching who is always there for everything, Grace for keeping my off lab hours sane, Siow Yin and Wai Yee for their support) because they have exposed me to many lab and life experiences that helped me grow; thank God for putting me through difficult and embarrassing moments to keep me humble and braver; and thank God for keeping me alive so I will always know how little I have and how precious things are and keep moving forward with these cherished ones.

I believe that words are incapable of describing even a fraction of my emotions and the depth of my sincerest gratitude for this opportunity and journey however, in this section, I would just like these people whom I have met along this journey of life to know that the guidance, lessons, memories



and knowledge shared will always be cherished and remembered for the rest of my life and beyond.

This thesis is especially dedicated to my three elder sisters and mother for sacrificing everything because even at their most difficult trying times, they are still able to love the most difficult person on earth. Thank you so much for still loving me especially through the times when I can't even love myself and for those difficult times, you have made me what I am today.

Thank you once again, for there is no known depth for my gratitude.



I certify that an Examination Committee has met on 14<sup>th</sup> April 2009 to conduct the final examination of LEE HAI YEN on her Master of Science thesis entitled “Characterization of *Bacillus cereus* isolated from ready-to-eat cereals” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Act 1981. The committee recommends that the student be awarded the Master of Science in Food Safety.

Members of the Examination Committee were as follows:

**Roselina Karim, PhD**

Lecturer  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Shuhaimi Mustafa, PhD**

Associate Professor  
Deputy Director  
Institut Penyelidikan Produk Halal  
Universiti Putra Malaysia  
(Internal Examiner)

**Fatimah Abu Bakar, PhD**

Associate Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Internal Examiner)

---

**BUJANG KIM HUAT, PhD**

Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science in Food Safety. The members of the Supervisory Committee were as follows:

**Farinazleen Mohamad Ghazali, Ph.D**

Lecturer  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Son Radu, Ph.D**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

**Jinap Selamat, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

---

**HASANAH MOHD GHAZALI, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 8 June 2009



## DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

---

LEE HAI YEN

Date: 24 July 2009



## LIST OF TABLES

Table		Page
2.1	Characteristics of <i>B. cereus s.l.</i> group and the biochemical tests for differentiation and identification of the subspecies	14
2.2	Incidences of foodborne diseases based on the bacterial cause (Source:Kramer and Gilbert, 1989).	25
3.1	Enumeration for <i>Bacillus</i> spp. and <i>B. cereus</i> indicating the concentration (in MPN/g) and occurrence of food samples containing the respective counts.	48
3.2	Summary on the concentration of <i>Bacillus</i> spp. and <i>B. cereus</i> in the various types of flavors in food samples indicating the number of samples tested, range and median of cell concentration	51
4.1	summarizes the primer sequences, PCR product sizes and annealing temperature for the toxin gene screening.	63
5.1	Antibiogram of <i>B. cereus</i> strains isolated from RTE cereals indicating the general mechanism of action of the antibiotic and the percentage of isolates showing 'sensitive', 'intermediate' or 'resistant' to the antibiotic test.	78



6.1	shows the work chart of a preliminary stepwise risk assessment of the expected number of cases in various scenarios for general population (A-C) and in children population (D-F)	100
-----	---	-----





## LIST OF FIGURES

Figure		Page
2.1	Gram stain of <i>B. cereus</i> NCTC 11143 under 400 × magnifications (produced in laboratory)	12
3.1 (a)	Agarose gel electrophoresis for enumeration for genus <i>Bacillus</i> spp. producing the size of 1114 bp PCR product. M- indicates the 100 bp molecular ladder (VIVANTIS, Malaysia). 1- <i>B. cereus</i> control NCTC11143, 2-9: DNA extraction from food samples.	44
3.1 (b)	Agarose gel electrophoresis for detection of <i>B. cereus</i> with product size of 413 bp.  Lane 1 – <i>B. cereus</i> NCTC 11143 (positive control),  Lane 2-10: DNA extraction from food sample	44
3.3	The types of products tested and the median concentration of <i>Bacillus</i> spp. (in dark orange) and <i>B. cereus</i> (in light orange).	50
3.4	Illustration for comparing the median of <i>Bacillus</i> spp. and <i>B. cereus</i> in various types of flavors added to the samples. Weaning food shows the highest	



	concentration of bacteria compared to the other types	52
3.5	illustrates the enumeration range for <i>Bacillus</i> spp. in products of local and imported origin showing the differences on the frequency of contamination level in RTE products	53
3.6	Prevalence of <i>B. cereus</i> samples of local and foreign origin	54
4.1	Toxin screening for bceT generates amplicon size of 428 bp. M: 100 bp molecular ladder, C: PCR mix without DNA (negative control), 1: <i>B. cereus</i> NCTC 11143, 2-13: isolates from RTE cereals	64
4.2	Multiplex toxin screening for detection of <i>nheA</i> , <i>nheB</i> and <i>nheC</i> genes from food isolate. M: molecular 100 bp ladder, 1-3: isolates from RTE samples , C: <i>B. cereus</i> NCTC 11143 positive control, 4-9: isolates from RTE samples.	66
4.3	Pie chart showing the percentages of isolates carrying the three components non-hemolytic enterotoxin gene ( <i>nheA</i> , <i>nheB</i> and <i>nheC</i> ) and the various combinations of genes.	66



5.1	Flowchart of plasmid isolation protocol	73
5.2	Schematic diagram of plasmid pattern types observed among isolates of <i>B. cereus</i> . Number 1-24 indicates the plasmid pattern observed among isolates. Shaded columns indicate plasmid pattern types that are in Cluster I (refer to next page, Figure 5.3 on clustering).	83
5.3	Dendogram showing the clustering based on plasmid pattern types (number 1-24 from Figure 5.2) indicating two major clusters observed.	84
5.4	shows the clusters generated based on RAPD fingerprinting using Gel Compar and the corresponding characteristics of the isolates based on origin of samples, prevalence of toxin genes, antibiotic resistances and plasmid sizes.	86
6.1	Fitted line plot for estimating the $r$ value, which is the probability of illness occurring in consumption of one <i>B. cereus</i> cells. Equation obtained expresses the probability of illness per MPN/g consumed (data derived from McElroy et al., 1999).	94



## LIST OF ABBREVIATIONS

BAM	Bacteriological Analytical Manual
bceT	Bacillus cereus enterotoxin T
DNA	Deoxyribonucleic acid
GMP	good manufacturing practice
HACCP	hazard analysis critical control point
MPN	Most Probable Number
MYP	Mannitol Yolk Polymyxin
NHE	non-hemolytic enterotoxin
PCR	Polymerase chain reaction
RAPD	Randomly Amplified Polymorphic DNA
RBC	red blood cell
rpm	revolution per minute
RTE	ready-to-eat
<i>s.l.</i>	<i>sensu lato</i>
<i>s.s.</i>	<i>sensu stricto</i>
VP	Voges-Proskauer
WHO	World Health Organization



## TABLE OF CONTENTS

	PAGE
ABSTRACT	i
ABSTRAK	iv
ACKNOWLEDGMENTS	vii
APPROVAL	viii
DECLARATION	xii
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xix
<b>CHAPTER</b>	
1 <b>INTRODUCTION</b>	1
2 <b>LITERATURE REVIEW</b>	5
2.1 Epidemiology Of <i>Bacillus</i> Spp. and <i>Bacillus cereus</i>	5
2.2 <i>Bacillus</i> spp.	7
2.3 <i>Bacillus cereus</i>	11
2.3.1 Identification Of <i>B. cereus</i>	13
2.4 Methods Of Detection For <i>B. cereus</i>	15
2.4.1 Most Probable Number	16
2.4.2 Polymerase Chain Reaction	17
2.5 <i>Bacillus</i> In Food Industry	19
2.5.1 Survival Of <i>B. cereus</i>	22
2.5.2 Incidences	24
2.5.3 <i>B. cereus</i> In Cereals	25
2.6 Hazard Characterization	27
2.6.1 Toxins In <i>B. cereus</i>	27
2.7 Antibiotic Resistances	29
2.7.1 Mechanism Of Resistance	30
2.8 Food Safety	32
2.8.1 Risk Assessment	34



<b>3</b>	<b>PREVALENCE AND ENUMERATION OF <i>Bacillus cereus</i> IN READY-TO-EAT CEREALS</b>	
	3.1 Introduction	38
	3.2 Materials And Methods	40
	3.3 Results And Discussion	45
	3.4 Conclusion	59
<b>4</b>	<b>SCREENING OF TOXIN GENES AMONG <i>Bacillus cereus</i> ISOLATED FROM READY-TO-EAT CEREALS</b>	
	4.1 Introduction	60
	4.2 Materials And Methods	62
	4.3 Results And Discussion	64
	4.4 Conclusion	70
<b>5</b>	<b>CHARACTERIZATION OF <i>Bacillus cereus</i> ISOLATES BASED ON PLASMID PROFILES, ANTIBIOTIC RESISTANCES AND RAPD FINGERPRINTING</b>	
	5.1 Introduction	71
	5.2 Materials And Methods	73
	5.3 Results And Discussion	77
	5.4 Conclusion	90
<b>6</b>	<b>A PRELIMINARY STEPWISE RISK ASSESSMENT ON READY-TO-EAT CEREALS</b>	
	6.1 Introduction	92
	6.2 Materials And Methods	93
	6.3 Results And Discussion	100
	6.4 Conclusion	103
<b>7</b>	<b>SUMMARY, GENERAL CONCLUSION AND RECOMMENDATION FOR FUTURE RESEARCHES</b>	104
	<b>REFERENCES</b>	106
	<b>APPENDICES</b>	
	<b>BIODATA OF STUDENT</b>	



## CHAPTER 1

### INTRODUCTION

The resilience of spore formers has become a major concern in food safety. Particularly for *Bacillus cereus sensu lato* which is the parent group of six subspecies (*Bacillus cereus sensu stricto*, *Bacillus anthracis*, *Bacillus thuringiensis*, *Bacillus mycoides*, *Bacillus pseudomycoides* and *Bacillus weihenstephanensis*), this group has increasingly getting attention because what initially thought as the *B. cereus s.s* as the only foodborne pathogen, it has been found that other subspecies such as *B. thuringiensis* may have accounted to the outbreaks caused by *B. cereus s.s* (Jackson et al., 2008). This is due to the fact that the subspecies does not only possess a high homology in genomic DNA sequences, the prevalence of toxins and different symptoms it manifests also differs within the single subspecies such as *B. cereus s.s*. For example, three various strains of *B. cereus s.s*. have been analyzed due to the involvement in the recent several outbreaks and fatalities (Lund et al., 2000) and another type of strain for food poisoning that does not produce any known toxin (Lapidus et al., 2008). Recently, anthrax-like toxin possessed by *B. cereus* isolates has been detected to cause severe respiratory illnesses (Hoffmaster et al., 2004) from food. Toxins from other subspecies such as *B.*



*weihenstephanensis*, a psychrotrophic strain surviving causing problems in milk contamination were also found to be involved other foodborne outbreaks (Stenfors Arnesen et al., 2008).

Since the changing trend of pathogenesis on *B. cereus* extends to its six subspecies, this issue calls for a greater concern and re-assessment on products not only as ready-to-eats but also the fact that these foods are also consumed by the more vulnerable individuals such as children and convalescing patients. Therefore, it is important to have at least a baseline data to serve as a guide for developing a system on food safety monitoring so that food hazards can be identified in an early stage to prevent the unwanted outcomes. These data requires information on characteristics and behavior as well as point of entry into the food chain (Kleter and Marvin, 2008).

In comparison to the other types of pathogens such as *Salmonella* spp., *Campylobacter* spp., *E. coli* O157:H7 and *L. monocytogenes*, data availability of *B. cereus s.l.* as part of surveillance study in developing countries are scarce or not available. To date, there is only one appraisal of microbiological quality of ready-to-eat cereals by Rani et al. (2005) but that did not include spore formers which have attributed to several outbreaks of severe foodborne





poisoning as well as fatal outbreaks found in infant foods. Despite the sporadic outbreaks of *B. cereus* in foods, there are no known surveillance data available particularly from the developing region.

The practice of microbial risk assessment is increasingly applied in the assessment of food hazards because of the changes in foodborne diseases such as lower infectious dose of  $10^3$  to  $10^4$  cells *B. cereus* have been reported to be responsible for an outbreak case (Notermans et al., 1997). MRA data on *B. cereus* is scarce in the recent years indicating the need to address and update the evolving hazards of *B. cereus* in foods. In Malaysia, the Ministry of Health has recently established the Food Safety and Quality division in 2002 to monitor the food safety issues in Malaysia. The first microbiological risk assessment report was published on the contamination of *Vibrio parahaemolyticus* in tiger prawns, which aided the country's management of food quality when facing international trade regulations. This shows the importance of risk assessment studies which generates data not only for the national surveillance but also for the international bodies.

