



**UNIVERSITI PUTRA MALAYSIA**

**ASSESSMENT OF *Bifidobacterium pseudocatenulatum* G4 AS A  
PROBIOTIC**

**BARKA MOHAMMED KABEIR**

**FSTM 2009 17**

**ASSESSMENT OF *Bifidobacterium pseudocatenulatum* G4 AS A  
PROBIOTIC**

**BARKA MOHAMMED KABEIR**

**DOCTOR OF PHILOSOPHY  
UNIVERSITI PUTRA MALYSIA**

**2009**



**ASSESSMENT OF *Bifidobacterium pseudocatenulatum* G4 AS A  
PROBIOTIC**

**By**

**BARKA MOHAMMED KABEIR**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of  
Philosophy**

**March 2009**



## **DEDICATION**

The work is dedicated to my parents Mohd Kabeir and Hawa' Nurain; to all members of my big family. It also goes to teachers, scientists, researchers, and all seekers for knowledge.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

**ASSESSMENT OF *Bifidobacterium pseudocatenulatum* G4 AS A  
PROBIOTIC**

By

**BARKA MOHAMMED KABEIR**

**March 2009**

**Chairman: Professor Mohd Yazid Abdul Manap, PhD**

**Faculty : Food Science and Technology**

This study has demonstrated the probiotic criteria of *Bifidobacterium pseudocatenulatum* G4 related to safety profile; effects on intestinal microbiota; interactions with prebiotics; and the growth in peanut milk (PM) and survivability during the storage. The safety profile of the strain G4 was evaluated in groups of BALB/c mice, fed different concentrations of *Bifidobacterium* strains for a period of 28 days. At the end of the experimental period, the highest dose of 11 log CFU *B. pseudocatenulatum* G4 /day did not affect the general health of mice nor cause any toxicity to blood based on biochemistry and haematology measurements. Pathogenicity symptoms were not detected in the internal organs; serum enzymes of liver and kidney; and histology of ceacum, ileum and colon of the treated mice; thus, the strain G4 could be a safe probiotic for food application. The effects of feeding strain G4 on selected microbiota community and metabolic activity in ceacum and colon of rats were also examined in 30



days feeding trail. The rat groups received skim milk contained either viable or metabolic product of *B. pseudocatenulatum* G4. The viable supplement of strain G4 increased profile of bifidobacteria, Lactobacillus and total anaerobes in ceacum and colon of the fed rats; while it decreased total aerobes and potential pathogens (staphylococcus, enterococcus and enterobacteriaceae) of the same intestinal regions. However, salmonella and coliform were maintained mostly unchanged. In addition, the viable supplement of strain G4 has significantly ( $p < 0.05$ ) increased short chain fatty acids (acetic, propionic, and butyric) of ceacum and colon. Commercial prebiotics [inulin and fructo-oligosaccharide (FOS)], together with sorbitol, arabinan and inoculum concentration, were tested by fractional factorial design (FFD) to determine their impact on growth of *B. pseudocatenulatum* G4 in skim milk (SM) medium. At 48h incubation, bacterial growth was mainly influenced by FOS and inoculum concentration. Central composite design (CCD) was adopted using FOS and inoculum concentration at 48h incubation to develop the statistical model for optimization. The model predicted that 2.46 log CFU/ml produced the optimum growth increase of *B. pseudocatenulatum* G4. The combination that produced the optimum point was 2.86% FOS (g/v) and 0.67% inoculum concentration (v/v). At this point of optimum combination, validation experiment recorded  $2.40 \pm 10.02$  log CFU/ml. Further application in 1-L bioreactor for 24h showed higher growth of 2.95 log CFU/ml as compared to 2.46 CFU/ml of the optimum model prediction. The growth of strain G4 was evaluated in peanut milk (PM), and growth amounting 7.12 log CFU/ml was attained at 24h incubation. Supplementation with FOS produced growth of 8.35 log CFU/ml, which comparable to the 8.25 log CFU/ml



of Trypticase phytone yeast extract (TPY) obtained at 21h incubation. During the storage of fermented PM at 25 °C for a period of two weeks, the viable cells of strain G4 decreased to a level  $< 7 \log \text{ CFU/ml}$ , which did not fulfill the requirement of probiotics food. However, at two weeks refrigeration (4 °C) storage, the viability of strain G4 was maintained high at acceptable level of  $> 7 \log \text{ CFU CFU/ml}$  fermented PM products. Therefore, *B. pseudocatenulatum* G4 assessed to be a safe probiotic microorganism for synbiotics formulation with FOS and incorporation into peanut milk (PM) for delivery purposes.



Abstrak tesis yang dikemukakan Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**Penilaian *Bifidobacterium pseudocatenulatum* G4 sebagai bakteria probiotik**

Oleh

**BARKA MOHAMMED KABEIR**

**Mac 2009**

**Pengerusi : Profesor Mohd Yazid Abdul Manap, PhD**

**Fakulti : Fakulti Sains dan Teknologi Makanan**

Kajian ini telah mengutarakan ciri-ciri probiotik strain *Bifidobacterium pseudocatenulatum* G4 dari segi profil keselamatan, kesan ke atas mikrobiota usus, interaksi dengan prebiotik, dan kebolehan untuk tumbuh serta hidup dalam media berasaskan susu skim kacang tanah. Profil keselamatan strain G4 dinilai dalam beberapa kumpulan tikus BALB/c yang di beri makan strain *Bifidobacterium* pada kepekatan yang berbeza selama 28 hari. Pada akhir eksperimen, di dapati pemberian *B. pseudocatenulatum* G4/hari tidak mempengaruhi kesihatan tikus mahupun memberi kesan toksik dalam darah secara biokimia atau ukuran haematologi. Tiada sebarang patogenik simptom di kesan dalam organ dalaman, enzim-enzim serum hati dan buah pinggang, dan histologi cecum, usus kecil dan kolon. Oleh itu, strain *B. pseudocatenulatum* G4 adalah mikroorganisma probiotik yang selamat dan mempunyai potensi untuk di guna dalam aplikasi makanan. Kesan memakan strain G4 keatas komuniti





mikrobiota terpilih dan aktiviti metabolik pada cecum dan kolon tikus telah diuji selama 30 hari. Kumpulan tikus tersebut menerima diet susu skim yang mengandungi *B. pseudocatenulatum* G4 hidup atau produk metabolik *B. pseudocatenulatum* G4. Kesan pemberian strain G4 yang hidup telah meningkatkan populasi Bifidobacteria, Lactobacillus dan keseluruhan bakteria anearob dalam cecum dan kolon tikus yang di uji, manakala bilangan keseluruhan bakteria aerob dan bakteria yang berpotensi sebagai patogen seperti staphylococcus, enterococcus dan enterobacteriaceae telah menurun. Bagaimanapun, hampir sedikit atau tiada langsung perubahan pada bilangan Salmonella and coliform. Pemberian strain G4 yang hidup juga telah meningkatkan kandungan asid lemak berantai pendek (acetic, propionic, dan butyric) secara signifikan ( $p > 0.05$ ) dalam cecum dan kolon tikus yang di uji. Komersial prebiotik terpilih [inulin dan fructo-oligosaccharide (FOS)] serta sorbitol, arabinan dan kepekatan inokulum telah diuji menggunakan fractional factorial design (FFD) untuk mengetahui kesan interaksi terhadap pertumbuhan *B. pseudocatenulatum* G4 dalam media berasaskan susu skim. Setelah 48 jam inkubasi, didapati sebahagian besar pertumbuhan bakteria dipengaruhi oleh fructo-oligosaccharide (FOS) dan kadar kepekatan inokulum. Penurunan pertumbuhan telah diperhatikan dalam semua sampel selepas 72 jam inkubasi hasil dari pembentukan asid laktik. Central composite design (CCD) telah digunakan ke atas FOS dan kadar inokulum pada 48 jam inkubasi pengeraman untuk menghasilkan model statistik bagi tujuan pengoptimaan. Model tersebut meramal 2.461 log CFU/ml sebagai peningkatan pertumbuhan optimum *Bifidobacterium pseudocatenulatum* G4. Kombinasi yang menghasilkan titik

optimum tersebut adalah 2.86 % FOS (g/v) and 0.67 % kadar inokulum (v/v). Model tersebut seterusnya disahkan dalam bioreaktor 1L untuk 24 jam. Keputusannya menunjukkan peningkatan pertumbuhan lebih tinggi iaitu 2.95 log CFU/ml jika dibandingkan dengan ramalan model optima iaitu 2.46 CFU/ml. Pertumbuhan strain G4 telah dinilai dalam produk berasaskan susu kacang tanah dan kadar pertumbuhan selepas 24 jam inkubasi adalah 7.12 log CFU/ml. Pertumbuhan ini telah dirangsang oleh kesan pemberian frocto-oligosacharide (FOS). Pertumbuhan yang tinggi ialah 8.35 log CFU/ml di dalam susu kacang tanah + FOS dan ianya setanding dengan 8.25 log CFU/ml dalam TPY yang di perolehi selepas 21 jam inkubasi. Strain G4 hidup dalam sampel terfermentasi yang disimpan pada 25°C selama 2 minggu telah menurun kepada paras < 7 log CFU/ml, yang mana ianya tidak menepati keperluan sebagai probiotik makanan. Bagaimanapun, selepas penyimpanan selama 2 minggu pada 4°C, pertumbuhan strain G4 kekal tinggi melangkaui paras yang disyorkan iaitu sepuluh juta sel per ml produk. Kesimpulannya, *B. pseudocatenulatum* G4 mempunyai potensi sebagai bakteria probiotik. Ianya adalah mikroorganisma probiotik yang selamat untuk tujuan formulasi synbiotik bersama FOS dan dicampurkan ke dalam produk berasaskan susu kacang tanah selain susu skim.

## ACKNOWLEDGEMENTS

First and foremost, I would like to express my sincere thanks and heartfelt gratitude to the members of my supervisory committee Prof Dr Mohd Yazid Abdul Manap (chairman), Assoc Prof Dr Shuhaimi Mustafa (member) and Assoc Prof Dr Mohd Nazrul Hakim (member) for their constructive guidance and motivation, from which I have acquired invaluable knowledge and research experiences. They have provided selfless efforts to add value to my research by reviewing several component manuscripts, which essentially formed the chapters of my thesis.

I would also wish to express my deepest appreciation to all lecturers and staff at the Faculty of Food Science and Technology for their kindness, friendliness, cooperation through out my study period. I am especially grateful to aunt naimah and sister nurul (Department of food services and managements) for their understanding, kind cooperation and help. Special thanks are also extended to Mr. halim, Mr. azman, and Mr. saib for their technical advices and kindness.

I am also grateful to Sudan University of Science and Technology for grant me one year scholarship during my study to a PhD level. Last but not the least; I would like to say a big THANK YOU to members of my family for their continuous support, encouragement, and patience during this study period. Lastly, I would like to thank all my dearest friends at Faculty of Food Science and Technology, particularly members of food biotechnology laboratory.



I certify that an Examination Committee has met on 30 March 2009 to conduct the final examination of Barka Mohammed Kabeir on his Doctor of Philosophy thesis entitled “Assessment of *Bifidobacterium pseudocatenulatum* G4 as a probiotic” in accordance with Universiti Putra Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends that the candidate be awarded the relevant degree. Member of the Examination Committee are as follows:

**Nazamid Saari, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Suhaila Mohamed, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Internal Examiner)

**Son Radu, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Internal Examiner)

**Wan Mokhtar Wan Yusof, PhD**

Professor  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(External Examiner)

---

**BUJANG KIM HUAT, PhD**  
Professor and Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 18 June 2009



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Mohd Yazid Abd Manap, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Shuhaimi Mustafa, PhD**

Associate Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)

**Muhammad Nazrul Hakim, PhD**

Professor  
Faculty of Medicine and Health Sciences  
Universiti Putra Malaysia  
(Member)

---

**HASANAH MOHD. GHAZALI, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 9 July 2009



## DECLARATION

I declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously and is not concurrently, submitted for any other degree at UPM or at any other institution.

---

**BARKA MOHAMMED KABEIR**

Date: 24 May 2009



## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	i
<b>ABSTRACT</b>	ii
<b>ABSTRAK</b>	v
<b>ACKNOWLEDGEMENTS</b>	viii
<b>APPROVAL</b>	ix
<b>DECLARATION</b>	xi
<b>LIST OF TABLES</b>	xvi
<b>LIST OF FIGURES</b>	xx
<b>LIST OF PLATES</b>	xxii
<b>LIST OF ABBREVIATIONS</b>	xxiii
<b>CHAPTERS</b>	
<b>1 INTRODUCTION</b>	<b>1</b>
<b>2 LITERATURE REVIEW</b>	<b>7</b>
2.1 Concept of beneficial microorganisms or probiotics	7
2.2 Definition of probiotics	8
2.3 Initial colonization of probiotics in gut ecology	10
2.4 Probiotics and intestinal microbiota relationships	13
2.5 Microbiota activities in the gut environment	18
2.6 Effects of imbalance in probiotics communities and related metabolic activities on host health	20
2.7 Beneficial effects of balanced probiotics community on the host well-being	21
2.8 Criteria for selection of a probiotic microorganism	22
2.9 Enhance the efficiency of probiotics and prebiotics	24
2.10 Utilization of peanut for human food	26
2.11 <i>Bifidobacterium pseudocatenulatum</i> G4 as a probiotic	29
2.12 Criteria for selection of prebiotics ingredients	35
2.13 Development of prebiotics ingredients	36
2.14 FOS as a candidate prebiotic	39
2.15 Experimental animals the stem to predict probiotics and prebiotics effect of on human health	41
<b>3 SAFETY ASSESSEMENT OF <i>Bifidobacterium</i> <i>pseudocatenulatum</i> in BALB/c mice</b>	<b>43</b>
3.1 Introduction	43
3.2 Materials and methods	45
3.2.1 Identification of <i>Bifidobacterium pseudocatenulatum</i> G4 using 16S rRNA-Gene- and species Targeted Primers	45



3.2.2	<i>Bifidobacterium</i> strains	46
3.2.3	Animals	46
3.2.4	General experimental protocol	48
3.2.5	Heamatology and blood biochemistry	50
3.2.6	Bacterial translocation	50
3.2.7	Internal organ indices	51
3.2.8	Histology observation of gut mucosa	51
3..2.9	Statistical analysis	52
3.3	Results and discussion	53
3.3.1	Conformation of <i>Bifidobacterium pseudocatenulatum</i> G4	53
3.3.2	Effect of feeding <i>Bifidobacterium pseudocatenulatum</i> G4 on general health of mice	53
3.3.3	Lack of toxicity in mice fed <i>Bifidobacterium</i> strains	63
3.3.4	Lack of pathogenicity symptoms in mice fed <i>Bifidobacterium</i> strains	71
3.4	Conclusion	74
<b>4</b>	<b>MICROBIOTA MODULATION IN CEACUM AND COLON OF RATS FED WITH <i>Bifidobacterium pseudocatenulatum</i> G4 SUPPLEMENTED MILK</b>	<b>77</b>
4.1	Introduction	77
4.2	Materials and methods	78
4.2.1	Animals	78
4.2.2	Experimental design	80
4.2.3	Internal organ and bowel weights	82
4.2.4	Clinical chemistry analysis of blood	82
4.2.5	Bacterial quantification of intestinal content	83
4.2.6	Analysis of total soluble solid (TSS), short chain fatty acids (SCFA) , and glucose	83
4.2.7	Statistical analysis	85
4.3	Results and discussion	86
4.3.1	Aspects on general health of rats	86
4.3.2	Safety profile of <i>Bifidobacterium</i> supplemented milk	88
4.3.3	Microbiota communities of ceacum and colon	96
4.3.4	Concentration of short chain fatty acids (SCFA)	101
4.3.5	Level of nutrients availability	105
4.4	Conclusion	108
<b>5</b>	<b>SYNBIOTIC GROWTH OPTIMIZATION OF <i>Bifidobacterium pseudocatenulatum</i> G4 WITH PREBIOTICS IN SKIM MILK BASED MEDIUM</b>	<b>110</b>
5.1	Introduction	110
5.2	Materials and methods	111
5.2.1	Maintenance of microorganism	111
5.2.2	Medium and fermentation conditions	112
5.2.3	Enumeration of viable bacterial cell	113





5.2.4	Determination of organic acids	113
5.2.5	Experimental design and statistical analysis	114
5.3	Results and discussion	119
5.3.1	Screening for significant growth factors	119
5.3.2	Approaching the optimum growth point	125
5.3.3	Optimal range of significant growth factors	125
5.3.4	Validating the statistical model of optimum growth	132
5.3.5	Organic acids profiles produced during the screening for significant growth factors	134
5.3.6	Organic acids profiles at point of optimum synbiotics growth	148
5.3	Conclusion	152
<b>6</b>	<b>THE GROWTH OF <i>Bifidobacterium pseudocatenulatum</i> G4 IN PEANUT MILK (PM) SUPPLEMENTED WITH FRUCTOOLIGOSACCHARIDES (FOS)</b>	<b>154</b>
6.1	Introduction	154
6.2	Materials and methods	157
6.2.1	Preparation of fermentation mediums	157
6.2.2	Starter culture and <i>Bifidobacterium</i> inoculums	157
6.2.3	Batch cultivation conditions	160
6.2.4	Shelf life study of fermented products	160
6.2.5	Microbiological analysis	162
6.2.6	Chemical analysis	162
6.2.7	Statistical analysis	163
6.3	Results and discussion	163
6.3.1	Growth of <i>B. pseudocatenulatum</i> G4 during incubation of peanut milk (PM) and skim milk (SM)	163
6.3.2	The change of pH and organic acid profiles during incubation of PM and SM with <i>Bifidobacterium</i> <i>pseudocatenulatum</i> G4	169
6.3.3	Survivability of <i>Bifidobacterium pseudocatenulatum</i> G4 during the storage of fermented PM and SM	178
6.3.4	pH decline and acid accumulation during the storage of PM and SM fermented with <i>Bifidobacterium pseudocatenulatum</i> G4	181
6.4	Conclusion	189
<b>7</b>	<b>GENERAL CONCLUSION AND RECOMMENDATIONS</b>	<b>190</b>
	<b>REFERENCES</b>	<b>193</b>
	<b>APPENDICES</b>	<b>211</b>
	<b>BIODATA OF THE STUDENT</b>	<b>221</b>
	<b>LIST OF PUBLICATIONS</b>	<b>222</b>



## LIST OF TABLES

Table		Page
2.1	Composition of 100 grams row peanut without skin	27
3.1	Two-Way ANOVA result of feed intake of mice fed different <i>Bifidobacterium</i> doses for four treatment weeks and two weeks acclimatization period	59
3.2	Two-Way ANOVA result of water intake of mice fed different <i>Bifidobacterium</i> doses for four treatment weeks and two weeks acclimatization period	60
3.3	Haematology parameters of mice groups orally fed different doses of <i>Bifidobacterium</i> per day for a period of four weeks	64
3.4	Blood biochemistry data of mice groups orally fed different doses of <i>Bifidobacterium</i> per day for a period of four weeks	66
3.5	Marker enzymes profiles of mice groups orally fed with different doses of <i>Bifidobacterium</i> per day for a period of four weeks	67
3.6	Internal organs (Liver, Heart, Kidney, and Spleen) indices of mice fed on different doses of <i>Bifidobacterium</i> strains per day	68
3.7	Summary of the incidence of bifidobacterial translocation in mice <sup>a</sup> blood	72
3.8	Histology measurements ( $\mu\text{m}$ ) of mice fed with different doses of <i>Bifidobacterium</i> per day for a period of four weeks	75
4.1	Enumeration media and incubation conditions of different microbiota communities in ceacum and colon of rats fed different milk supplements	84
4.2	Blood biochemistry profile of rats fed different <i>Bifidobacterium</i> milk supplements for a period of 20 days	90



4.3	Serum enzyme of blood in rats fed different <i>Bifidobacterium</i> milk supplements for a period of 20 days	91
4.4	Haematology parameters of blood in rats fed different <i>Bifidobacterium</i> milk supplements for a period of 20 days	93
4.5	Indices of internal organs and bowel weight of rats fed with different <i>Bifidobacterium</i> milk supplements	95
4.6	Total aerobe and potential helpful population (log CFU/ml) from ceacum and colon of rats received different <i>Bifidobacterium</i> supplement	97
4.7	Potential pathogens (Log CFU/g) from ceacum and colon of rats received different <i>Bifidobacterium</i> supplements	99
4.8	Short chain fatty acids (mmol/ml) in ceacum and colon of rats received different <i>Bifidobacterium</i> supplements for a period of 20 days	102
4.9	Total soluble solid (TSS) in ceacum and colon of rats received different <i>Bifidobacterium</i> supplements for a period of 20 days	106
4.10	Glucose content (mg/g sample) in ceacum and colon of rats received different <i>Bifidobacterium</i> supplements for a period of 20 days	107
5.1	Actual factors levels corresponding to coded factor levels	115
5.2	Experimental design of the $2^{5-2}$ fractional factorial design (FFD)	116
5.3	Central composite design (CCD) using coded levels of FOS and inoculum rate for samples incubated for two days	118
5.4	Results of strain G4 growth (log CFU/ml) in skim milk from the second- order model of the $2^{5-2}$ fractional factorial design (FFD)	120
5.5	ANOVA results of the second order model for fractional factorial design after 48h and 72h incubation periods	121



5.6	Results of $2^{5-2}$ fractional factorial design (FFD) regression analysis coefficient estimate for strain G4 growth and reduction after incubation at 72h	123
5.7	Steepest path scent for FOS and inoculum rate	126
5.8	Results of strain G4 growth (log CFU/ml) in skim milk from the central composite design (CCD) using coded levels of FOS and inoculum rate for samples incubated for 48h	127
5.9	Analysis of Variance for CCD results of FOS and inoculum rate for samples incubated for 48h	128
5.10	Estimated Regression Coefficients for central composite design (CCD) using coded levels of FOS and inoculum rate for samples incubated for 48h	130
5.11	Growth of Bifidobacterium G4 (log CFU/ml) in skim milk medium fermented in 1L bench bioreactor	133
5.12	Results of lactic acid (mmol/ml) from second- order model of the $2^{5-2}$ fractional factorial design (FFD)	135
5.13	Results of acetic acid (mmol/ml) from second- order model of the $2^{5-2}$ fractional factorial design (FFD)	136
5.14	Results of propionic acid (mmol/ml) from second- order model of the $2^{5-2}$ fractional factorial design (FFD)	137
5.15	Results of butyric acid(mmol/ml) from second- order model of the $2^{5-2}$ fractional factorial design (FFD)	138
5.16	Results of $2^{5-2}$ fractional factorial design (FFD) regression analysis coefficient estimate for lactic acid production at different incubation periods	140
5.17	Results of $2^{5-2}$ fractional factorial design (FFD) regression analysis coefficient estimate for acetic acid production at different incubation periods	142
5.18	Results of $2^{5-2}$ fractional factorial design (FFD) regression analysis coefficient estimate for propionic acid production at different incubation periods	144
5.19	Results of $2^{5-2}$ fractional factorial design (FFD) regression analysis coefficient estimate for butyric acid	146



	production at different incubation periods	
5.20	Results of organic acids (mmol/ml) from central composite design (CCD) using coded levels of FOS and inoculum rate for samples incubated for two days	149
5.21	Analysis of Variance for organic acids from CCD results of FOS and inoculum rate for samples incubated for two days	150
5.22	Estimated Regression Coefficients of organic acids from central composite design (CCD) using coded levels of FOS and inoculums rate for samples incubated for two days	151
6.1	Kinetic parameter values from batch fermentation of peanut milk and skim milk with <i>B. pseudocatenulatum</i> G4 in 1-L bioreactor	166
6.2	Survival of <i>B. pseudocatenulatum</i> G4 (log CFU/ml) during the storage of fermented peanut and skim milk with different supplements	179
6.3	pH decline during the storage of fermented peanut and skim milk with <i>Bifidobacterium pseudocatenulatum</i> G4	182
6.4	Lactic acid profile (mmol/ml) during the storage of peanut and skim milk fermented with <i>Bifidobacterium pseudocatenulatum</i> G4	184
6.5	Acetic acid profile (mmol/ml) during the storage of peanut and skim milk fermented with <i>Bifidobacterium pseudocatenulatum</i> G4	186
6.6	Probionic acid profile (mmol/ml) during the storage of peanut and skim milk fermented with <i>B. pseudocatenulatum</i> G4	187
6.7	Butyric acid profile (mmol/ml) during the storage of fermented peanut and skim milk with <i>B. pseudocatenulatum</i> G4	188



## LIST OF FIGURES

Figure		Page
2.1	The intestinal microbiota development during the first seven days of child	11
2.2	Changes in fecal microbiota communities with increased age	14
2.3	Microbiota communities in different regions of the gastrointestinal tract	15
2.4	Criteria suggested for a successful probiotic microorganism	23
3.1	Feed intake (g/mouse) by mice groups fed on different <i>Bifidobacterium</i> doses for a period of four weeks in addition to prior two weeks acclimatization period	57
3.2	Water intake (ml/mouse) by mice groups fed on different <i>Bifidobacterium</i> doses for four weeks in addition to prior two weeks acclimatization period	58
3.3	weight of mice (g) fed on different <i>Bifidobacterium</i> doses for four weeks in addition to prior two weeks acclimatization period	62
4.1	Weight of rats (g) fed different <i>Bifidobacterium</i> supplements for 30 days recorded at ten days interval throughout the experimental period	87
5.1	Contour plot of strain G4 growth (log CFU/ml) fitted to response surface model	131
6.1	Growth of <i>B. pseudocatenulatum</i> G4 (log CFU/ml) during fermentation of peanut milk (PM) and skim milk (SM) with different supplements	165
6.2	pH decline during fermentation of peanut milk and skim milk with <i>Bifidobacterium pseudocatenulatum</i> G4	170
6.3	Lactic acid (mmol/ml) production by <i>Bifidobacterium pseudocatenulatum</i> G4 during fermentation of peanut milk and skim milk with different supplements	172



6.4	Acetic acid (mmol/ml) production by <i>Bifidobacterium pseudocatenulatum</i> G4 during fermentation of peanut and skim milk with different supplements	174
6.5	Propionic acid (mmol/ml) production by <i>B. pseudocatenulatum</i> G4 during fermentation of peanut and skim milk with different supplements	176
6.6	Butyric acid (mmol/ml) production by <i>Bifidobacterium pseudocatenulatum</i> G4 during fermentation of peanut and skim milk with different supplements	177



## LIST OF PLATES

Plate		Page
2.1	Morphology of <i>Bifidobacterium</i> probiotic microorganism	31
3.1	BALB/c mice	47
3.2	Oral feeding of mice	49
3.3	Gel electrophoresis of PCR products using genus-specific primer Lm26/Lm3	54
3.4	Gel electrophoresis of PCR products using species-specific primers BiCATG-1/ BiCATG-2	55
4.1	Female Sprague-Dawley rats housed two per cage	79
4.2	Taking weight of the rats	81
6.1	Peanut without skin	158
6.2	Fermentation mediums prior sterilization	159
6.3	Batch fermentation process with a four vessels Bioreactor	161





## LIST OF ABBREVIATIONS

G	gram
°C	degree Celsius
%	Percentage
et al.	et cetera (and company)
WHO	World Health Organization
PCR	Polymerase chain reaction
16S rRNA	Ribosomal ribonucleic acid
Kb	kilo base pair
bp	base pair
JCM	Japan Collection of Microorganism
MRS	de Mann-Rogosa-Sharpe
nm	nanometer
OD	Optical density
PBS	phosphate buffer saline
CFU	colony Forming Unit
mmol	milimolar
ml	milliliter
x g	gravity
EDTA	Ethylenediaminetetraacetic acid
H&E	Haematoxylin and Eosin
mg	milligram
ANOVA	Analysis of variance

