



UNIVERSITI PUTRA MALAYSIA

**PRODUCTION, CHARACTERIZATION AND EXPRESSION OF AN
ORGANIC SOLVENT TOLERANT LIPASE FROM PSEUDOMONAS
AERUGINOSA S5**

SYARUL NATAQAIN BAHARUM.

FBSB 2005 4

**PRODUCTION, CHARACTERIZATION AND EXPRESSION OF AN ORGANIC
SOLVENT TOLERANT LIPASE FROM *PSEUDOMONAS AERUGINOSA* S5**

By

SYARUL NATAQAIN BAHARUM

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the degree of Doctor of Philosophy**

November 2005



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

PRODUCTION, CHARACTERIZATION AND EXPRESSION OF AN ORGANIC SOLVENT TOLERANT LIPASE FROM *PSEUDOMONAS AERUGINOSA* S5

By

SYARUL NATAQAIN BINTI BAHARUM

November 2005

Chairman : Professor Abu Bakar Salleh, PhD

Faculty : Biotechnology and Biomolecular Sciences

Lipolytic bacterium was screened from five pure bacteria cultures available in Enzyme and Microbial Technology laboratory in UPM. The stock cultures were tested for lipase production. Two isolates (S5 and 205W) showed the highest activity in tripticase soy broth and brain heart infusions. These isolates were further incubated in different basal media. Isolate S5 was shown to give higher activity (0.327 U/ml) than isolate 205W in media M1 and stable in various organic solvents tested. Therefore isolate S5 was chosen for further studies. Based on its morphological, biochemical characteristics and 16S rDNA sequence, strain S5 was identified as *Pseudomonas aeruginosa*. *P. aeruginosa* lipase exhibited the highest relative activity with n-hexane (410%) for 20 min reaction. Optimum lipase production was obtained at pH 7.0 and 37°C at static condition with peptone as the best nitrogen source and olive oil as the best carbon source. The best inoculum size was 6%. The surfactants, Tween 60 and Tween 80 were found to enhance for bacterial growth and lipase production by S5.



The lipase was purified to homogeneity by affinity column chromatography and anion exchange column chromatography. The purified lipase was highly homogeneous as determined by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and the molecular mass was estimated to be 60 kDa by SDS-PAGE and G-100 gel filtration column chromatography. The optimum temperature and pH of the purified enzyme was 45°C and pH 9.0, respectively. S5 lipase was stable at pH 6-9 for 30 min. The half-life of the S5 lipase at 45°C and 50°C was 2 h and 1 h, respectively. The lipase exhibited high stability in the presence of n-dodecane, 1-pentanol and toluene. As for metal ions, it was found that Ca²⁺ stimulated lipase activity in 15 min incubation time, while EDTA had no effect on lipase activity. However, the S5 lipase was strongly inhibited by the addition of 1 mM phenyl methyl sulfonyl fluoride (PMSF) (87% inhibition) and 1 mM of Pepstatin (76% inhibition) after 30 min incubation. The S5 lipase exhibited the highest activity in the presence of palm oil as a substrate and followed by coconut oil. S5 lipase was found to have the highest activity against triolein which possess longer carbon chain length. S5 lipase is a non-specific lipase as shown by triolein hydrolysis.

The gene encoding for the intracellular lipase of *P. aeruginosa* strain S5 was isolated via genomic DNA library and cloned into *pRSET*. The cloned sequence included two open reading frames (ORF) consisting of 1575 bp for the first ORF (ORF1) and 582 bp for the second ORF (ORF2). The ORF2 was located at the downstream and function as the *act* gene for ORF1. The conserved pentapeptide Gly- X- Ser- X-Gly was located in the ORF1. Catalytic triad resembling of that serine protease, consisting of serine, histidine,



aspartic acid or glutamic acid residues was present in this lipase gene. Expression in *E.coli* resulted a 100-fold increase in enzyme activity after 9 h induction with 0.75 mM IPTG. The recombinant plasmid revealed a size of 60 kDa on SDS-PAGE. The Lip S5 gene was stable in the presence of 25% (v/v) n-dodecane and n-tetradecane after 2 h incubation at 37°C. Predicted 3D structure of S5 lipase revealed topological organization of α / β -hydrolase fold consisting of 10 α -Helices and 5 β -strands. The Ramachandran plot of S5 lipase showed that 85.8% (229) of residues lie in the most-favored region and only 2.2% (6) of residue lie in generously allowed regions and 1 residue lie in disallowed region.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENGHASILAN, PENCIRIAN, PENGKLONAN DAN PENGEKSPRESAN
LIPASE DARIPADA *PSEUDOMONAS AERUGINOSA* S5 YANG TOLERAN
TERHADAP PELARUT ORGANIK**

Oleh

SYARUL NATAQAIN BINTI BAHARUM

November 2005

Pengerusi : Profesor Abu Bakar Salleh, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Bakteria yang menghasilkan enzim lipolitik telah disaring daripada lima kultur bakteria tulen yang terdapat di Makmal Enzim dan Teknologi Mikrob, UPM. Stok kultur ini telah diuji untuk penghasilan lipase. Dua pencilan (S5 dan 205w) menunjukkan aktiviti yang tertinggi di dalam media soya triptikas dan infusi hati dan otak. Kedua-dua pencilan ini seterusnya dieram di dalam basal media berlainan. Pencilan S5 dipilih untuk kajian selanjutnya kerana ia menunjukkan aktiviti yang tinggi (0.327 U/ml) berbanding pencilan 205w di dalam media M1 dan lebih stabil di dalam pelbagai pelarut organik yang telah diuji. Berdasarkan morfologi, pencirian biokimia dan jujukan 16S rDNA, strain S5 ini dikenalpasti sebagai *Pseudomonas aeruginosa*. Lipase *P. aeruginosa* menunjukkan aktiviti relatif tertinggi di dalam n-hexane (410%) di dalam tindak balas selama 20 minit. Kajian pengoptimuman terhadap penghasilan lipase oleh pencilan S5 menunjukkan penghasilan lipase adalah maksimum pada pH 7.0 dan 37°C pada keadaan statik dengan pepton sebagai sumber nitrogen terbaik dan minyak zaitun sebagai sumber

karbon terbaik. Saiz inokulum yang terbaik ialah 6%. Tween 60 dan Tween 80 merupakan sumber surfaktan terbaik bagi pertumbuhan bakteria dan penghasilan lipase bagi S5

Lipase ini telah ditulenkan sehingga homogeniti yang nyata menggunakan kromatografi turus afiniti dan penukaran cas anion. Lipase yang telah ditulenkan adalah berhomogenus tinggi berdasarkan poliakrilamid elektroforesis gel sodium dodecil sulfat (SDS-PAGE). Berat molekul lipase ini adalah 60 kDa berdasarkan SDS-PAGE dan gel penurasan kromatografi turus G-100. Suhu dan pH optimum bagi enzim tulen ini adalah 45°C dan pH 9.0, masing-masing. Lipase S5 adalah stabil pada pH 6-9 selama 30 minit. Separuh hayat bagi lipase S5 adalah 45°C dan 50°C masing-masing selama dua dan satu jam. Lipase ini menunjukkan kestabilan yang tinggi di dalam n-dodekana, 1-pentanol dan toluene. Ca^{2+} merangsang lipase aktiviti dalam masa 15 min pengeraman, sementara EDTA tidak memberi kesan terhadap aktiviti lipase. Walau bagaimanapun, lipase S5 direncat dengan penambahan 1mM fenil metal sulfonil flourida (PMSF) (87% direncat) dan 1mM Pepstatin (76% direncat) setelah dieram selama 30 minit. Lipase S5 menunjukkan aktiviti yang tinggi dengan kehadiran minyak kelapa sawit dan minyak kelapa sebagai substrat. Lipase S5 didapati menunjukkan aktiviti yang tinggi terhadap triolein yang mempunyai rangkaian karbon yang panjang. Lipase S5 adalah jenis tidak spesifik seperti ditunjukkan ke atas hidrolisis triolein.

Gen yang mengkodkan intrasel lipase dari *P. aeruginosa* S5 dipencilkan menerusi perpustakaan genomik DNA dan diklonkan ke dalam *pRSET*. Klon ini mengandungi dua rangka bacaan terbuka (ORF) yang mengandungi 1575 bp untuk ORF pertama (ORF1)

dan 582 bp untuk ORF kedua (ORF2). ORF2 ini yang terletak di aliran ke bawah yang berperanan sebagai gen bertindak (*act gene*) untuk ORF1. Pentapeptida terpelihara Gly-X- Ser- X- Gly terletak di ORF1. Triad pemangkin yang seperti didapati dalam protease serina yang mengandungi serina, histidina, asid aspartik atau asid glutamik terdapat di dalam gen lipase ini. Ekspresi di dalam *E.coli* menghasilkan peningkatan aktiviti enzim sebanyak 100 kali ganda selepas 9 jam induksi dengan 0.75 mM IPTG. Plasmid rekombinan di atas SDS-PAGE menunjukkan saiz 60 kDa. Lip S5 adalah stabil dengan kehadiran 25% (v/v) n-dodekana dan n-tetradekana selepas 2 jam pada 37°C. Ramalan struktur tiga dimensi (3D) S5 lipase menunjukkan organisasi topologi lipatan α/β -hidrolase mengandungi 10 α -Heliks and 5 β -bebenang. Plot Ramachandran menunjukkan 85.8% (229) residu terletak di dalam kawasan yang diyakini betul, sementara 11.6% (31) terletak di dalam kawasan yang betul. Hanya 2.2% (6) residu terletak di dalam kawasan yang diragui betul dan satu residu terletak di dalam kawasan yang tidak betul.



ACKNOWLEDGEMENTS

All praises be to the Al-mighty Allah, the Merciful and the Beneficent for the strength and blessing in the completion of this challenging tasks.

My heartiest gratitude goes to the Chairman of my supervisory committee: Prof. Dr. Abu Bakar Salleh for his invaluable advice, guidance, ideas and comments, in the midst of his heavy responsibilities and duties. My deep appreciation is also extended to Assoc. Prof. Dr. Raja Noor Zaliha Raja Abd. Rahman for her guidance, encouragement, support and motherly care. I am also grateful to Prof. Dr. Mahiran Basri for her indispensable advice, suggestion and encouragement throughout the project. In taking this text from the raw manuscript stage to the final ready thesis stage, I have received tremendous assistance from the meticulous checking of the text to numerous suggestions for modifications and special learning aids from all of you.

I would like to dedicate my appreciation to Assoc. Prof. Dr. Mohd. Basyaruddin Abd. Rahman and Dr. Norazizah Shafie for their constructive comments, constant support and invaluable guidance. My former supervisor Assoc. Prof. Dr. Che Nyonya Abd. Razak, thank you for giving me the opportunity to study on this interesting and advent project. I really miss you.

Special thanks are also extended to all my labmates in Lab 139 and 140: Ina, Aiman, Sha, Fairol, Rofandi, Zira, Bimo, Ayob, Brother Mohamad, Sue, Ada, Shukuri, Tengku and K. Lia for their assistant and friendship. Most of all their presence has made my



I certify that an Examination Committee met on 11th November 2005 to conduct the final examination of Syarul Nataqain Baharum on her Doctor of Philosophy thesis entitled "Production, Characterization and Expression of an Organic Solvent Tolerant Lipase from *Pseudomonas aeruginosa* S5" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Lai Oi Ming, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Norhani Abdullah, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Foo Hooi Ling, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Ibrahim Che Omar, PhD

Professor
School of Biological Sciences
Universiti Sains Malaysia
(External Examiner)



HASANAH MOHD. GHAZALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **19 JAN 2006**

This thesis submitted to the Senate of the Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as followed:

ABU BAKAR SALLEH, PhD


Professor
Faculty of Botechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

RAJA NOOR ZALIHA RAJA ABDUL RAHMAN, PhD

Associate Professor
Faculty of Botechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

MAHIRAN BASRI, PhD

Professor
Faculty Science
Universiti Putra Malaysia
(Member)



AINI IDERIS, PhD

Professor/ Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **07 FEB 2006**



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



SYARUL NATAQAIN BAHARUM

Date: 10/01/06

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xxii
LIST OF FIGURES	xxiv
LIST OF ABBREVIATIONS	xxvii
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	5
2.1 Lipase	5
2.2 Sources of Lipase	6
2.3 Classification of Microbial Lipolytic Enzymes	7
2.4 Applications of Lipase	10
2.5 Organic Solvent tolerant Microorganisms	11
2.6 Enzyme in Organic Solvents	12
2.7 Organic Solvent Tolerant Enzymes: Prospects and Challenges	17
2.8 Enzyme in Non-conventional Media	19
2.9 Effect of Nutritional Factor on Lipase Production	20
2.9.1 Nitrogen Sources	20
2.9.2 Carbons	21
2.9.3 Surfactants	23
2.9.4 Lipids	24
2.9.5 Metal Ions	25
2.10 Effect Of Physical Factor on Lipase Production	25
2.10.1 Effect of the Cultivation Temperature	26
2.10.2 Effect of the Initial pH of the Medium	27
2.10.3 Effect of Inoculum Size	27
2.10.4 Effect of Shaking Condition	28
2.10.5 Time Course Study	29
2.11 Purification of Lipase	29
2.11.1 Initial Step of purification	30
2.11.2 Chromatography Separation	30
2.11.3 Purification of Microbial Lipases	32
2.12 Characterization of Lipase	36
2.12.1 Optimum Temperature	36
2.12.2 Optimum pH	37
2.12.3 Effect of Metal Ions	38
2.12.4 Positional Specificity	40



2.13	Molecular Cloning	41
2.13.1	Cloning of the Lipase Gene	42
2.13.2	Genomic DNA Library	44
2.13.3	Size of Library	45
2.13.4	Selection of Vector	46
2.13.5	Selection of Host	47
2.13.6	Expression of Lipase Gene	48
2.14	Protein Structure Prediction	49
2.14.1	Comparative Protein Structure Modeling	50
3	MATERIALS AND METHODS	55
3.1	Materials	55
3.2	Equipments	60
3.3	Methodology	61
3.3.1	Sources of Bacteria	61
3.3.2	Screening of Lipase Producer	61
3.3.3	Preparation of Inoculum	62
3.3.4	Preparation of Stock Culture	63
3.3.5	Assay of Lipase Activity	63
3.3.6	Statistical Analysis	64
3.3.7	Stability of Lipase in Organic Solvents	64
3.3.8	Effect of Reaction Time on Enzyme activity in n-Hexane	65
3.3.9	Growth Curve and Lipase Production of <i>Pseudomonas aeruginosa</i> Strain S5	65
3.3.10	Bacterial Identification	65
3.3.11	Optimization of the Lipase Production	67
3.3.12	Purification of Lipase	72
3.3.13	Characterization of Organic solvent Tolerant Lipase	75
3.3.14	Cloning and Sequencing of the Organic Solvent Tolerant Lipase Gene	80
3.3.15	PCR Cloning	81
3.3.16	Genomic DNA Library	84
3.3.17	Structure Prediction of an Organic Solvent Tolerant Lipase S5	90
4	RESULTS AND DISCUSSION	92
4.1	Screening of the Lipase Producer	92
4.2	Screening of Organic Solvent Tolerant Lipase Produced by Selected Lipase Producer	95
4.3	Stability of S5 Lipase in n-Hexane	97
4.4	Bacterial Identification	99
4.4.1	Morphology and Biochemical Test	99
4.4.2	16S rDNA Identification and Phylogenetic Tree Analysis	101
4.4.3	Time Course Studies on Bacterial Growth and Lipase Activity	107



4.5	Optimization Studies	109
4.5.1	Effect of Physical Factor on Growth and Lipase Production	109
4.5.2	Effect of Nutritional Factor on Growth and Lipase Production	120
4.6	Purification of S5 Lipase	144
4.6.1	Affinity Column Chromatography	145
4.6.2	Anion Exchange Column Chromatography	147
4.7	Characterization of Purified Lipase	153
4.7.1	Determination of molecular Mass	153
4.7.2	Stability of the Purified Enzyme in Organic Solvent	156
4.7.3	Effect of pH on Purified Lipase S5	160
4.7.4	Effect of pH on Enzyme Stability	163
4.7.5	Effect of Temperature	167
4.7.6	Effect of Temperature on Enzyme Stability	170
4.7.7	Effect of Metal Ions, Oxidizing, Reducing and Chelating Agents	172
4.7.8	Substrate Specificity	176
4.7.9	Positional Specificity against Triacylglycerols	178
4.8	Cloning and Expression of an Organic Solvent Tolerant Lipase S5	180
4.8.1	Genomic DNA Extraction	180
4.8.2	Lipase Gene Isolation via PCR	181
4.8.3	Cloning of Organic Solvent Tolerant Lipase Gene by p <i>Drive</i> Cloning Vector	189
4.8.4	Cloning and Expression of S5 Lipase by Different Expression Vectors	193
4.8.5	Lipase Gene Isolation via Genomic DNA Library	197
4.8.6	Restriction Mapping of Putative Lipase Gene	204
4.8.7	Gene Analysis	207
	Nucleotide Sequence Analysis	207
	Phylogenetic Analysis of S5 Lipase	213
	Signal Peptide Prediction and Hydrophobicity Profile of S5 Lipase	215
	Amino acid Composition	217
4.8.8	Optimization and Expression of <i>P. aeruginosa</i> Lipase in <i>E. coli</i>	217
	Effect of IPTG Concentration	222
	Induction Time	224
4.8.9	Stability of Recombinant Clone in Various Organic Solvents	228
4.9	Structural Prediction of S5 Lipase	230
4.9.1	Model Building	230
4.9.2	Structural Evaluation of the S5 Lipase	236
5	CONCLUSION AND RECOMMENDATIONS	238
5.1	Conclusion	238
5.2	Recommendations	242



6	REFERENCES	245
	APPENDICES	263
	BIODATA OF THE AUTHOR	



LIST OF TABLES

Tables	Page
1. The Classification of Microbial Lipolytic Enzymes	9
2. Log $P_{o/w}$ Value of Comon Solvents Used	14
3. General Effects of Organic Solvents on Biocatalysis	18
4. Properties of Cloned Lipase Genes	43
5. Different Media for Lipase Production	62
6. The Dilutions of <i>MBOI</i> in Ten Individual Microcentrifuge Tubes	86
7. Lipase Production by Different Isolates in Enrichments Media	93
8. Lipase Production by Isolate 205w and S5 in Different Production Media	94
9. Effect of Organic Solvents on S5 and 205w Extracellular Lipases	96
10. Taxonomic Characteristics of the <i>Pseudomonas</i> sp. strain S5	100
11. Summary of Purification of S5 Lipase	151
12. Stability of Purified Lipase In The Presence of Various Organic Solvents	157
13. Effect of Various Reagents on Lipase Activity	174
14. Hydrolytic Activities of S5 Lipase against Various Natural Oils and Triacylglycerols	177
15. Spectrophotometric Assay of the Extracted Genomic DNA	183
16. Lipase Activities Produced by the Recombinant Clones	196
17. Lipase Activity of Different Colonies of S5/p <i>RSET</i> after 6 hour Induction with 1 mM IPTG	203
18. Percentage of Amino Acid Composition Between S5 Lipase and other Lipases from Various Organisms	221



19. Lipase activity of <i>S5/pRSET</i> after 6 hours Induction at Different Concentration of IPTG	223
20. LipS5 Lipase Stability in Various Organic Solvents	229



LIST OF FIGURES

Figures	Page
1. Flowchart for Comparative Protein Structure Modeling on the Genome Scale	52
2. Stability of S5 Lipase in n-Hexane	98
3. 16S rDNA Gene (1500 bp) of Isolate S5 Gene Amplified via PCR	102
4. 16S rDNA Sequence of <i>Pseudomonas aeruginosa</i> Strain S5	103
5. Rooted Phylogenetic Tree Showing the Relationship of Isolate S5 to other <i>Pseudomonas aeruginosa</i>	105
6. Growth Curve and Lipase Production of <i>Pseudomonas aeruginosa</i> strain S5	108
7. Effect of Agitation Rates on Lipase Production and Bacterial Growth	111
8. Effect of Inoculum Sizes on Lipase Production and Bacterial Growth	113
9. Effect of pH on Lipase Production and Bacterial Growth	115
10. Effect of Temperature on Lipase Production Bacterial Growth	118
11. Effect of Organic Nitrogen Sources on Lipase Production	122
12. Effect of Organic Nitrogen Sources on Bacterial Growth	123
13. Effect of Inorganic Nitrogen Sources on Lipase Production	125
14. Effect of Inorganic Nitrogen Sources on Bacterial Growth	126
15. Effect of Carbon Sources on Lipase Production	128
16. Effect of Carbon Sources on Bacterial Growth	130
17. Effect of Lipids on Lipase Production	132
18. Effect of Lipids on Bacterial Growth	135



19. Effect of Surfactants on Lipase Production	137
20. Effect of Surfactants on Bacterial Growth	139
21. Effect of Metal Ions on Lipase Production	140
22. Effect of Metal Ions on Bacterial Growth	143
23. Affinity Column Chromatography Elution Profile of the S5 Lipase on Con A Sepharose	146
24. Electrophoresis of strain S5 Lipase in 10% Polyacrylamide Gel Under Denaturing Conditions	148
25. Anion Exchange Elution Profile of the S5 Lipase on DEAE Sephacel	150
26. S5 Lipase Molecular Weight Estimated by Gel Filtration G-100	154
27. Electrophoresis of strain S5 Lipase in 10% Polyacrylamide Gel Under Denaturing Conditions	155
28. Effect of pH on Purified S5 Lipase Activity	161
29. Effect of pH on Purified S5 Lipase Stability	164
30. Effect of Temperature on Purified S5 Lipase Activity	168
31. Effect of Temperature on Purified S5 Lipase Stability	171
32. Thin-layer Chromatogram of the Hydrolysis Products of Triolein Catalyzed by S5 Lipase	179
33. Extraction of Genomic DNA from <i>P. aeruginosa</i> S5	182
34. Gel Electrophoresis of PCR Products Amplified by using Primers For F1 and Rev R1	185
35. Alignment of Nucleotide Sequences between LipS5 (964 bp) and LipC PAO1 (900 bp)	187
36. Gel Electrophoresis of PCR Products Amplified by using Primers For PGL and Rev PGL	188
37. Gel Electrophoresis of Recombinant Plasmid Extractions	190



38. Gel Electrophoresis of Recombinant Plasmids that was amplified by using Primers For PGL and Rev PGL	191
39. Nucleotide Sequence of the Lipase Gene from <i>P. aeruginosa</i> S5	192
40. Gel Electrophoresis of S5/pTrcHis/Top10 Recombinant Plasmid Extractions	194
41. Gel Electrophoresis of S5/pTrcHis/Top10 Recombinant Plasmid Digested with <i>EcoRI</i>	195
42. MBOI Partial Digestion of Genomic DNA Extracted from <i>P. aeruginosa</i> S5	198
43. Recombinant Clone of S5/pRSET / Top 10 on tributyrin-ampicilin LB agar plate	200
44. Recombinant Clone of S5/pRSET / Top 10 on Victoria blue-ampicilin LB agar plate	201
45. Recombinant clone of S5/pRSET / Top 10 on Rhodamine B-ampicilin LB agar plate	202
46. Plasmid Digestion of Recombinant Clones	205
47. Restriction Mapping Scheme of the DNA coding <i>P. aeruginosa</i> S5 Lipase from LipS5	206
48. Nucleotide Sequence of the Lipase Gene from <i>P. aeruginosa</i> S5	210
49. Comparison of the Amino Acid sequence of Active Site among Lipases	212
50. The Phylogenetic Tree Analysis of S5 Lipase Gene	214
51. The SignalP Neural Network Output of Organic Solvent Tolerant S5 Lipase from <i>Pseudomonas aeruginosa</i> Strain S5	216
52. Hydrophobicity of S5 lipase according to Kyte and Doolittle (1982)	218
53. Lipase activity of Top 10/S5/pRSET at Different Time Intervals after Induction with 0.75mM IPTG	225
54. Different Time Induction with 0.75 mM IPTG	227



55. Amino acid Sequence Alignment of S5 lipase (S5) and <i>Pseudomonas aeruginosa</i> PAO1 (IEX9) as a Template with 46% Sequence Identity	232
56. The predicted Structure of S5 Lipase Rendered as Ribbon	234
57. Secondary Structure of S5 Lipase as generated by DSSP	235
58. Ramachandran Plot of Predicted S5 Lipase Structure	237



LIST OF ABBREVIATIONS

APS	ammonium persulfate
BHI	Brain heart infusion
bp	base pair
BTEX	Benzene, Toluene, Ethyl-benzene, p-xylene
cm	centimeter
CM	carboxymethyl
CMC	critical micellar concentration
ConA	Concanavalin A
Da	Dalton
DEAE	diethylaminoethyl
DMSO	dimethyl sulfoxide
DMF	n-n-dimethylformamide
DNA	deoxyribonucleic acid
DTT	dithiothreitol
EDTA	ethylenediaminetetraacetic acid
FFA	free fatty acid
FPLC	fast protein liquid chromatography
g	gram
g/L	gram per liter
h	hour
IPTG	isopropyl β -D thiogalactoside
Kb	kilobase
kDa	kiloDalton



L	liter
LB/amp	Luria-Bertani containing ampicilin
M	molar
mA	milliampear
mM	millimolar
mg	milligram
min	minute
NB	nutrient broth
nm	nanometer
ORF	open reading frame
PAH	polyaromatic hydrocarbon
PCR	polymerase chain reaction
PMSF	phenylmethysulfonyl fluoride
SDS	sodium dodecyl sulphate
SDS-PAGE	sodium dodecyl sulphate polyacrylamide gel electrophoresis
TEMED	N, N, N, N- Tetramethylenediamide
TSB	trypticase soy broth
μg	Microgram
μl	microliter
μm	micrometer
U/ml	Unit per milliliter
U/mg	Unit per milligram
v/v	volume per volume
w/v	weight per volume



CHAPTER 1

INTRODUCTION

The estimated world-wide sales volume for industrial enzymes in 1995 is US\$ 1 billion and this volume is foreseen to double by 2005 (Godfrey and West, 1996). Following proteases and carbohydrases, lipases are considered to be the third largest group based on total sales volume. The demand for industrial enzymes, particularly of microbial origin, is ever increasing owing to their applications in a wide variety of processes. It is in the last decade that lipases have gained importance to a certain extent over proteases and amylases, especially in the area of organic synthesis (Saxena *et al.*, 2005). The interest of lipases stems primarily from their ability to preferentially hydrolyze long/short or saturated/unsaturated fatty acyl residues, but they also exhibit a positional specificity for either the 1 (3) or 1,2 (2,3) positions of a triacylglycerol molecule.

Lipases (E.C. 3.1.1.3) are glycerol ester hydrolases that catalyze the hydrolysis of triacylglycerols into fatty acid, partial acylglycerols and glycerol. Lipases are unique in catalyzing the hydrolysis of fats into fatty acids and glycerol at the water-lipid interface and reversing the reaction in non-aqueous media. The enzyme conformation changes when it contacts with a water-insoluble substrate. This phenomenon is becoming more interesting in understanding the structure-function of enzymes.

