



UNIVERSITI PUTRA MALAYSIA

**DETECTION OF HEPATITIS B CORE ANTIGEN USING A FUSION
BACTERIOPHAGE**

SITI SALWA HASMONI.

FBSB 2005 14

**DETECTION OF HEPATITIS B CORE ANTIGEN USING A FUSION
BACTERIOPHAGE**

**By
SITI SALWA HASMONI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements of the Degree of Master of Science**

September 2005



Specially dedicated to,

Papa and Mama, Kak Long, Kayna, Aji, Alol, Yan, and Limah

For their invaluable love, understanding, patience, support and care.



Abstract of thesis presented to the Senate Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science.

DETECTION OF HEPATITIS B CORE ANTIGEN USING A FUSION BACTERIOPHAGE

By

SITI SALWA BINTI HASMONI

September 2005

Chairman: Associate Professor Tan Wen Siang, PhD

Faculty: Biotechnology and Biomolecular Sciences

Due to the many reported cases of hepatitis B disease around the world, a keen interest among researches has aroused on the cause of the disease, hepatitis B virus (HBV). One of the serological markers for HBV is hepatitis B core antigen (HBcAg) that is a marker of the infectious material and it is the most accurate index of the viral replication. The importance of the HBcAg especially when considering the close relationship with the viral DNA load has created revolutionary studies on the HBcAg ever since. The HBV nucleocapsid or HBcAg is extremely immunogenic during infection and after immunization. A fusion bacteriophage that interacts with HBcAg has been isolated from a phage display peptide library. The phage interacts tightly to HBcAg and thus has the potential to be further developed as a diagnostic reagent. In this study, two immunoassays have been developed using the fusion bacteriophage to detect HBcAg. Phage-ELISA and phage-dot blot assay could detect not only purified HBcAg but also HBcAg in serum samples. As low as 10 ng of HBcAg can be significantly detected by 10^{12} pfu/ml of fusion phage when the reading at 405 nm was measured ($A_{405} = 0.4$). Using the fusion bacteriophage, these newly developed immunoassays



provide an easier and cheaper option for detecting HBcAg. The sensitivity of these immunoassays demonstrates the potential and perhaps vast future uses to detect HBcAg. The fusion phage is also capable of purifying the HBcAg due to its capability to precipitate HBcAg.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGESANAN ANTIGEN TERAS VIRUS HEPATITIS B DENGAN
MENGUNAKAN BAKTERIOFAJ REKOMBINAN**

Oleh

SITI SALWA BINTI HASMONI

September 2005

Pengerusi: Profesor Madya Tan Wen Siang, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Terdapat banyak kes penyakit Hepatitis B telah dilaporkan. Ini menimbulkan minat yang tinggi kepada para penyelidik untuk menjalankan penyelidikan tentang punca penyakit tersebut iaitu virus Hepatitis B (HBV). Salah satu penunjuk serologi virus ini adalah HBcAg. Ia merupakan penanda jangkitan HBV yang sangat efektif dan paling tepat dalam menunjukkan kehadiran aktiviti replikasi virus tersebut. Kepentingan HBcAg terutama mengenai hubungan rapat protein tersebut dengan jumlah DNA HBV telah mencetus revolusi di dalam bidang penyelidikan yang telah memberi kesan yang mendalam. HBcAg atau juga dikenali sebagai nukleokapsid HBV adalah sangat imunogenik semasa jangkitan dan selepas immunisasi. Satu bakteriofaj rekombinan yang telah dipilih daripada sebuah perpustakaan pameran faj didapati dapat berinteraksi kuat dengan HBcAg. Faj yang membawa peptide-peptida tertentu ini berpotensi untuk dijadikan suatu reagen dalam bidang diagnostik. Di akhir pengajian ini, dua jenis immunoasai yang menggunakan bakteriofaj rekombinan untuk mengesan HBcAg telah berjaya dihasilkan. Faj-ELISA dan asai faj-dot blot mampu mengesan HBcAg yang telah dituliskan dan juga HBcAg di dalam sample-sampel serum.

Sejumlah 10^{12} pfu/ml faj rekombinan berupaya mengesan dengan baik 10 ng HBcAg apabila bacaan tindakbalas yang diambil pada gelombang 405 nm diukur ($A_{405} = 0.4$). Imunoasai-imunoasai baru ini menyediakan suatu kaedah yang lebih murah dan mudah dalam mengesan HBcAg disamping mempamerkan pengesanan HBcAg yang sensitif. Oleh yang demikian, imunoasai-imunoasai ini berpotensi tinggi untuk digunakan di masa hadapan dalam mengesan HBcAg khususnya. Faj rekombinan ini juga boleh digunakan untuk menuliskan HBcAg, berdasarkan kebolehannya memendakkan HBcAg.

ACKNOWLEDGEMENTS

I would like to express my deepest appreciation and thanks to my main supervisor, Assoc. Prof. Dr. Tan Wen Siang, for his never ending guidance and encouragement throughout this project. His patience and dedication has inspired me to give my very best in completion of this study and the thesis.

My sincere thanks to my co-supervisor, Prof. Datin Dr. Khatijah Yusoff for her invaluable advice and constructive criticism during the entire progress of this project. Her suggestions and comments definitely helped in the success of this thesis.

A special note of appreciation also goes out to the staff from Faculty of Biotechnology and Biomolecular Studies and everyone who have contributed in one way or another to this study.

I am truly indebted to my lab mates, Kak Tan, Thong Chuan, Kah Fai, Andrew, Swee Tin, Taznim, Yan Peng, Lalita, Kak Fieda, Kak Onie, Abg. Zul, Max, Watti, Kak Sue, Kak Raha and last but not least Dr. Majid Eshaghi for their helpful collaboration and discussion. Also not forgetting my friends, Deela, Nurul, Elia, Fazu, Surini, Che Lina, Mok, Shahrul, Tajul, Lan, and a bunch of others that have really helped me in any possible way there could be. Last but not least, my heartiest appreciation to Soffiyan and his family for their continual support and understanding for me to proceed completion.

May god bless you all.



I certify that an Examination Committee met on 30th September 2005 to conduct the final examination of Siti Salwa Hasmoni on her Master of Science thesis entitled "Detection of Hepatitis B Core Antigen Using a Fusion Bacteriophage" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Janna Ong Abdullah, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Raja Noor Zaliha Raja Abdul Rahman, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Muhajir Hamid, PhD


Lecturer

Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Sheila Nathan, PhD

Associate Professor

Faculty of Science and Technology
Universiti Teknologi Malaysia
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **25 OCT 2005**

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

Tan Wen Siang, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Khatijah Yusoff, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)



AINI IDERIS, PhD

Professor/Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 17 NOV 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



SITI SALWA HASMONI

Date: 24 OKTOBER 2005

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xv
LIST OF FIGURES	xvi
LIST OF ABBREVIATION	xix
 CHAPTER	
1 INTRODUCTION	
Introduction	1
2 LITERATURE REVIEW	
2.1 Hepatitis B Virus (HBV)	5
2.1.1 Hepatitis B Virus Classification	5
2.2 Biology of Hepatitis B Virus	5
2.2.1 HBV Morphology	5
2.2.2 Genomic Organisation	6
2.2.3 Replication of Hepatitis B Virus	8
2.2.4 Viral Proteins	9
2.3 Immunology of Hepatitis B Virus in Human	11
2.3.1 Serological markers	12
2.4 Detection of Hepatitis B Virus	14
2.4.1 Production of Monoclonal Antibodies	15
2.4.2 Radioimmunoassays	16
2.4.3 Immunoassays	16
2.4.4 Molecular Techniques	17
2.4.4.1 PCR	17
2.5 Biology of Filamentous Bacteriophage	18
2.5.1 Morphology of Filamentous Bacteriophage	18
2.5.2 Genomic Organisation and the Products of Filamentous Phage	20
2.5.3 Replication of Filamentous Bacteriophage	21
2.6 Phage Display Technology	22
2.6.1 Phage Display System	23
2.7 Application of Phage Display Technology	24
2.8 Principles of Affinity Chromatography	26
2.8.1 Coupling Gels for Ligand Immobilisation	27



3	MATERIALS AND METHODS	
3.1	Materials	29
	3.1.1 Standard Solutions and Buffers	29
	3.1.1 Liquid and Solid Media	30
	3.1.2 Serum Samples	30
3.2	Preparation and Purification of Full-Length and Truncated HBcAg	31
	3.2.1 The Bradford Assay	32
	3.2.2 SDS-PAGE	33
	3.2.3 Western Blot Analysis	34
3.3	Preparation and Purification of Fusion M13 Bacteriophage	35
	3.3.1 Phage Titration	35
	3.3.2 Large Scale preparation and Purification of Fusion M13 Bacteriophage	36
	3.3.2.1 Extraction of ssDNA	38
	3.3.2.2 Agarose Gel Electrophoresis	38
	3.3.2.3 DNA Sequencing	39
3.4	Detection of HBV Serum Samples with PCR	40
	3.4.1 Extraction of HBV DNA	40
	3.4.2 PCR	40
3.5	Pretreatment of Serum Samples	41
3.6	Phage-Enzyme-Linked Immunoabsorbent Assay (Phage-ELISA)	41
	3.6.1 Optimization of the Concentrations of HBcAg, Fusion Phage, Temperatures, and Blocking Buffers	42
3.7	Enzyme-Linked Immunoabsorbent Assay (ELISA)	44
3.8	Phage-ELISA for the Detection of HBcAg in Serum Samples	45
3.9	Phage-Dot Blot Assay	45
	3.9.1 Optimization of the Concentrations of Fusion Phage, HBcAg, and Anti-HBcAg Monoclonal Antibodies	46
3.10	Dot Blot Assay	48
3.11	Phage-Dot Blot Assay for the Detection of HBcAg in Serum Samples	48
3.12	Specificity Test of Phage-ELISA and Phage-Dot Blot Assay for the Detection of HBcAg using Fusion Phage	49
	3.12.1 Phage-ELISA	49
	3.12.2 Phage-Dot Blot Assay	50
3.13	Phage-Precipitation Assay	51
	3.13.1 Immobilisation (Cross-Linking Optimization)	51
	3.13.2 Precipitation of Full-Length and Truncated HBcAg	51
3.14	Purification of Full-Length and Truncated	



	HBcAg using Fusion M13 Bacteriophage	52
4	RESULTS	
4.1	Preparation, Purification and Analysis of Full-length and Truncated HBcAg	54
4.2	Large Scale Preparation and Purification of Fusion M13 Bacteriophage	56
	4.2.1 Cesium Chloride Ultracentrifugation	56
4.3	Analysis of HBV Samples with PCR	59
4.4	Phage-Enzyme-Linked Immunoabsorbent Assay (Phage-ELISA)	63
	4.4.1 Optimization of the Concentrations of HBcAg, Fusion Phage, Temperatures, And Blocking Buffers	63
4.5	Enzyme-Linked Immunoabsorbent Assay (ELISA)	72
4.6	Phage-ELISA for the detection of HBcAg in Serum Samples	76
4.7	Phage-Dot Blot Assay	76
	4.7.1 Optimization of the Concentrations of the Fusion Phage, HBcAg, and Anti-HBcAg Monoclonal Antibodies	76
4.8	Dot Blot Assay	83
4.9	Phage-Dot Blot Assay for the Detection of HBcAg in Serum Samples	83
4.10	Specificity Test of Phage-ELISA and Phage-Dot Blot Assay for the Detection of HBcAg using Fusion Phage	87
4.11	Phage-Precipitation Assay	90
	4.11.1 Optimization of the Cross-Linking Reaction	90
	4.11.2 Precipitation of Full-Length and Truncated HBcAg	94
5	DISCUSSION	
5.1	Preparation, Purification and Analysis of Full-length and Truncated HBcAg	97
5.2	Large Scale Preparation and Purification of Fusion M13 Bacteriophage	99
5.3	Analysis of HBV Samples with PCR	100
5.4	Phage-Enzyme-Linked Immunoabsorbent Assay (Phage-ELISA)	101
5.5	Enzyme-Linked Immunoabsorbent Assay (ELISA)	103
5.6	Phage-ELISA for the detection of HBcAg in Serum Samples	104
5.7	Phage-Dot Blot Assay	105
5.8	Dot Blot Assay	106
5.9	Phage-Dot Blot Assay for the Detection of HBcAg in Serum Samples	107



5.10	Specificity Test of Phage-ELISA and Phage-Dot Blot Assay for the Detection of HBcAg using Fusion Phage	108
5.11	Phage-Precipitation Assay	109
5.12	Purification of Full-Length and Truncated HBcAg using fusion M13 bacteriophage	110
6	SUMMARY AND CONCLUSION	113
	REFERENCES	117
	BIODATA OF THE AUTHOR	129
	PUBLICATIONS	130



LIST OF TABLES

Table		Page
3.1	Standard solutions and buffers	30
3.2	Parameters and conditions for the first part of Phage-ELISA	43
3.3	Parameters and conditions towards the final optimization of Phage-ELISA	44
3.4	Parameters and conditions for the optimization of Phage-dot blot assay	47
4.1	The number of blue plaque formed or the number of uncoupled ligands after the cross-linking reaction	93
6.1	Final Optimized Parameters and Conditions of Phage-ELISA	116
6.2	Final Optimized Parameters and Conditions for Phage-Dot Blot Assay	116



LIST OF FIGURES

Figure		Page
2.1	A schematic representative of virion structure of HBV	6
2.2	Genome organisation of HBV	7
2.3	Interaction of different cell system in the immune response against HBV	12
2.4	A schematic representative of Ff bacteriophage	19
2.5	The genome of the M13	21
2.6	Principle of affinity chromatography	27
4.1	SDS-PAGE of full-length and truncated HBcAg fractionated on sucrose gradient centrifugation	55
4.2	SDS-PAGE and Western blot of concentrated full-length and truncated HBcAg	57
4.3	Plaque assay showing blue recombinant M13 plaques	58
4.4	Agarose gel electrophoresis of ssDNA fusion M13 bacteriophage	60
4.5	Chromatogram of the nucleotide sequence of the fusion M13 phage clone WSFFSNI	61
4.6	Detection of HBV serum samples using the PCR	62
4.7	Detection of HBcAg using the fusion phage	64
4.8	Optimization of detection of HBcAg using different set of concentration of fusion phage	66
4.9	Detection of different range of amount of HBcAg using the fusion phage	67



4.10	Detection of a narrower range of HBcAg using the fusion phage	68
4.11	Graph representing the detection of HBcAg using 1.0×10^{10} to 1.0×10^{12} pfu/ml of the fusion phage with milk diluent as the blocking buffer	70
4.12(a)	Graph presenting the detection of HBcAg using 1.0×10^{10} to 1.0×10^{12} pfu/ml of the fusion phage that uses SEA blocking solution as the blocking buffer	71
4.12(b)	Detection of HBcAg using 1.0×10^{10} to 1.0×10^{12} pfu/ml of fusion phage with different blocking duration	71
4.13	Optimization of the coating duration for the detection of HBcAg using fusion phage (1.0×10^{10} to 1.0×10^{12} pfu/ml)	73
4.14	Optimised detection of HBcAg using fusion phage via Phage-ELISA	74
4.15	Detection of HBcAg with anti-HBcAg monoclonal antibody via ELISA	75
4.16	Detection of HBcAg in HBV positive serum samples using fusion phage via phage-ELISA	77
4.17	Optimization of the blocking buffer for the detection of HBcAg using fusion phage	79
4.18	Optimization of different duration and temperature of HBcAg incubation for phage-dot blot assay	80
4.19	Optimization of the duration and temperature of anti-	81



	HBcAg incubation for phage-dot blot assay	
4.20	Optimization of the dilutions of anti-HBcAg mAb (primary antibody) for phage-dot blot assay	82
4.21	Detection of HBcAg with fusion phage via dot blot assay	84
4.22	Detection of HBcAg with anti-HBcAg monoclonal antibody via dot blot assay	85
4.23	Detection of HBcAg in HBV positive serum samples with fusion phage via dot blot assay	86
4.24	Specificity test results for phage-ELISA	88
4.25	The specificity test phage-dot blot results that correspond to that of the phage-ELISA	89
4.26	Specificity test results for phage-ELISA	91
4.27	Specificity test phage-dot blot assay results	92
4.28	SDS-PAGE of precipitated products using the fusion phage immobilized on CNBr activated-agarose as ligand	95
4.29	Western blotting of precipitated products with anti-HBcAg antibody	96



LIST OF ABBREVIATIONS

ϵ	encapsulation signal
α	alpha
β	beta
μg	microgram (10^{-6} g)
μl	microlitre (10^{-6} l)
μM	micromolar (10^{-6} M)
Amp	ampicillin
anti-HBcAg	antibody to HBcAg
anti-HBeAg	antibody to HBeAg
anti-HBsAg	antibody to HBsAg
anti-HBxAg	antibody to HBxAg
$^{\circ}\text{C}$	degree centigrade
CHAPS	3-[(3-chol-amidopropyl)-dimethylammonio]-1-propanesulfonate
CNBr	cyanogen bromide
bp	base pair
BSA	bovine serum albumin
C	cytosine/core
ccc	covalently closed circular
CTL	cytotoxic T lymphocyte
c-terminus	carboxyl terminus
DC	dendritic cell
DNA	deoxy-ribonucleic acid



DNase	Deoxyribonuclease
dNTP	deoxynucleoside triphosphate
DTT	1,4-dithiothreitol
ds	Double stranded
EDTA	ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunoabsorbent assay
g	Gram
h	Hour
HBcAg	hepatitis B core antigen
HBeAg	hepatitis B e antigen
HBsAg	hepatitis B surface antigen
HBV	hepatitis B virus
HBxAg	hepatitis B x antigen
HCC	hepatocellular carcinoma
HCl	hydrochloric acid
IFN	Interferon
IFN- γ	interferon gamma
IL2	interleukin 2
IL4	interleukin 4
IL5	interleukin 5
IL10	interleukin 10
IL12	interleukin 12
HIV	human immunodeficiency virus
HLA	human leukocyte antigen
IgG	immunoglobulin G



IgM	immunoglobulin M
IPTG	isopropyl- β -d-thiogalactopyranoside
kb	kilobase
kDa	kilodalton
K_D^{rel}	relative dissociation constant
l	Litre
LB	Luria Bertani
L-HBsAg	large surface antigen
M	Molar
mAb	monoclonal antibody
mg	milligram (10^{-3} g)
MgCl ₂	magnesium chloride
M-HBsAg	medium surface antigen
min	Minute
ml	millilitre (10^{-3} l)
mm	milimeter (10^{-3} m)
mRNA	messenger ribonucleic acid
NaCl	sodium chloride
NDV	Newcastle disease virus
NK	natural killer
nM	nanomolar (10^{-9} M)
nm	nanometer (10^{-9} m)
nt(s)	nucleotide(s)
N-terminus	amino terminus
OD	optical density

ORF	open reading frame
P	polymerase protein
PAGE	polyacrylamide gel electrophoresis
PBS	phosphate buffered saline
PCR	Polymerase Chain Reaction
PEG	polyethylene glycol
pfu	plaque forming unit
pgRNA	pregenomic RNA
preC	precore
preC/C	precore and core
preS	hepatitis B preS genes
preS/S	preS and surface
PreS1	N-terminal region of L-HBsAg comprising 108 or 119 amino acid
PreS2	region of M and L-HBsAg comprising 55 amino acid
RF	replicative form
RNA	ribonucleic acid
RNase	ribonuclease
rpm	revolutions per minute
RT	room temperature
s	second
SDS	sodium dodecyl sulphate
S-HBsAg	small surface antigen
ssDNA	single stranded DNA
TAE	tris acetate EDTA buffer



Taq	thermus aquaticus thermostable DNA
TBE	tris-buffered EDTA solution
TBS	tris-buffered saline
TE	tris-EDTA buffer
TEMED	tetramethyl ethylenediamine
Th1	T helper 1
Th2	T helper 2
TP	terminal protein
USA	United States of America
v	Volt
v/v	volume/volume
WHO	World Health Organization
w/v	weight/volume
x <i>g</i>	centrifugal force
X-gal	5-bromo-4-chloro-3-indol- β -D-galactosidase



AMINO ACID ABBREVIATIONS

	One letter code	Three letter code
Alanine	A	Ala
Arginine	R	Arg
Asparagine	N	Asn
Aspartic acid	D	Asp
Cysteine	C	Cys
Glutamic acid	E	Glu
Glutamine	Q	Gln
Glycine	G	Gly
Histidine	H	His
Isoleucine	I	Ile
Leucine	L	Leu
Lysine	K	Lys
Methionine	M	Met
Phenylalanine	F	Phe
Proline	P	Pro
Serine	S	Ser
Threonine	T	Thr
Tryptophan	W	Trp
Tyrosine	Y	Tyr
Valine	V	Val

