



UNIVERSITI PUTRA MALAYSIA

**PARTIAL PURIFICATION AND CHARACTERISATION OF ALKALINE
PHOSPHATASE FROM HEPATOPANCREAS AND INTERTINE OF
RED TILAPIA, (TILAPIA MOSSAMBICA)**

VANITHA MARIAPPAN.

FBSB 2005 30

**PARTIAL PURIFICATION AND CHARACTERISATION OF
ALKALINE PHOSPHATASE FROM HEPATOPANCREAS AND
INTESTINE OF RED TILAPIA, (*Tilapia mossambica*)**

By

VANITHA MARIAPPAN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfillment of the Requirement for the
Degree of Master of Science**

July 2005



DEDICATION

Specially dedicated to the memory of my dear aiyah, late Mr. **MARIAPPAN ARUNASALAM** who left me & my siblings 15 years ago and to my beloved amma **BABY MARIAPPAN** who have been standing strong all her life just for us.....

Aiyah even if you are gone for long,

Your presence I feel within me,

But you will always be in my memory,

As sweet as it can be,

All my life forever & ever...

**HARD IS LIFE,
FOR HE WHO DESIRE DEATH,
BUT LIVES ON,
FOR THE SAKE OF HIS LOVED ONE!!!**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfilment of the requirement for the degree of Master of Science

**PARTIAL PURIFICATION AND CHARACTERISATION OF
ALKALINE PHOSPHATASE FROM HEPATOPANCREAS AND
INTESTINE OF RED TILAPIA, (*Tilapia mossambica*)**

By

VANITHA MARIAPPAN

July 2005

Chairman: Professor Nor Aripin Shamaan, PhD

Faculty: Biotechnology and Biomolecular Sciences

Alkaline phosphatase (EC 3.1.3.1) is a metalloenzyme, which catalyze nonspecific hydrolysis of phosphate monoesters. Partial purification was conducted on alkaline phosphatase (ALP) extracted from hepatopancreas and intestine of red tilapia, (*Tilapia mossambica*) using two main steps – ammonium sulphate precipitation and ion exchange chromatography on DEAE – 52. Samples from the ion-exchange step were analysed for ALP activities and characterised by SDS-PAGE. SDS-PAGE analysis showed 2 identical bands and was found to have molecular weight of 68, 000 Da (hepatopancreas ALP) and 180, 500 Da (intestinal ALP) subunits. Overall, purification fold obtained from the final step are 1.8 and 21.9 for hepatopancreas and intestinal



respectively, with recovery of only 0.22% from hepatopancreas and 0.01% from intestine. The specific activity of the enzyme was $1.72 \times 10^{-2} \mu\text{mol min}^{-1} \text{mg}^{-1}$ and $2.93 \times 10^{-1} \mu\text{mol min}^{-1} \text{mg}^{-1}$ from hepatopancreas and intestine respectively. The ALP from hepatopancreas remained stable at temperatures up to 50°C , and ALP from intestine enzyme had an optimum temperature of 60°C . The optimum pH for both hepatopancreas and intestine ALP of *Tilapia mossambica* is pH 10. The positive monovalent alkali metal ions (Li^+ , Na^+ and K^+) have no effect on the ALP enzyme activity. However, the positive divalent alkali metal ions (Mg^{2+} and Ca^{2+}) activate the enzyme activities. Heavy metal ions (Zn^{2+} , Cu^{2+} , Cd^{2+} and Hg^{2+}) were found to inhibit the enzyme activity.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra
Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENULENAN SEPARA DAN PENCIRIAN ENZIM ALKALINE
FOSFATASE DARI HEPATOPANKREAS DAN USUS IKAN
TILAPIA MERAH (*Tilapia mossambica*)**

Oleh

VANITHA MARIAPPAN

Julai 2005

Pengerusi: Profesor Nor Aripin Shamaan, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Alkaline fosfatase (EC 3.1.3.1) adalah metalloenzim yang hidrolisis secara takspesifik fosfat monoester. Penulenan separa telah dijalankan ke atas alkaline fosfatase (ALP) dari hepatopankreas dan usus ikan tilapia merah (*Tilapia mossambica*) dengan menggunakan dua kaedah utama iaitu pemendakan amonium sulfat dan kromatografi penukaran ion DEAE - 52. Sampel dari penukaran ion DEAE yang mengandungi aktiviti ALP dianalisis dan dicirikan melalui kaedah elektroforesis SDS. Analisis SDS-PAGE telah menunjukkan 2 jalur yang serupa dan berat molekulnya adalah 68, 000 Da (ALP hepatopankreas) dan 180, 500 Da (ALP usus) bagi subunitnya. Secara keseluruhannya, faktor penulenan yang diperolehi daripada kaedah terakhir ialah 1.8

(hepatopankreas) dan 21.9 (usus) dengan pemulihan hanya 0.22% dari hepatopankreas dan 0.01% dari usus. Aktiviti spesifik enzim ialah $1.72 \times 10^{-2} \text{ U mg}^{-1}$ bagi hepatopankreas dan $2.93 \times 10^{-1} \text{ U mg}^{-1}$ bagi usus. ALP hepatopankreas adalah stabil pada suhu sehingga 50°C dan ALP usus pula mempunyai suhu optimum 60°C . pH optimum bagi kedua-dua hepatopankreas dan usus *Tilapia mossambica* adalah pH 10. Ion monovalen logam alkali positif (Li^+ , Na^+ and K^+) tidak memberikan sebarang kesan kepada aktiviti ALP. Bagaimana pun, ion divalen logam alkali positif (Mg^{2+} and Ca^{2+}) mengaktifkan aktiviti enzim. Ion logam berat pula (Zn^{2+} , Cu^{2+} , Cd^{2+} and Hg^{2+}) menghalang tindak balas enzim.



ACKNOWLEDGEMENTS

First of all, I would like to thank GOD for his blessing.

I wish to convey my deepest gratitude and greatest appreciation to my supervisor Assoc. Prof. Dr. Nor Aripin Shamaan for his guidance, advice, encouragement, keen interest, and support throughout my whole project and course of the study. I am thankful to Dr. Nor Aripin for being very understanding when I was going through tough times especially when I was ill and hospitalised.

Words cannot express my heartfelt thanks to my co-supervisors; Dr. Yunus Abd Shukor and Prof. Dr. Mohd Arif Syed for their supervision, providing useful information, constructive suggestions, invaluable advice, and for extending their time discussing about my research work.

Furthermore, I am also indebted to Mr. Jasni from the Hatchery Unit, Universiti Putra Malaysia for providing fresh live tilapia fishes used for the experiments without any charge.



Thanks are also extended to the member of Toxicology Lab especially to Noor Azlina Masdor and Suhaidah Ahmat @ Amirrudin for their invaluable help, assistance and co-operation and the fun we had during work for the past two years. Thanks to my fellow graduate students especially Yap Wai Sum, Lailatul Jumaiyyah, Anthony Chin Chee Meng, Palaniammal Krishnan and Putri Noor Faizah, for being such a wonderful friends.

A very special thanks goes to my dearest friends; Mr. Sunil Bhalla and Dr. Sreeramanan Subramaniam for their moral support, sincere advice, and encouragement throughout the completion of this study. I really appreciate it. Thank you friends!!!

Last but definitely not least, I am deeply grateful and thankful to my beloved mother who have helping me financially, and for the love she showed me all my live; also to my sisters, brother, brothers in-law, my first ever niece and my boyfriend for their support and encouragements. Without the understanding and love of these people everything would have been impossible. GOD BLESS YOU ALL AND THANK YOU!



I certify that an Examination Committee met on 20th July 2005 to conduct the final examination of Vanitha Mariappan on her Master of Science thesis entitled “Partial Purification and Characterisation of Alkaline Phosphatase from Hepatopancreas and Intestine of Red Tilapia (*Tilapia mossambica*)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Radzali Muse, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Johari Ramli, PhD


Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Muhajir Hamid, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Yasmin Anum Mohd Yusof, PhD

Associate Professor
Faculty of Medicine
Universiti Kebangsaan Malaysia
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 22 AUG 2005

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory committee are as follows:

Nor Aripin Shamaan, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Mohd Arif Syed, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)

Mohd Yunus Abdul Shukor, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)



AINI IDERIS, PhD

Professor/Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 08 SEP 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations that have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any degree at Universiti Putra Malaysia or other institutions.



VANITHA MARIAPPAN

Date: 12/8/05

TABLE OF CONTENT

	Page
DEDICATION	iii
ABSTRACT	iv
ABSTRAK	vi
ACKNOWLEDGEMENTS	viii
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xxii

CHAPTER

1.0 INTRODUCTION	1
2.0 LITERATURE REVIEW	5
2.1 Development of fishing industry in Malaysia	5
2.2 Culture of hybrid of tilapia	8
2.2.1 Taxonomy and distribution	8
2.2.2 Life history characteristics	11
2.2.3 Hepatopancreas and digestive system	14
2.3 Phosphatase	17
2.3.1 Role of phosphatase	19
2.4 Specificity of Acid and Alkaline Phosphatase	28
2.5 Definition of Alkaline Phosphatase	31
2.5.1 Characteristics of ALP	35
2.5.2 ALP as a hydrolase	41
2.5.3 Molecular and physical properties of ALP	46
2.5.4 Active site of ALP	48
2.5.5 Functions of ALP	52
2.6 Diseases related to significant of ALP	54
2.6.1 Diseases of bone hyperparathyroidism	54
2.6.2 Paget's disease	55
2.6.3 Osteomalacia	55



2.6.4	Bone tumors	56
2.6.5	Disease of the liver	56
2.6.6	Infectious hepatitis	57
2.6.7	Bone fractures	57
2.6.8	Other diseases if greater or lower than normal ALP level	57
2.6.9	Effects of ALP in fish	58
2.7	Theory of ammonium sulphate precipitation	58
2.8	Theory of ion exchange chromatography	59
2.9	Theory of isoelectric focusing	59
3.0	MATERIALS AND METHODS	60
3.1	Materials	60
3.1.1	Sources of fish	60
3.1.2	Chemical and biochemical reagents	61
3.1.3	Apparatus	62
3.2	Methods	63
3.2.1	Determination of protein content	63
3.2.2	Assay of enzyme activity	66
3.2.3	Extraction of crude enzyme	68
3.2.4	Purification of ALP enzyme	69
3.2.5	Ion-exchange chromatography on DEAE	72
3.2.6	Relative molecular weight determination	76
3.2.7	Isoelectric focusing	82
3.2.8	Characterisation on ALP enzyme	83
3.2.9	Experimental design and statistical analysis	85
3.2.10	Flow chart of the research project	86
4.0	RESULT AND DISCUSSION	87
4.1	Assay of the enzyme activity	87
4.2	Extraction of crude enzyme	88
4.3	Purification of ALP enzyme	90
4.3.1	Ammonium sulphate precipitation $(\text{NH}_4)_2\text{SO}_4$	90
4.3.2	Dialysis	97
4.4	Ion-exchange chromatography	104
4.4.1	Sample binding	104
4.4.2	DEAE-52 chromatography	106
4.5	Relative molecular weight determination	112
4.5.1	SDS-polyacrylamide gel electrophoresis	112



4.5.2	Determination of gel percentage	113
4.5.3	Molecular weight determination	118
4.6	Isoelectric focusing	121
4.7	Characterisations of the ALP enzyme	125
4.7.1	Determination of protein content and enzyme	125
4.7.2	Effect of incubation time on ALP	126
4.7.3	Effect of temperature on ALP	127
4.7.4	Effect of pH on ALP	130
4.7.5	Effect of metal ions on ALP	132
4.7.6	Effect of heavy metals on ALP	137
4.7.7	Assay of the Michaelis-Menten constant (K_m) and maximum velocity (V_{max})	143
5.0	CONCLUSION	145
	REFERENCES	148
	APPENDICES	156
	BIODATA OF THE AUTHOR	171



LIST OF TABLES

Table	Page
1. Hydrolysis of various O- and S-substituted monoesters of phosphorothioic acid by alkaline and acid phosphatases.	29
2. Molecular weights of ALP in different sources	36
3. Types of glycoprotein found in different sources	38
4. Physical properties of ALP	47
5. Total protein and activities of the ALP enzyme from the organs of <i>Tilapia mossambica</i> .	89
6. Summary of the purification for alkaline phosphatase from <i>Tilapia mossambica</i> hepatopancreas.	110
7. Summary of the purification for alkaline phosphatase from <i>Tilapia mossambica</i> intestine.	110
8. Protein content and enzyme activities of the ALP enzyme from the purified hepatopancreas and intestine of <i>Tilapia mossambica</i> .	125
9. Comparison of characterizations of hepatopancreas and intestine of <i>Tilapia mossambica</i> .	147



10. Differences of freshwater aquaculture production (tan metric) for year 2002 and 2003	170
11. Freshwater fish production values for year 2002 and 2003	170
12. Changes in numbers of freshwater fish aquaculture between year 2002 and 2003	171
13. Differences in (hectares) of freshwater fish aquaculture between year 2002 and 2003	171



LIST OF FIGURES

Figure	Page
1. The enzyme nomenclature of alkaline phosphatase.	2
2. Schematic diagram of red tilapia (<i>Tilapia mossambica</i>).	8
3. Schematic diagram of fish anatomy.	13
4. Schematic diagram of the alkaline phosphatase hydrolytic reaction	31
5. Molecular model of alkaline phosphatase structure.	33
6. Two-steps process hydrolysis by ALP.	41
7. Hydrolysis and phosphotransferase mechanism phosphate monoesters by ALP.	43
8. The two-step mechanism of alkaline phosphatase.	45
9. Molecular states of ALP under various conditions.	46
10. Active site of ALP.	49
11. Red tilapia (<i>Tilapia mossambica</i>).	60



12. Hepatopancreas of fish.	61
13. Electrophoresis apparatus.	62
14. Phosphate hydrolysis of <i>p</i> -NPP to <i>p</i> -NP.	66
15. Schematic diagram of ion exchange chromatography method.	75
16. Schematic diagram of the SDS-PAGE electrophoresis gel.	78
17. Over all flow chart of the research project.	86
18. (a) Graph of amount of protein (mg) and total enzyme activity of ALP supernatant ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from hepatopancreas supernatant.	91
18. (b) Graph of amount of protein (mg) and total enzyme activity of ALP supernatant ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from intestine supernatant.	92
19. (a) Graph of amount of protein (mg) and total enzyme activity of ALP supernatant ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from hepatopancreas supernatant.	95
19. (b) Graph of amount of protein (mg) and total enzyme activity of ALP supernatant ($\mu\text{mole min}^{-1}$) versus	



percentage of ammonium sulphate precipitation (%) from intestine supernatant.	96
20. (a) Graph of amount of protein (mg) and total enzyme activity of ALP pellet ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from hepatopancreas pellet.	99
20. (b) Graph of amount of protein (mg) and total enzyme activity of ALP pellet ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from intestine pellet.	100
21. (a) Graph of amount of protein (mg) and total enzyme activity of ALP pellet ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from hepatopancreas pellet.	102
21. (b) Graph of amount of protein (mg) and total enzyme activity of ALP pellet ($\mu\text{mole min}^{-1}$) versus percentage of ammonium sulphate precipitation (%) from intestine pellet.	103
22. pH of DEAE binding versus absorbance at 410nm (ALP enzyme activity).	105
23. (a) Chromatography on DEAE-52 with hepatopancreas from the $(\text{NH}_4)\text{SO}_4$ step.	108
23. (b) Chromatography on DEAE-52 with intestine from the $(\text{NH}_4)\text{SO}_4$ step.	109
24. (a) SDS-polyacrylamide gel electrophoresis of under reduced conditions of hepatopancreas alkaline phosphatase of tilapia using a 15% gel.	114



24. (b) SDS-polyacrylamide gel electrophoresis of under reduced conditions of intestine alkaline phosphatase of tilapia using a 15% gel.	115
25. (a) SDS-polyacrylamide gel electrophoresis of under reduced conditions of hepatopancreas alkaline phosphatase of tilapia using a 12% gel.	116
25. (b) SDS-polyacrylamide gel electrophoresis of under reduced conditions of intestine alkaline phosphatase of tilapia using a 12% gel.	117
26. (a) Log standard molecular weight plot against R_f value of hepatopancreas to determine the molecular weight of alkaline phosphatase enzyme.	119
26. (b) Log standard molecular weight plot against R_f value of intestine to determine the molecular weight of alkaline phosphatase enzyme.	120
27. (a) Isoelectric focusing on hepatopancreas	122
27. (b) Isoelectric focusing on intestine	123
28. Influence of incubation time (minutes) on hepatopancreas and intestinal ALP activity.	126
29. Effect of temperature on hepatopancreas and intestinal ALP activity.	129



30. Effect of pH values on hepatopancreas and intestinal ALP activity.	132
31. (a) Activation effects of metal ions on hepatopancreas and intestinal ALP activity Mg^{2+}	135
31. (b) Activation effects of metal ions on hepatopancreas and intestinal ALP activity Ca^{2+} .	136
32. (a) Inhibition effects of heavy metals on hepatopancreas and intestinal ALP activity Zn^{2+}	139
32. (b) Inhibition effects of heavy metals on hepatopancreas and intestinal ALP activity Cu^{2+}	140
32. (c) Inhibition effects of heavy metals on hepatopancreas and intestinal ALP activity Cd^{2+} .	141
32. (d) Inhibition effects of heavy metals on hepatopancreas and intestinal ALP activity Hg^{2+} .	142
33. Saturation curves on <i>p</i> -nitrophenyl phosphate substrate of both hepatopancreas and intestinal ALP activity.	144



LIST OF ABBREVIATION

α	- alpha
β	- beta
$^{\circ}\text{C}$	- degree Celsius
%	- percentage
Abs	- absorbance
ALP	- alkaline phosphatase
APS	- ammonium persulphate
BSA	- bovine serum albumin
cm	- centimetre
Da	- Dalton
DEAE	- diethylaminoethylcellulose
IEF	- isoelectric focusing
kD/kDa	- kilodalton
kg	- kilogram
mA	- mili Amp
mg	- miligram
ml	- mililiter
min	- minute

mins	- minutes
MW	- molecular weight
NaOH	- sodium hydroxide
$(\text{NH}_4)_2\text{SO}_4$	- ammonium sulfate
nm	- nanometer
pH	- (-) log concentration of H^+
<i>p</i> -NPP	- para nitrophenyl phosphate
<i>p</i> -NP	- para nitrophenol
SDS	- sodium dodecyl sulphate
U	- unit
μmol	- micromole
μg	- microgram
X g	- gravity
w/v	- weight/volume
w/w	- weight/weight

CHAPTER 1

INTRODUCTION

The existence of enzymes has been known for well over a century. Some of the earliest studies were performed in 1853 by the Swedish chemist Jon Jacob Berzelius who termed their chemical action catalytic. It was not ever since James B. Sumner of Cornell University purified and crystallizes the enzyme urease from the jack bean (EC 3.5.1.5) for the first time in the year 1926 as mentioned in Dixon and Webb, (1979); biochemists have successfully purified perhaps, thousands of enzymes thus far.

Alkaline phosphatase, ALP, (EC 3.1.3.1) is found in abundance in nature. ALP is a hydrolase and catalyzes the hydrolysis of various bonds. It has found in wide application especially in molecular biology, medical, and industries. It is a hydrolytic enzyme which catalyses the cleavage of a chemical bond with the addition of water (McComb *et al.*, 1979). ALP is in a group of enzyme acting on ester bonds and these esterases are subdivided into those acting on phosphoric monoester hydrolases, the phosphatases under an alkaline condition (McComb *et al.*, 1979).

