



**UNIVERSITI PUTRA MALAYSIA**

**EFFECT OF PALM KERNEL CAKE BASED DIET ON LIBIDO AND SEMEN QUALITY OF MALIN X SANTA-INES LAMBS**

**ZELEALEM TESFAY GEBRETSADIK.**

**FP 2004 5**

**EFFECT OF PALM KERNEL CAKE BASED DIET ON LIBIDO AND SEMEN  
QUALITY OF MALIN x SANTA-INES LAMBS**

**By**

**ZELEALEM TEFAY GEBRETSADIK**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of Master of  
Agricultural Science**

**January 2004**



## **DEDICATION**

**This thesis is dedicated to the people of ETHIOPIA. I owe my country a great debt.**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Agricultural Science

**EFFECT OF PALM KERNEL CAKE BASED DIET ON LIBIDO AND SEMEN QUALITY OF MALIN x SANTA-INES LAMBS**

By

**ZELEALEM TEFAY GEBRETSADIK**

**January 2004**

**Chairman: Associate Professor Ismail Bin Idris, Ph.D.**

**Faculty: Agriculture**

A study was conducted for six months at the farm of the Department of Animal Science, Universiti Putra Malaysia (UPM), Serdang, Selangor (3° 00' N, 101° 42' E) to investigate the effect of palm kernel cake based diet on serving capacity and semen quality of Malin x Santa-Ines crossbred ram lambs. Twenty Malin x Santa-Ines crossbred ram lambs were randomly assigned into three dietary treatments namely (T1) 60% Palm kernel cake + 40% oil palm frond, (T2) 60% Palm kernel cake + 40% oil palm frond supplemented with 23 mg/kg dry matter of molybdenum as ammonium molybdate ((NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>·4H<sub>2</sub>O) and 600mg/kg dry matter of Sulfur as sodium sulphate (Na<sub>2</sub>SO<sub>4</sub>) and (T3) control; 60% concentrate of corn-soybean mix + 40% oil



palm frond. There was no significant difference ( $p>0.05$ ) among the rams for the three dietary treatments for libido or mating behavior. All rams reached their optimum mating behavior (number of ejaculation =  $6.7\pm 1.3$  and mating efficiency =  $0.43\pm 0.01$ ) after the second month of the experiment. It was also found that there was no significant difference ( $p>0.05$ ) for all the semen quality parameters among the rams under the three dietary treatments. Rams in all dietary treatments produced good semen and their sperm characteristics were within the ranges for good rams (semen volume = 0.7-2.0 ml; sperm concentration =  $2-5 \times 10^9$  cells/ml; sperm motility > 80%; percentage of abnormal morphology <15%).

Although there was a significant difference on the level of copper (Cu) accumulated in the liver ( $p<0.01$ ) and testis ( $p<0.05$ ), with highest values in rams under T1 ( $1089 \pm 189$  ppm and  $0.63 \pm 0.18$  ppm, respectively), there was no symptom of toxicity observed in the rams. Therefore, it was concluded that PKC has no negative effect on serving capacity and semen quality of Malin x Santa-Ines crossbred sheep, at least for six months of the feeding period. It was also concluded that supplementing 23 mg/kg dry matter of Molybdenum and 600 mg/kg dry matter Sulphur to sheep fed on PKC based diet helps to protect Cu accumulation in the plasma, liver and testis of rams without negative effect on their reproductive capacity and semen quality.



Even though higher serum testosterone was found in rams under diets T1 and T2 compared to T3, this did not reflect the actual reproductive level and sperm quality. Hence, it was concluded that serum testosterone level might not be a good indicator of reproductive performance of rams when blood samples are taken from conscious animals.

**Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan ijazah Mster Sains Pertanian**

**KESAN DIET ASAS ISIRONG KELAPA SAWIT KE ATAS LIBIDO DAN  
KUALITI SEMEN ANAK BIRI-BIRI BAKA KACUKAN MALIN x SANTA-INES**

**Oleh**

**ZELEALEM TEFAY GEBRETSADIK**

**Januari 2004**

**Pengerusi: Professor Madya Ismail Bin Idris, Ph.D.**

**Fakulti: Pertanian**

Satu Kajian telah dijalankan selama enam bulan di ladang Jabatan Sains Haiwan, Universiti Putra Malaysia (UPM), Serdang, Selangor (3° 00' Utara, 101° 42' Timur) untuk menyelidik kesan isirong kelapa sawit (PKC) sebagai diet asas ke atas keupayaan mengawan dan kualiti semen biri-biri baka kacukan Malin x Santa-Ines. Dua puluh ekor biri-biri Malin x Santa-Ines telah ditentukan secara rawak kepada tiga rawatan diet iaitu (T1) 60% PKC + 40% pelet pelepah kelapa sawit, (T2) 60% PKC + 40% pelet pelepah kelapa sawit di suplemen dengan 23 mg/kg berat kering molybdenum sebagai ammonium molybdate ((NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>.4H<sub>2</sub>O) dan 600 mg/kg berat kering sulfur sebagai sodium sulfat



(Na<sub>2</sub>SO<sub>4</sub>) dan (T3) kawalan; 60% konsentrat mengandungi campuran jagung-kacang soya + 40% pelet pelepah kelapa sawit. Tiada perbezaan bererti ( $p > 0.05$ ) diantara biri-biri jantan pada ketiga-tiga rawatan diet untuk 'libido' atau kelakuan mengawan. Semua biri-biri jantan mencapai kelakuan mengawan yang optimum (bilangan ejakulasi =  $6.7 \pm 1.3$  dan efisiensi mengawan =  $0.43 \pm 0.01$ ) selepas dua bulan kajian dijalankan. Tiada perbezaan bererti ( $p > 0.05$ ) juga didapati untuk semua parameter kualiti semen diantara biri-biri jantan untuk ketiga-tiga rawatan diet. Biri-biri jantan dalam semua rawatan diet menghasilkan semen yang berkualiti dan ciri-ciri sperma yang dihasilkan adalah dalam julat untuk biri-biri jantan yang baik (isipadu semen =  $0.7-0.2$  ml; kepekatan sperma =  $2-5 \times 10^9$  sel/ml; motiliti sperma  $> 80\%$ ; peratusan morfologi sperma abnormal  $< 15\%$ ).

Walau bagaimanapun, terdapat perbezaan bererti pada tahap kuprum terkumpul dalam hati ( $p < 0.01$ ) dan testis ( $p < 0.05$ ), dengan nilai yang tertinggi bagi biri-biri jantan di bawah T1 ( $1089 \pm 189$  ppm dan  $0.63 \pm 0.18$ , masing-masing), tidak terdapat simptom keracunan pada biri-biri jantan. Oleh yang demikian, dapat dirumuskan bahawa tiada kesan negatif PKC ke atas keupayaan mengawan dan kualiti semen biri-biri jantan baka kacukan Malin x Santa-Ines, sekurang-kurangnya untuk jangkamasa pemakanan selama enam bulan. Ia juga dapat dirumuskan bahawa dengan memberi suplemen 23 mg/kg berat kering molybdenum dan 600 mg/kg berat kering sulfur kepada biri-biri yang diberi PKC sebagai diet asas akan menolong melindungi pengumpulan



Cu di dalam plasma, hati dan testis biri-biri jantan tanpa kesan negatif terhadap keupayaan reproduktif dan kualiti semen.

Walaupun testosteron dalam serum didapati tinggi pada biri-biri jantan di bawah T1 dan T2 berbanding T3 ini tidak menggambarkan tahap reproduktif sebenar dan kualiti sperma. Oleh itu, ia dapat dirumuskan bahawa tahap testosterone dalam serum mungkin merupakan petunjuk yang kurang baik bagi prestasi reproduktif biri-biri jantan apabila sampel darah diambil daripada haiwan yang sedar.



## ACKNOWLEDGEMENTS

*At most Thanks to Almighty God!*

I would like to express my sincere and profound gratitude to my main supervisor, Assoc. Prof. Dr. Ismail Idris, for his invaluable guidance and friendship throughout the course of my study. My deep appreciation is also extended to Assoc. Prof. Dr. Halimatun Yaakub, Assoc. Prof. Dr. Sabrina Sukardi and Assoc. Prof. Dr. Abdul Razak Alimon for their invaluable help and guidance.

My deepest appreciation goes to Dr. Jothi Malar Panandam for her devotion to avail herself to answer questions related to my work and allowing me to use the laboratory facilities. I am also grateful to Assco. Prof. Dr. Abdul Rahman Omar from the Faculty of Veterinary Medicine for his permission to use the lab facilities in the Virology laboratory and for assisting me in the analysis and interpretation of the results of enzyme immuno assay. Thanks are also due to Mr. Jamiuddin Shahudin for assisting me in the animal farm and to Ms. Lai Pui Wah for translating my abstract to “Bahasa Malaysia”. The collaboration, help and encouragement from all Ethiopian students in UPM and fellow students in the Department of Animal Science are acknowledged. Finally I would like to give my sincere gratitude to my father and all my family for their encouragement they offered me.



I certify that an Examination Committee met on 26<sup>th</sup> January 2004 to conduct the final examination of Zelealem Tesfay Gebretsadik on his Master of Agricultural Science thesis entitled "Effect of Palm Kernel Cake Based Diet on Libido and Semen Quality of Malin X Santa-Ines Lambs" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Zainal Aznam Mohad Jelani, Ph.D.**

Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Wahid bin Haron, Ph.D.**

Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**Liang Juan Boo, Ph.D.**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Hiroyuki Hirooka, Ph.D.**

Professor  
Graduate School of Agriculture  
Kyoto University  
Japan  
(Independent Examiner)



---

**GULAM RUSUL RAHMAT ALI, Ph.D.**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 22 MAR 2004

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Agricultural Science. The members of the Supervisory Committee are as follows:

**Ismail bin Idris, Ph.D.**  
Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Halimatun Yaakub, Ph.D.**  
Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Abd Razak bin Alimon, Ph.D.**  
Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Sabrina Sukardi, Ph.D.**  
Associate Professor  
Faculty of Medicine and Health Science  
Universiti Putra Malaysia  
(Member)



---

**AINI IDERIS, Ph.D.**  
Professor / Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 09 APR 2004



## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

 *zelealem  
Tefay*

---

**ZELEALEM TEFAY GEBRETSADIK**

Date: *24/03/04*

## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	<b>II</b>
<b>ABSTRACT</b>	<b>III</b>
<b>ABSTRAK</b>	<b>VI</b>
<b>ACKNOWLEDGEMENTS</b>	<b>IX</b>
<b>APPROVAL</b>	<b>X</b>
<b>DECLARATION</b>	<b>XII</b>
<b>LIST OF TABLES</b>	<b>XV</b>
<b>LIST OF FIGURES</b>	<b>XVI</b>
<b>LIST OF PLATES</b>	<b>XVII</b>
<b>ABBREVIATIONS</b>	<b>XVIII</b>

### CHAPTER

<b>I. INTRODUCTION</b>	<b>1</b>
<b>II. LITERATURE REVIEW</b>	<b>5</b>
<b>TRACE ELEMENTS IN ANIMAL</b>	<b>5</b>
Importance of Copper	6
Copper Homeostasis	7
Copper Imbalance	8
Sensitivity of Sheep to Copper	9
Factors Affecting the Copper Status in Sheep	10
Copper in the feed	10
Animal factor	13
Stress factors	14
Source of supplemental copper	14
Copper Antagonism Interactions	15
Iron antagonism	15
Zinc antagonism	16
Molybdenum antagonism	17
Sulphur antagonism	18
Thiomolybdate	18
Utilization of Antagonistic Interactions to Prevent Cu Toxicity	19
Copper Toxicity in Malaysia	20
<b>TRACE MINERALS AND REPRODUCTION</b>	<b>21</b>
<b>REPRODUCTIVE HORMONES</b>	<b>21</b>
Factors Affecting Basal Hormone Level	22
Exposure to ewe	22
Animal management	22



SERVING CAPACITY TEST	23
Factors Affecting Serving Capacity	24
Genetic factor	24
Social interaction	24
Age and experience	26
Seasonal variation	26
<b>III. MATERIALS AND METHODS</b>	<b>27</b>
EXPERIMENTAL SITE	27
MATERIALS	27
Experimental Animals	27
Dietary Treatment	28
Feeding and Care	29
METHODS	31
Familiarization and Training of Rams to Mount and Ejaculate	31
Serving Capacity Test	31
Semen Analysis Procedure	33
Blood Sampling and Analysis	39
Determination of Cu Levels in the Liver and Testis	40
Statistical Analysis	40
<b>IV. RESULTS</b>	<b>42</b>
SERVING CAPACITY	42
SEMEN QUALITY	50
COPPER STATUS	57
SERUM TESTOSTERONE LEVEL	60
<b>V. DISCUSSION</b>	<b>62</b>
SERVING CAPACITY	62
SEMEN QUALITY	66
COPPER STATUS	68
SERUM TESTOSTERONE LEVEL	71
<b>VI. CONCLUSION</b>	<b>72</b>
REFERENCE	75
APPENDICES	86
BIODATA OF THE AUTHOR	96



## LIST OF TABLES

	<b>Page</b>
Table 1: Mean weight of ram lambs at the start of the experiment	28
Table 2: Nutrient content of the dietary treatments	29
Table 3: Sexual performances of rams at the beginning and end of the experiment	44
Table 4: Monthly sexual performance of rams in the three dietary treatments	45
Table 5: Pearson correlation matrix for sexual performance of rams	49
Table 6: Semen quality of rams at the beginning and end of the experiment	51
Table 7: Monthly semen quality of ram in the three dietary treatment,	52
Table 8: Pearson correlation matrix for sperm quality parameters	56
Table 9: Plasma copper level of rams at the beginning and end of the experiment	58
Table 10: Liver and testis copper level of rams at the end of the experiment	58
Table 11: Monthly plasma copper level of rams fed on	59
Table 12: Total Livestock Population in Peninsular Malaysia (1970 – 2001)	86
Table 13: Nutritive value of PKC collected from literatures	87
Table 14: Chemicals used in wet digestion and their proportion	89
Table 15: Trend analysis for mating behavior over six months period over all treatments	90
Table 16: Trend analyses for sperm quality over four months period	90
Table 17: Proportion of abnormal sperm cells of rams subdivided into parts of sperm cells	91
Table 18: Mean monthly body weights of rams over the experimental period	92
Table 19: Calibration for EIAgen Testosterone Kit	93





## LIST OF FIGURES

	<b>Page</b>
Figure 1: Determination of sperm concentration	37
Figure 2: Determination of % sperm motility	37
Figure 3: Latency to first mount of rams over experimental period	46
Figure 4: Latency to first ejaculation of rams over experimental period	46
Figure 5: Refractory period of rams over experimental period	47
Figure 6: Number of mount of rams over experimental period	47
Figure 7: Number of ejaculation of rams over experimental period	48
Figure 8: Ram mating efficiency over experimental period	48
Figure 9: Sperm concentration of rams over four months	53
Figure 10: Proportion of abnormal sperm cells of rams over four months	53
Figure 11: Proportion of live sperm cells of rams over four months	54
Figure 12: Volume of ejaculates of rams over four months	54
Figure 13: Proportion of motile sperm cells of rams over four months	55
Figure 14: Serum testosterone profile of rams over experimental period	61
Figure 15: Monthly body weight change of rams	92
Figure 16: An overview of micro plate used for assaying serum testosterone	93



## LIST OF PLATES

	<b>Page</b>
Plate 1. Rams in a raised pen	30
Plate 2. Housing and feeding facility of rams	30
Plate 3. Ram exhibiting its serving potential in 30 min serving capacity test	33
Plate 4. Artificial vagina	38
Plate 5. Semen collection using artificial vagina	38



## ABBREVIATIONS

The following abbreviations are used in the thesis with or without definition.

ANOVA	Analysis of variance
ATP	Adenosine triphosphate
Ca	Calcium
CONC	Concentrate
CRD	Completely randomized design
Cu	Copper
Cu <sub>2</sub> O	Cuprous oxide
CV	Coefficient of variation
DI	Deciliter
DM	Dry matter
EIA	Enzyme immuno assay
EJA	Ejaculations
EMS	Error mean squares
Fe	Iron
GLM	General linear model
H	Hour
ICP	Inductively coupled plasma
K	Potassium
LFE	Latency to first ejaculation
LFM	Latency to first mount



LH	Lutienizing hormone
MEFF	Mating efficiency
Mo	Molybdenum
Mol	Moles
MPOB	Malaysian Palm Oil Board
ng	Nano gram
Nmol/l	Nano mole per liter
NRC	National research council
NS	Non significant
°C	Degree Celsius
OPF	Oil palm frond
p	Probability
PBS	Phosphate Buffer Saline
PKC	Palm kernel cake
ppm	Parts per million
r	Correlation coefficient
REF	Refractory period
RPM	Revolution per minute
S	Sulphur
SAS	Statistical analysis system
SD	Standard deviation
SE	Standard error
S-Mo	Sulphur molybdenum complex



t test	Students t-test
TCA	Trichloroacetic acid
UPM	Universiti Putra Malaysia
Zn	Zinc

# CHAPTER 1

## INTRODUCTION

The small ruminant production systems in Malaysia can be described as extensive, intensive, semi-intensive and animal tree-crop integration systems (Rajion *et al.*, 1993). Integration of ruminants with tree-crops is the most practical and widely practiced production system by many government and private farms. Integration of small ruminants, specifically sheep, in oil palm plantation has been successful (Rajion *et al.*, 1994), though the herbage under the plantation does not support the animals' maintenance requirement in quantity and quality.

Sheep population in Malaysia has showed an increasing trend for the last 30 years (Appendix A). This is due to their feeding habit, the ease to integrate them with oil palm or rubber plantations and the liberal importation of exotic breed and crossing or upgrading program (Abdullah Sani *et al.*, 1999).

The success of a sheep production enterprise is highly dependent, among others on the ability of rams to breed maximum number of ewes, which is a factor of libido and mating ability of the rams. Reproductive inefficiency in sheep results in reduced dissemination of superior genetic material and lower producer profitability. The existence of a large number of low libido rams in a



flock will have a negative impact on flock productivity and profitability by causing longer lambing interval and creating the need to keep many rams. This problem may be more pronounced in small farm households where it is uneconomical to keep less productive male animals for a long period.

Libido in sheep, as in all other animals, is influenced by factors such as genotype, nutrition, climate, health of animals and farm management practices. Nutritional manipulation/fluctuation, in particular trace elements, has a major influence in libido expression in animals (Martin and White, 1992; Puls, 1994). This can be induced through antagonistic interaction, toxicity and deficiency of the trace elements.

Among the trace elements, copper (Cu) has been reported to have a significant role in influencing the fertility of farm animals, being constituent of many enzymes and hormones (Davis and Mertz, 1987; Linder and Hazegh-Azam, 1996; Underwood and Suttle, 1999). Therefore, it is important to include high Cu feed source in animal rations. Palm kernel cake (PKC), a known by-product of the oil palm industry in Malaysia can be used in this regard as it has high Cu content. Palm kernel cake is a widely used feed supplement in Malaysia with considerable export potential (MPOB, 2002). The production of PKC by palm oil industry in Malaysia is increasing year after year. The production in 2000 was 1.3 million metric tones while that of 2001



was 1.8 million metric tones. The nutrient content of PKC makes it suitable for most ruminant animals. However, its Cu content makes it less suitable for sheep because a concentration as low as 10 ppm of Cu in the feed is said to be toxic to sheep (Church and Pond, 1988).

Previous research work have showed that Cu deficiency can cause infertility and delayed estrus in cattle and sheep (Underwood, 1977; Puls, 1994). Rams fed with Cu deficient diet were found to be less successful in breeding ewes (Weiner and Sales, 1976). This indicates how Cu is important for normal reproductive function of animals. However, information on the effect of excessive Cu on reproduction is lacking.

### **General Objective**

- To evaluate the effect of high level of palm kernel cake diet on serving capacity and semen quality of ram lambs.

### **Specific Objectives**

1. To investigate the effects of PKC (higher Cu diet) on serving capacity and sperm quality of ram lambs.
2. To investigate the effect of PKC (high Cu diet) on plasma Cu and serum testosterone levels in ram lambs





3. To investigate the controlling effect of Molybdenum (Mo) and Sulphur (S) supplemented to PKC based diet on plasma, liver and testis Cu level in ram lambs.