



UNIVERSITI PUTRA MALAYSIA

**INFLUENCE OF GLOMUS MOSSEASE AND NITROGEN
FERTILIZATION ON GROWTH AND YIELD OF SWEET POTATO
(IPOMOEA BATATAS L.)**

YASSIN MOHAMED AHMED.

FP 2005 22

**INFLUENCE OF *Glomus mosseae* AND NITROGEN FERTILIZATION ON
GROWTH AND YIELD OF SWEET POTATO (*Ipomoea batatas* L.)**

By

YASSIN MOHAMED AHMED

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of Master of Science**

June 2005



DEDICATION

This work is dedicated To

My beloved young daughter Tasabeeh Yassin, who missed me for long time during this research,

To

The departed soul of my mother,

To

My small family members:

Tahani Yassin

Abdul khalig Yassin

Ammar Yassin

Zahra Taha



Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

INFLUENCE OF *Glomus mosseae* AND NITROGEN FERTILIZATION ON GROWTH AND YIELD OF SWEET POTATO (*Ipomoea batatas* L.)

By

YASSIN MOHAMED AHMED

June 2005

Chairperson : Associate Professor Radziah Othman, PhD

Faculty : Agriculture

Application of arbuscular mycorrhizae (AM) in sweet potato production can contribute to reduction of input of chemical fertilizers in a sustainable agriculture. Pot and field experiments were conducted to determine the effect of AM inoculum level, and AM inoculation with different rates of nitrogen, on growth and yield of sweet potato. A pot experiment was conducted under glasshouse conditions using Sepang Oren sweetpotato cultivar grown in unsterilized Bungor series soil, and inoculated with 0, 20, 40 and 80 g plant⁻¹ of AM inoculum, consisting of *Glomus mosseae* UK118. The soil was amended with organic compost JITU Biofertilizer ® at rate of 20 t ha⁻¹, and inorganic fertilizer N, P, and K at the rate of 30 kg N, 60 kg P₂O₅ and 100 kg K₂O ha⁻¹, respectively. Results showed that AM inoculum level significantly (p<0.05) influenced the root colonization, but not the storage root yield. Application of 40 g plant⁻¹ *G. mosseae* inoculum significantly increased the root infection, spore number and P concentration in shoot. Different levels of AM applied



showed no significant difference on storage root yield, starch content and nutrient concentrations.

A field experiment was conducted to assess the effect of AM inoculation and nitrogen fertilization rates on growth and yield of sweetpotato. Experimental plots were prepared in the UPM Integrated Farm and three levels of nitrogen fertilizer used were 0, 60 and 120 kg ha⁻¹ in the form of urea in combination with two levels of AM inoculation. Inorganic fertilizers were applied at the rate of 60 kg P₂O₅ and 100 kg K₂O ha⁻¹. Sweetpotato was harvested at 120 days of growth. Results showed that application of nitrogen significantly ($P < 0.05$) influenced the yield of sweetpotato. Storage root yield increased with increasing nitrogen fertilizer, and application of 60 kg N ha⁻¹ and 120 kg N ha⁻¹ showed storage roots yield increase of 54.1% and 83.6% respectively, compared to control treatment. The present study showed no significant effect of *G. mosseae* on sweetpotato yield, plant nutrient uptake and soil nutrient concentrations. However, the mycorrhizal inoculation significantly ($p < 0.05$) influenced the root colonization and AM spore number in the soil. The highest root colonization of 44.36% was in AM inoculated plant with 120 kg N ha⁻¹ fertilization rate. The highest spore number of 46 spore 10 g⁻¹ soil was observed in AM inoculated treatment without N fertilization.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains

**PENGARUH *Glomus mosseae* DAN BAJA NITROGEN TERHADAP
PERTUMBUHAN DAN HASIL KELEDEK (*Ipomoea batatas* L.)**

Oleh

YASSIN MOHAMED AHMED

June 2005

Pengerusi : Profesor Madya Radziah Othman, PhD

Fakulti : Pertanian

Aplikasi mikoriza arbuskul (MA) dalam pengeluaran ubi keledek boleh menyumbang kepada pengurangan input baja kimia dalam pertanian lestari. Eksperimen berpasu dan lapangan telah dijalankan untuk menilai kesan aras inokulum MA dan kesan inokulasi MA serta kadar berbeza baja nitrogen terhadap pertumbuhan dan hasil ubi keledek. Satu kajian berpasu telah dijalankan dalam rumah kaca menggunakan ubi keledek kultivar Sepang Oren pada tanah siri Bungor tidak steril dengan inokulasi 0, 20, 40 dan 80 g inokulum MA *Glomus mosseae* UK 118 bagi setiap pokok. Tanah dirawat dengan kompos organik Baja bio JITU® pada kadar 20 t ha⁻¹ dan baja tak organik N, P, dan K masing-masing pada kadar 30 kg N, 60 kg P₂ O₅ dan 100 kg K₂O ha⁻¹. Keputusan menunjukkan paras inokulum MA memberi kesan signifikan terhadap kolonisasi akar tetapi tidak pada hasil ubi keledek. Aplikasi dengan 40 g inokulum *G. mosseae* untuk setiap pokok memberi



kesan signifikan terhadap peningkatan jangkitan akar, bilangan spora dan kadar P dalam daun. Perbezaan paras inokulum MA juga menunjukkan tiada perbezaan yang signifikan ke atas hasil ubi keledek, kandungan kanji dan kandungan nutrien.

Satu kajian di lapangan telah dijalankan untuk mengenalpasti kesan inokulasi MA dan kadar pembajaan N pada pertumbuhan dan hasil keledek. Plot eksperimen telah disediakan di ladang bersepadu UPM Serdang dan tiga kadar baja N telah di gunakan iaitu 0, 60 dan 120 kg ha⁻¹ dalam bentuk urea dengan kombinasi dua paras inokulasi MA. Baja tak organik telah diaplikasi pada kadar 60 kg P₂O₅ dan 100 kg K₂O ha⁻¹. Ubi keledek dituai selepas 120 hari pertumbuhan. Kajian mendapati paras pembajaan N memberi kesan signifikan terhadap hasil ubi keledek. Hasil ubi keledek meningkat dengan peningkatan kadar baja N dan pemberian 60 kg dan 120 kg N ha⁻¹ menunjukkan peningkatan hasil sebanyak 54.1% dan 83.6% berbanding kawalan. Kajian ini menunjukkan *G. mosseae* tidak memberi kesan signifikan terhadap hasil keledek, kadar pengambilan nutrien dalam tanaman dan kandungan nutrien dalam tanah. Walau bagaimanapun, inokulasi mikoriza memberi kesan signifikan terhadap kolonisasi akar dan bilangan spora MA dalam tanah. Mikoriza arbuskul memberi kesan ketara ($P < 0.05$) terhadap kolonisasi akar dan jumlah spora dalam tanah. Peratusan kolonisasi tertinggi 44.36% didapati pada rawatan 120 kg N ha⁻¹ dengan MA. Bilangan spora tertinggi 46 spora 10 g⁻¹ tanah didapati dalam rawatan mikoriza tanpa pembajaan N.

ACKNOWLEDGEMENTS

Alhamdu lillah with whose permission good deeds are completed.

I would like to express my most sincere gratitude to my supervisor Assoc. Prof. Dr. Radziah Othman, chairperson of my supervisory committee, who provided my scholarly insights, professional guidance, moral support and motivation throughout my study. My sincere thanks are also due to my supervisory committee member; Assoc. Prof. Dr. Mohd Said Saad for his invaluable advice and support.

Sincere thanks and appreciation is to be extended to Prof. Dr. Azizah Hashim, my former supervisory committee member for her guidance and endless support. My special thanks to Dr. Lutfi, Prof. Dr. Zulkifli Hj. Shamsudin, Dr. Samsuri A. Wahid, Dr. Ahsanul Haque, Mr. Abdul Galeel and Dr. Osmanu Haruna for the assistant with the statistical analysis of the data. Special thanks to Prof. Dr. Mohammed Omer my external examiner, Assoc. Prof. Dr. Anuar Abd Rahim my chairman during the viva, Assoc. Prof. Dr. Halimi Mohd saud and Assoc. Prof. Dr. Mohd Razi Ismail my internal examiners.

I would like to thank all the staff of Department of Land Management and particularly; Mr. Jamil, Mr. Ramli, Mr. Rahim, Puan Fauziah, Mr. Elias, Mr. Abdul Aziz, Mr. Ariffin, Mr. Zulkifli and Puan Zarina. My thanks also to Mr. Johari and Mr. Nazri in Plant Pathology laboratory, Puan Asiah in Plant Physiology laboratory, for their help and assistance.



I would like also to extend my thanks to the staff of the Integrated Farm Universiti Putra Malaysia for their cooperation and assistance during the field experiment and special thanks to Mr. Osman Salleh, Mr. Shahar, Mr. Ali Rashid, Velasamy and his friends, and all the staff of the farm. My thanks to all my colleagues the postgraduate students; Tengku Sabrina, Mukhlis, Farzana, Illani, Maliza, Zakri, Aini and Negash, for their kind assistance and cooperation.

Special Thanks to the Ministry of Science, Technology and Innovation (MOSTI) for the financial support IRPA project NO 01-02-04-0097-EA001.



I certify that an Examination committee met on 24/06/2005 to conduct the final examination of Yassin Mohamed Ahmed on his Master of Science thesis entitled “Influence Of *Glomus mosseae* and Nitrogen Fertilization on Growth and Yield of Sweet Potato (*Ipomoea batatas* L.)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universti Pertanian (Higher Degree) Regulation 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

ANUAR ABD RAHIM, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

MOHD RAZI ISMAIL, PhD


Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

HALIMI MOHD SAUD, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

MOHAMMED OMAR, PhD

Professor
Monash University
Malaysia
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD

Professor /Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **21 JUL 2005**



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

RADZIAH OTHMAN, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

MOHD SAID SAAD, PhD
Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)




AINI IDERIS, PhD
Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11 AUG 2005



DECLARATION

I hereby declare that the thesis is based on my original work except for quotation and citation, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



YASSIN MOHAMED AHMED

Date: 16/08/05

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL	viii
DECLARATION	x
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF PLATES	xvii
LIST OF ABBREVIATIONS	xviii
CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	5
Sweet Potato	5
Climate	6
Growth Cycle	7
The Root System	7
Storage root	8
Shoot System	10
Tillage and Seedbed Preparation	11
Effect of Nitrogen Fertilization on Sweetpotato	15
Mycorrhizae	16
Mycorrhizal Symbioses	16
Types of Mycorrhiza	17
Arbuscular Mycorrhiza (AM)	18
Benefits of Arbuscular Mycorrhiza	22
Factors Affecting AM Activity	26
Beneficial Interaction with PGPR	29
Resouce Competition	29
AM Inoculation	30
Inoculum Density	30
Application of AM on Sweetpotato	31



III	EFFECT OF <i>Glomus mosseae</i> INOCULUM DENSITY ON GROWTH AND YIELD OF SWEETPOTATO	
	Introduction	32
	Materials and Methods	33
	Soil Preparations	33
	Planting Materials	34
	AM Inoculation and Planting	34
	Shoot, Root and Storage root weight	35
	Plant Tissue Analysis	35
	AM Root Colonization	36
	AM Spore Count	36
	Soil pH and Nutrients Analysis	37
	Statistical Analysis	37
	Results	38
	Plant Growth	38
	Sweetpotato Yield	38
	AM Root Colonization and Spore Count	39
	Nutrient Concentration in Shoots	42
	Nutrient Uptake in Shoots	42
	Nutrient and Starch Concentration of Storage root	43
	Nutrient Content of Storage root	44
	Soil pH and Nutrient Concentration	44
	Correlation Analysis	45
	Discussion	47
	Conclusion	51
IV	EFFECT OF <i>Glomus mosseae</i> INOCULATION AND NITROGEN FERTILIZATION RATE ON GROWTH AND YIELD OF SWEETPOTATO UNDER FIELD CONDITION	
	Introduction	52
	Materials and Methods	54
	Experiment Location	54
	Field Preparation and Soil Analysis	54
	Planting Material	55
	AM Inoculation	55
	Inorganic Fertilization	56
	Harvesting above Ground Biomass and Storage root Yield	56
	Nutrient Concentration in Shoot and Storage root	57
	Storage root Starch and Crude Protein Content	57
	Arbuscular Mycorrhizal Root Colonization	58
	Arbuscular Mycorrhizal Spore Count	58
	Soil pH and Nutrient Concentration	59
	Statistical Analysis	59
	Results	60
	Above Ground Biomass	60
	Storage root Yield	61



Sweetpotato Production	62
Marketable Yield	62
Nutrient Concentration in Shoots	64
Nutrient Uptake in Shoot	65
Nutrient, Starch and Crude Protein Concentration of Storage root	65
Nutrient Content of Storage root	67
Root Colonization and Spore Count	68
Soil pH and Nutrient Concentration	69
Discussion	71
Conclusion	76
V GENERAL DISCUSSION AND CONCLUSION	77
REFERENCES	83
APPENDICES	96
BIODATA OF THE AUTHOR	106



LIST OF TABLES

Table		Page
1	Fungal and host taxonomy of mycorrhizae.	17
2	Characterisation of arbuscular mycorrhiza.	18
3	Effect of AM inoculum density on sweetpotato growth and root/shoot ratio.	38
4	Effect of AM inoculum density on sweetpotato storage root yield and diameter.	39
5	Effect of AM inoculum density on root colonization and spore count.	40
6	Effect of AM inoculum density on nutrient concentration in shoot.	42
7	Effect of AM inoculum density on the uptake of N, P, K, Ca and Mg in shoot	43
8	Effect of AM inoculum density on nutrients and starch concentration of storage root.	43
9	Effect of AM inoculum density on storage root N, P, K, Ca and Mg content (mg plant ⁻¹).	44
10	Effect of AM inoculum density on soil pH and nutrient concentration.	45
11	Correlation between yield, root colonization (RC), starch content (SC), storage root N (St.N), Storage root P (St.P), shoot N (Sh. N), shoot P (Sh. P), Soil N (S.N) and soil P (S.P).	46
12	Effect of AM inoculation and N fertilization rate on shoot, storage root dry weight and shoot to root ratio.	60
13	Effect of AM inoculation and N fertilization on, total storage root yield, sweetpotato production and marketable grade storage root.	63
14	Arbuscular mycorrhiza inoculated and nitrogen-fertilized sweetpotato shoot nutrient concentration (%).	64



15	Arbuscular mycorrhiza inoculated and nitrogen-fertilized sweetpotato shoot nutrient uptake(g plant ⁻¹).	65
16	Effect of AM inoculation and N fertilization on nutrient, starch and crude protein concentration(%) of storage root.	66
17	Effect of AM inoculation and N fertilization on nutrient content of storage root (g plant ⁻¹).	68
18	Effect of AM inoculation and N fertilization on root colonization and spore number in the soil.	69
19	Effect of AM inoculation and N fertilization on soil pH and nutrient concentration.	70



LIST OF FIGURES

Figure		Page
1	Types of roots in the sweetpotato plant.	9
2	Parts inside the storage roots.	9
3	Relationship between AM inoculum density and root colonization (%)	41
4	Relationship between AM inoculum density and spore number in soil.	41



LIST OF PLATES

Figure		Page
1	Arbuscular mycorrhiza vesicles in infected sweetpotato roots.	40
2	<i>Glomus mosseae</i> spores isolated by wet sieving and decanting method.	59
3	Storage root yield of sweetpotato in different N fertilizer rate without AM inoculation.	61
4	Storage root yield of sweetpotato in different N fertilizer rate with AM inoculation.	61



LIST OF ABBREVIATIONS

AM	: Arbuscular Mycorrhiza.
AMF	: Arbuscular Mycorrhiza Fungi.
ANOVA	: Analysis of Variance.
Ca	: Calcium.
CIP	: International Potato Center.
CRD	: Completely Randomized Design.
DOA	: Department of Agriculture.
ECM	: Ectomycorrhiza.
K	: Potassium.
LSD	: Least Significant Difference.
MAFF	: Ministry of Agriculture, Food and Fisheries.
Mg	: Magnesium.
MOP	: Muriate of Potash.
N	: Nitrogen.
P	: Phosphorus.
PGPR	: Plant Growth Promoting Rhizobacteria.
PSB	: Phosphate Solubilizing Bacteria.
R/S	: Root to shoot.
RCBD	: Randomized Complete Block Design.
TSP	: Triple Super Phosphate.
VAM	: Vesicular Arbuscular Mycorrhiza.
Zn	: Zinc.



CHAPTER I

INTRODUCTION

Sweetpotato *Ipomoea batatas* (L.) is ranked the fifth among the world most important food crops, with more than 133 million tones of annual production (CIP, 2005). Sweetpotato is currently grown in more than 100 countries. Most of the producing nations are situated in the tropical developing world. The increasing recognition of potential of sweetpotato has resulted intensified research to enhance its production and consumption as food for the world (Woolfe, 1992). In Malaysia sweetpotato is the second most important root crop ranked after cassava. At present utilization possibilities are somewhat limited mainly to home use and modest snack industry (Tan and Mooi, 1997). It has the potential of being transformed from minor food crop into a commercial crop. Sweetpotato is one of the cash crops with total area of 1,413.7 hectares and total production of 15,946.5 metric tones; most producing states are Johor, Perak and Kelantan (DOA, 2005).

Main problems of sweetpotato production in Malaysia are the yield decline of sweetpotato cultivars (Mooi and Tan, 2001) lack of good soil, low market price of sweetpotato yield, small hectareage and lack of mechanization of production operations (Saad, 1994). It is essential to increase sweetpotato production as a potential food, but the use of high chemical fertilizers to increase the production lead to environmental problems and contamination in land and water resources and finally deterioration of the product quality. Therefore, due to the public concern, there is

increasing interest in the application of biofertilizers in agriculture. A biofertilizer such as mycorrhizal fungi can keep the balance between the high production of important crops, natural resources and environment.

Mycorrhizae are symbiotic associations between the roots of most plant species and soil fungi. Bi-directional movement of nutrients characterizes this symbiosis where carbon flows to the fungus and inorganic nutrients move to the plant thereby providing a critical linkage between plant root and soil. Mycorrhizal associations between fungus and plant root are ubiquitous in the natural environment (Smith and Read, 1997). The arbuscular mycorrhizal association is the most ancient and most common and probably aided the first land plants (Simon et al., 1993).

Mycorrhiza may give an advantage to the plant group with a tendency to efficiently exploit a given soil volume by accessing nutrients outside of the roots' nutrient depletion zone (Campbell et al., 1991). Arbuscular-mycorrhizal fungi (AMF) form symbiotic associations with most of the tropical crop plants. This fungi can improve plant growth under conditions of low fertility, confer resistance and/or tolerance of some pathogens, improve the water balance of plants and contribute to the formation of soil structure (Harikumar and Potty, 2002).

Tropical crops, such as cassava, sweetpotato and cowpea are often heavily colonized by arbuscular mycorrhiza AM fungi under natural conditions. Environmental conditions affect the formation of AM or influence the extent of root colonization.

The natural inoculum concentration of AM fungi in the soil also determines the extent of root infection (Sieverding, 1991). The quantity of mycorrhizae formed by species of AM fungi in field soils will depend on both the amount of infective hyphae and their relative competitive ability (Abbott and Robson, 1984). The length of root colonized by AM fungi at early stages of root growth is related to inoculum density (Abbott and Robson, 1984; Wilson, 1984; Giovannetti and Avio, 1986). The difference in competitive ability to form mycorrhizae may be based on antagonism (Hepper et al., 1988). Depletion of nutrients within roots (Wilson and Trinick, 1983) and carbon demand (Pearson et al., 1993).

Nitrogen is one of the most limiting elements for the plant production in tropical soils. It is the most frequently applied fertilizer in the tropics and often the only fertilizer element added to the soil. The most common nitrogen fertilizer source is urea. In the soil, urea is converted to ammonium. Ammonium fertilizer sources were reported to inhibit mycorrhizal growth more than nitrate sources (Chambers et al., 1980). However, the effects of nitrogen application on vesicular arbuscular mycorrhiza (VAM) are not consistent. They vary from one soil site to another and may depend on the availability of P (Sieverding, 1991). The uptake of many ions by plants, including NO_3^- seems to be controlled by specific demand driven mechanisms (Imsande and Touraine, 1994).

High application rates of inorganic fertilizers especially nitrogen cause pollution of water and land resources, soil acidification and increased production cost. Many



studies were conducted on the AM application in agriculture, but most of the studies were on other crops. Currently there are insufficient studies on the AM application on root crops in general and specifically on sweetpotato as one of the world's potential food crops.

In view of the benefits of AM on plant growth the following studies were conducted to determine the effect of arbuscular mycorrhizal fungi (*Glomus mosseae*) inoculum density on growth and yield of sweetpotato under glasshouse condition, and to assess the effect of AM inoculation and nitrogen fertilization rate on growth and yield of sweetpotato under field conditions.

CHAPTER II

LITERATURE REVIEW

Sweetpotato

Sweetpotato *Ipomoea batatas* (L.) is ranked the fifth among the world most important food crops, with more than 133 million tonnes of annual production (CIP, 2005). It is an ancient food plant of tropical America and Pacific Islands, and several varieties of sweetpotato are now extensively cultivated in many parts of the world (Winaro, 1982). Sweetpotato is the second most important root crop in Malaysia ranked after cassava (*Manihota esculent* Crantz) grown mainly as a cash crop or subsistence crop, in small farm or home garden for the fresh food market or home consumption. Sweetpotato production in Malaysia is 15,946.5 metric ton in an area of 1,413.7 hectare, with average yield of 11 t ha⁻¹ (DOA, 2005). Some sweetpotato varieties planted in Malaysia are UPMSS5 and Ubi Biru (Saad, 1994), Serdang1, Empat Bulan, Large While, Bukit Naga (Anon, 1990), Gendut, Telong, Jalomas, Kuala Bikam and Susu Lembu (Tan, 2000).

Sweetpotato is a major crop that feeds millions of people in developing world. It is especially popular among farmers with limited resources, and produces more biomass and nutrients per hectare than any other food crop in the world (Prakash, 1994). Sweetpotato is adaptable to a broad range of agro-ecological conditions and fits in low input agriculture. It is in many ways an ideal crop for farmers as it grows on low nitrogen soils, tolerates drought well, crowds out weeds, suffers relatively from few pests, and is highly productive even under adverse farming conditions. Sweetpotato is grown in more than 100 countries as a valuable source