



UNIVERSITI PUTRA MALAYSIA

**CHEMICAL CONSTITUENTS FROM ANNONA MURICATA (LINN.)
AND GARCINIA MANGOSTANA (LINN.) AND THEIR BIOLOGICAL
ACTIVITIES**

SHAARI BIN DAUD.

FS 2005 31

**CHEMICAL CONSTITUENTS FROM
ANNONA MURICATA (LINN.) AND *GARCINIA MANGOSTANA* (LINN.)
AND THEIR BIOLOGICAL ACTIVITIES**

By

SHAARI BIN DAUD

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

April 2005



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**CHEMICAL CONSTITUENTS FROM
ANNONA MURICATA (LINN.) AND *GARCINIA MANGOSTANA* (LINN.)
AND THEIR BIOLOGICAL ACTIVITIES**

By

SHAARI BIN DAUD

April 2005

Chairman : Associate Professor Gwendoline Ee Cheng Lian, PhD

Faculty : Science

Chemical studies were carried out on *Annona muricata* L. (Annonaceae) and *Garcinia mangostana* L. (Guttiferae). The chemical investigations covered acetogenin, triterpenoids and xanthenes. These compounds were isolated using common chromatographic techniques and were identified by using spectroscopic technique such as NMR, MS, IR and UV.

Biological activities for both plants were also evaluated for their antimicrobial activities against four pathogenic bacteria and four fungi, cytotoxic activity against CEM-SS and HL-60 cell lines and larvicidal activity against *Aedes aegypti*. Larvicidal activity data have never been reported before.



Annona muricata L. (stem bark) have afforded one acetogenin, solamin and two triterpenoids, stigmasterol and sitosterol. Meanwhile studies on *Garcinia mangostana* L. (bark) have provided a new xanthone, mangosharin (2,8-dihydroxy-6-methoxy-5-(3-methylbut-2-enyl)-xanthone) together with six other xanthones, α -mangostin, β -mangostin, garcinone D, xanthone 1, 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone and mangostanol and one triterpenoid, friedelin.

Cytotoxic tests were performed using CEM-SS cell line and HL-60 cell line. The crude hexane extract of *Annona muricata* L. gave a significant activity with an IC₅₀ value of 0.8 μ g/ml against CEM-SS cell line while the crude ethyl acetate (EA) extract also gave a significant activity with an IC₅₀ value of 0.5 μ g/ml but against HL-60 cell line. The crude hexane extract of *Garcinia mangostana* L. is also considered to be active against the CEM-SS cell-line with an IC₅₀ value of 17 μ g/ml. Other than that, the pure compounds α -mangostin, mangostanol and garcinone D also gave significant activities with IC₅₀ values of 5.5, 9.6 and 3.2 μ g/ml against CEM-SS cell line, respectively.

The antimicrobial activity test was also carried out using four pathogenic bacteria, namely, Methicillin Resistant *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *Bacillus subtilis*. However, most of the crude extracts gave only moderate or weak activity. The pure compound α -mangostin gave weak activity against *Salmonella typhimurium* and medium inhibition against *Bacillus subtilis*.



The larvicidal tests were performed against the larvae of *Aedes aegypti* using the WHO (1981) standard procedures with slight modifications. The crude hexane and ethyl acetate extracts of *Annona muricata L.* showed moderate toxicity towards the larvae by giving LC₅₀ values of less than 100 µg/ml. The crude hexane and ethanol extracts of *Garcinia mangostana L.* gave weak activities against the larvae with LC₅₀ values of more than 150 µg/ml while the crude ethyl acetate extract showed moderate activity with an LC₅₀ value of 30.1 µg/ml. Larvacidal activity on the pure compound, α-mangostin gave a LC₅₀ value of 19.4 µg/ml.

No activity was recorded for the antifungal activity tests.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**KANDUNGAN KIMIA DAN AKTIVITI BIOLOGI DARIPADA
ANNONA MURICATA (LINN.) DAN *GARCINIA MANGOSTANA (LINN.)***

Oleh

SHAARI BIN DAUD

April 2005

Pengerusi : Profesor Madya Gwendoline Ee Cheng Lian, PhD

Fakulti : Sains

Kajian kimia telah dijalankan terhadap *Annona muricata L.* (Annonaceae) dan *Garcinia mangostana L.* (Guttiferae). Kajian kimia terperinci merangkumi jenis sebatian seperti asetogenin, tritepenoid dan xanthone. Sebatian-sebatian ini diasingkan dengan menggunakan pelbagai teknik kromatografi dan dikenalpasti dengan menggunakan teknik spektroskopi seperti NMR, MS, IR dan UV.

Aktiviti biologi untuk kedua-dua tumbuhan juga dinilai untuk aktiviti microbial terhadap empat bacteria dan empat fungi, aktiviti sitotoksik terhadap sel CEM-SS dan sel HL-60 dan aktiviti larva terhadap *Aedes aegypti*. Data untuk aktiviti larva tidak pernah direkodkan sebelum ini.

Annona muricata L. (kulit batang) telah menghasilkan satu asetogenin, solamin dan dua tritepenoid, stigmasterol dan sitosterol. Sementara itu, kajian terhadap kulit pokok

Garcinia mangostana L. telah memberikan satu xanthone baru, mangoshaarin (2,8-dihydroxy-6-methoxy-5-(3-methylbut-2-enyl)-xanthone bersama dengan enam xanthone yang lain, α -mangostin, β -mangostin, garcinone D, xanthone 1, 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone dan mangostanol dan satu triterpenoid, friedelin.

Ujian sitotoksik telah dijalankan dengan menggunakan sel CEM-SS dan HL-60. Ekstrak mentah heksana dari *Annona muricata L.* memberikan aktiviti yang kuat dengan nilai IC_{50} 0.8 $\mu\text{g/ml}$ terhadap sel CEM-SS manakala ekstrak mentah etil asetat juga menunjukkan aktiviti yang kuat dengan nilai IC_{50} 0.5 $\mu\text{g/ml}$ tetapi terhadap sel HL-60. Ekstrak heksana mentah dari *Garcinia mangostana L.* dianggap sebagai aktif terhadap sel CEM-SS dengan nilai IC_{50} 17 $\mu\text{g/ml}$. Selain daripada itu tiga sebatian tulen iaitu α -mangostin, mangostanol dan garcinone D menunjukkan aktiviti yang kuat dengan nilai IC_{50} 5.5, 9.6 dan 3.2 $\mu\text{g/ml}$ masing-masing terhadap sel CEM-SS.

Aktiviti antimikrob dijalankan dengan menggunakan empat jenis bakteria iaitu Methicillin Resistant *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *Bacillus subtilis*. Bagaimanapun, kebanyakan ekstrak mentah yang di uji dengan bakteria-bakteria ini hanya memberikan keaktifan yang sederhana dan lemah. Sebatian tulen α -mangostin menunjukkan aktiviti yang lemah terhadap bakteria *Salmonella typhimurium* dan sederhana terhadap *Bacillus subtilis*.

Ujian larva telah dijalankan dengan menggunakan larva jenis *Aedes aegypti* mengikut prosedur-prosedur piawai WHO (1981) dengan sedikit pengubahsuaian. Ekstrak mentah heksana dan etil asetat daripada *Annona muricata L.* menunjukkan toksik yang sederhana terhadap larva dengan memberi nilai LC_{50} kurang daripada 100 $\mu\text{g/ml}$. Ekstrak mentah heksana dan etanol daripada *Garcinia mangostana L.* memberikan aktiviti yang lemah terhadap larva dengan nilai LC_{50} lebih daripada 150 $\mu\text{g/ml}$ manakala ekstrak mentah etil asetat menunjukkan toksik yang sederhana dengan nilai LC_{50} 30.1 $\mu\text{g/ml}$. Aktiviti larva pada sebatian tulen, α -mangostin memberi nilai LC_{50} 19.4 $\mu\text{g/ml}$.

Tiada sebarang aktiviti yang direkodkan untuk ujian antifungi.

ACKNOWLEDGEMENTS

I would like to express my sincere and deepest appreciation to my dedicated supervisor, Assoc. Prof. Dr. Gwendoline Ee Cheng Lian for her constant advice and guidance throughout this project. My sincere and deepest gratitude are also extended to my supervisory committee member Prof. Dr. Mawardi bin Rahmani for his support. Financial support from the Malaysian government under IRPA programme is also greatly appreciated.

I also wish to thank my labmates Fido, Aza, Sheikh, Lim, Audrey, Sooi Kim, Cheow and Gin Keat for sharing the useful advices throughout the course of my research. My special thanks go to all the staff of the Chemistry Department of UPM especially Mr. Zainal Abidin Kassim, Mr. Johadi Iskandar and Mrs. Rusnani Amirudin for helping me with MS, NMR and IR spectral data.

Last but not least, I am also indebted to my beloved family and friends whose encouragement and support have contributed towards the success of this research.



I certify that an Examination Committee met on 18 April 2005 to conduct the final examination of Shaari bin Daud on his Master of Science thesis entitled "Chemical Constituents From *Annona muricata* (Linn.) and *Garcinia mangostana* (Linn.) and Their Biological Activities" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Dr. Abdul Halim Abdullah, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Dr. Taufiq Yap Yun Hin, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Dr. Mohd. Aspollah Sukari, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Dr. Ibrahim Jantan, PhD

Professor
Faculty of Allied Health Sciences
Universiti Kebangsaan Malaysia
(Independent Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/ Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 JUL 2005



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

GWENDOLINE EE CHENG LIAN, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

MAWARDI BIN RAHMANI, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.
Professor/ Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11 AUG 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



SHAARI BIN DAUD

Date: 20.7.2005



TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL SHEETS	ix
DECLARATION FORM	xi
LIST OF TABLES	xiv
LIST OF FIGURES	xvi
LIST OF ABBREVIATIONS	xxiii
CHAPTER	
1 INTRODUCTION	1
1.1 General Introduction	1
1.2 Objectives of Study	3
2 LITERATURE REVIEW	4
2.1 Botany of plants studied	4
2.2.1 The family Annonaceae	4
2.2.2 The genus <i>Annona</i>	4
2.2.3 The species <i>Annona muricata</i>	5
2.1.4 The family Guttiferae	6
2.1.5 The genus <i>Garcinia</i>	7
2.1.6 The species <i>Garcinia mangostana</i>	7
2.2 Chemical constituents	8
2.2.1 Chemistry of <i>Annona</i> species	8
2.2.2 Annonaceous acetogenins	9
2.2.2.1 Previous Work on annonaceous acetogenins in <i>Annona</i> species	11
2.2.3 Chemistry of <i>Garcinia</i> species	21
2.2.4 Previous Work on xanthenes in <i>Garcinia</i> species	21
2.2.5 Previous Work on flavonoids in <i>Garcinia</i> species	31
2.2.6 Previous Work on benzophenone in <i>Garcinia</i> species	33
2.2.7 Previous Work on terpenes in <i>Garcinia</i> species	35
2.3 Biological activities	36
2.3.1 Previous Work on Bioassay of <i>Annona</i> species	36
2.3.2 Previous Work on bioassay of <i>Garcinia</i> species	40
3 EXPERIMENTAL	44
3.1 Plant Material	44
3.2 Instruments	44
3.2.1 Infrared Spectroscopy (IR)	44
3.2.2 High Performance Liquid Chromatography (HPLC)	45



3.2.3	Mass Spectra (MS)	44
3.2.4	Melting Point	45
3.2.5	Nuclear Magnetic Resonance (NMR)	45
3.2.6	Ultra Violet (UV)	45
3.3	Chromatographic Methods	46
3.3.1	Column Chromatography	46
3.3.2	Thin Layer Chromatography (TLC)	46
3.3.3	Preparative Layer Chromatography	47
3.4	Dyeing Reagents for TLC	48
3.4.1	Vanillin-Sulfuric Acid solution	48
3.4.2	Iron (III) Chloride solution	48
3.5	Extraction and isolation of compounds from <i>Annona muricata L.</i> and <i>Garcinia mangostana L.</i>	49
3.5.1	<i>Annona muricata L.</i>	49
3.5.2	<i>Garcinia mangostana L.</i>	54
3.6	Cytotoxic assay	66
3.7	Antimicrobial activity	67
3.8	Larvicidal assay	68
4	RESULTS AND DISCUSSION	70
4.1	Isolation of Chemical Constituents from <i>Annona muricata L.</i>	70
4.1.1	Characterization of stigmasterol (111)	72
4.1.2	Characterization of solamin (22)	79
4.1.3	Characterization of sitosterol (112)	91
4.2	Isolation of Chemical Constituents from <i>Garcinia mangostana L.</i>	98
4.2.1	Characterization of friedelin (113)	100
4.2.2	Characterization of α -mangostin (64)	108
4.2.3	Characterization of β -mangostin (61)	119
4.2.4	Characterization of garcinone D (114)	130
4.2.5	Characterization of xanthone 1(115)	143
4.2.6	Characterization of 1,6-dihydroxy-3,7-dimethoxy -2-(3-methylbut-2-enyl)-xanthone (116)	155
4.2.7	Characterization of mangostanol (68)	166
4.2.8	Characterization of mangosharin (117)	180
4.3	Bioassay Results	195
4.3.1	Cytotoxic Activity	195
4.3.2	Antimicrobial Activity	198
4.3.3	Larvicidal Activity	200
4.3.4	Antifungal Activity	201
5	CONCLUSIONS	203
	BIBLIOGRAPHY	204
	APPENDICES	211
	BIODATA OF THE AUTHOR	223



LIST OF TABLES

Table		Page
4.1	^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3) Assignments of stigmasterol (111)	74
4.2	^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3) Assignments of solamin (22)	81
4.3	^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3) Assignments of sitosterol (112)	93
4.4	^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3) Assignments of friedelin (113)	102
4.5	^1H NMR (400 MHz, CD_3OD), ^{13}C NMR (100 MHz, CD_3OD), COSY and HMBC. Assignments of α -mangostin (64)	110
4.6	^1H NMR (400 MHz, CDCl_3), ^{13}C NMR (100 MHz, CDCl_3), COSY and HMBC. Assignments of β -mangostin (61)	121
4.7	^1H NMR (400 MHz, $\text{Me}_2\text{CO}-d_6$) and ^{13}C NMR (100 MHz, $\text{Me}_2\text{CO}-d_6$) of garcinone D and (114)	133
4.8	^1H NMR (400 MHz, $\text{Me}_2\text{CO}-d_6$), ^{13}C NMR (100 MHz, $\text{Me}_2\text{CO}-d_6$), COSY, and HMBC. Assignments of garcinone D (114)	134
4.9	^1H NMR (400 MHz, $\text{Me}_2\text{CO}-d_6$) and ^{13}C NMR (100 MHz, $\text{Me}_2\text{CO}-d_6$) of Xanthone 1 and (115)	145
4.10	^1H NMR (400 MHz, CDCl_3), ^{13}C NMR (100 MHz, CDCl_3), COSY and HMBC. Xanthone 1(115)	146
4.11	^1H NMR (400 MHz, CDCl_3) and ^{13}C NMR (100 MHz, CDCl_3) of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone and (116)	157
4.12	^1H NMR (400 MHz, CDCl_3), ^{13}C NMR (100 MHz, CDCl_3), COSY, and HMBC. Assignments of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methyl but-2-enyl)- xanthone (116)	158
4.13	^1H NMR (400 MHz, CD_3OD) and ^{13}C NMR (100 MHz, CD_3OD) of mangostanol and (68)	169



4.14	^1H NMR (400 MHz, CD_3OD), ^{13}C NMR (100 MHz, CD_3OD), COSY, and HMBC. Assignments of mangostanol (68)	170
4.15	^1H NMR (400 MHz, $\text{CDCl}_3/\text{CD}_3\text{OD}$), ^{13}C NMR (100 MHz, $\text{CDCl}_3/\text{CD}_3\text{OD}$), COSY, and HMBC. Assignments of mangoshaarin (117)	183
4.16	Cytotoxic Activity of plant extracts against HL-60 Cell Line (Promyelocytic Leukimia) and CEM-SS Cell Line (T-lymphoblastic Leukimia).	195
4.17	Antimicrobial activity of crude extracts and pure compound of <i>Annona muricata L.</i> and <i>Garcinia mangostana L.</i>	199
4.18	Larvicidal activity of crude extracts and pure compound against the larvae of <i>Aedes aegypti</i>	200



LIST OF FIGURES

Figures		Page
2.1	Core units for classification of annonaceous acetogenins	10
4.1	Compounds obtained from the stem bark of <i>Annona muricata L.</i>	71
4.2	EIMS spectrum of stigmasterol (111)	75
4.3	IR spectrum of stigmasterol (111)	76
4.4	^1H NMR spectrum of stigmasterol (111) (400 MHz, CDCl_3)	77
4.5	^{13}C NMR spectrum of stigmasterol (111) (100 MHz, CDCl_3)	78
4.6	CI-MS spectrum of solamin (22)	82
4.7	IR spectrum of solamin (22)	83
4.8	^1H NMR spectrum of solamin (22) (400 MHz, CDCl_3)	84
4.9	^1H - ^1H COSY spectrum of solamin (22) (400 MHz, CDCl_3)	85
4.10	^{13}C NMR spectrum of solamin (22) (100 MHz, CDCl_3)	86
4.11	^{13}C NMR spectrum of solamin (22) (100 MHz, CDCl_3) (expanded)	87
4.12	DEPT spectrum of solamin (22) (100 MHz, CDCl_3)	88
4.13	HSQC spectrum of solamin (22)	89
4.14	HMBC spectrum of solamin (22)	90
4.15	EIMS spectrum of sitosterol (112)	94
4.16	IR spectrum of sitosterol (112)	95
4.17	^1H NMR spectrum of sitosterol (112) (400 MHz, CDCl_3)	96



4.18	^{13}C NMR spectrum of sitosterol (112) (100 MHz, CDCl_3)	97
4.19	Compounds obtained from the bark of <i>Garcinia mangostana</i> L.	99
4.20	EIMS spectrum of friedelin (113)	103
4.21	IR spectrum of friedelin (113)	104
4.22	^1H NMR spectrum of friedelin (113) (400 MHz, CDCl_3)	105
4.23	^{13}C NMR spectrum of friedelin (113) (100 MHz, CDCl_3)	106
4.24	^{13}C NMR spectrum of friedelin(113) (100 MHz, CDCl_3) (expanded)	107
4.25	EIMS spectrum of α -mangostin (64)	111
4.26	IR spectrum of α -mangostin (64)	112
4.27	^1H NMR spectrum of α -mangostin (64) (400 MHz, CD_3OD)	113
4.28	^1H - ^1H COSY spectrum of α -mangostin (64) (400 MHz, CD_3OD)	114
4.29	^1H - ^1H COSY spectrum of α -mangostin (64) (400 MHz, CD_3OD) (expanded)	115
4.30	^{13}C NMR spectrum of α -mangostin (64) (100 MHz, CD_3OD)	116
4.31	HSQC spectrum of α -mangostin (64)	117
4.32	HMBC spectrum of α -mangostin (64)	118
4.33	EIMS spectrum of β -mangostin (61)	122
4.34	IR spectrum of β -mangostin (61)	123
4.35	^1H NMR spectrum of β -mangostin (61) (400 MHz, CDCl_3)	124
4.36	^1H - ^1H COSY spectrum of β -mangostin (61) (400 MHz, CDCl_3)	125
4.37	^{13}C NMR spectrum of β -mangostin (61) (100 MHz, CDCl_3)	126
4.38	DEPT spectrum of β -mangostin (61) (100 MHz, CDCl_3)	127



4.39	HSQC spectrum of β -mangostin (61)	128
4.40	HMBC spectrum of β -mangostin (61)	129
4.41	EIMS spectrum of garcinone D (114)	135
4.42	IR spectrum of garcinone D (114)	136
4.43	^1H NMR spectrum of garcinone D (114) (400 MHz, $\text{Me}_2\text{CO-d}_6$)	137
4.44	^1H NMR spectrum of garcinone D (114) (400 MHz, $\text{Me}_2\text{CO-d}_6$) (expanded)	138
4.45	^1H - ^1H COSY spectrum of garcinone D (114) (400 MHz, $\text{Me}_2\text{CO-d}_6$)	139
4.46	^{13}C NMR spectrum of garcinone D (114) (100 MHz, $\text{Me}_2\text{CO-d}_6$)	140
4.47	HSQC spectrum of garcinone D (114)	141
4.48	HMBC spectrum of garcinone D (114)	142
4.49	EIMS spectrum of xanthone 1 (115)	147
4.50	IR spectrum of xanthone 1 (115)	148
4.51	^1H NMR spectrum of xanthone 1(115) (400 MHz, CDCl_3)	149
4.52	^1H - ^1H COSY spectrum of xanthone 1(115) (400 MHz, CDCl_3)	150
4.53	^{13}C NMR spectrum of xanthone 1(115) (100 MHz, CDCl_3)	151
4.54	DEPT spectrum of xanthone 1(115) (100 MHz, CDCl_3)	152
4.55	HSQC spectrum of xanthone 1 (115)	153
4.56	HMBC spectrum of xanthone 1 (115)	154
4.57	EIMS spectrum of 1,6-dihydroxy-3,7-dimethoxy-2 -(3-methylbut-2-enyl)-xanthone (116)	159



4.58	IR spectrum of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone (116)	160
4.59	¹ H NMR spectrum of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone (116) (400 MHz, CDCl ₃)	161
4.60	¹ H- ¹ H COSY spectrum of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone (116) (400 MHz, CDCl ₃)	162
4.61	¹³ C NMR spectrum of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone (116) (100 MHz, CDCl ₃)	163
4.62	HSQC spectrum of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone (116)	164
4.63	HMBC spectrum of 1,6-dihydroxy-3,7-dimethoxy-2-(3-methylbut-2-enyl)-xanthone (116)	165
4.64	EIMS spectrum of mangostanol (68)	171
4.65	IR spectrum of mangostanol (68)	172
4.66	¹ H NMR spectrum of mangostanol (68) (400 MHz, CD ₃ OD)	173
4.67	¹ H- ¹ H COSY spectrum of mangostanol (68) (400 MHz, CD ₃ OD)	174
4.68	¹³ C NMR spectrum of mangostanol (68) (100 MHz, CD ₃ OD)	175
4.69	DEPT spectrum of mangostanol (68) (100 MHz, CD ₃ OD)	176
4.70	HSQC spectrum of mangostanol (68)	177
4.71	HSQC spectrum of mangostanol (68) (expanded)	178
4.72	HMBC spectrum of mangostanol (68)	179
4.73	HMBC correlations of mangosharin (117)	184
4.74	EIMS spectrum of mangosharin (117)	185
4.75	IR spectrum of mangosharin (117)	186
4.76	¹ H NMR spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD)	187



4.77	¹ H NMR spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD) (expanded)	188
4.78	¹ H- ¹ H COSY spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD)	189
4.79	¹ H- ¹ H COSY spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD) (expanded)	190
4.80	¹³ C NMR spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD)	191
4.81	HSQC spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD)	192
4.82	HMBC spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD)	193
4.83	HMBC spectrum of mangosharin (117) (400 MHz, CDCl ₃ /CD ₃ OD) (expanded)	194



LIST OF CHARTS

Charts		Page
4.1	Cytotoxic activity for the crude extract of <i>Annona muricata L.</i> towards the CEM-SS cell line (T-lymphoblastic leukemia).	196
4.2	Cytotoxic activity for the crude extracts of <i>Annona muricata L.</i> towards the HL-60 cell line (Promyelocytic Leukimia).	197
4.3	Cytotoxic activity for the crude extract of <i>Garcinia mangostana L.</i> towards the CEM-SS cell line (T-lymphoblastic Leukimia).	197
4.4	Cytotoxic activity for the pure compounds of <i>Garcinia mangostana L.</i> towards the CEM-SS cell line (T-lymphoblastic Leukimia).	198



LIST OF ABBREVIATIONS

α	alpha
β	beta
δ	chemical shift in ppm
γ	gamma
μg	micro gram
br s	broad singlet
br t	broad triplet
^{13}C	carbon-13
CHCl_3	chloroform
CDCl_3	deuterated chloroform
COSY	Correlated Spectroscopy
d	doublet
dd	doublet of doublet
DEPT	Distortionless Enhancement by Polarization Transfer
DMSO	dimethylsulfoxide
dt	doublet of triplet
EA	ethyl acetate
EIMS	Electron ionisation mass spectrometry
g	gram
GC	Gas Chromatography
GC-MS	Gas Chromatography- Mass spectrometry
^1H	proton



HETCOR	Heteronuclear Chemical Shift-correlation
HMBC	Heteronuclear Multiple Bond Connectivity by 2D Multiple Quantum
HPLC	High Performance Liquid Chromatography
Hz	Hertz
IC	Inhibition Concentration
IR	Infra Red
<i>J</i>	coupling constant in Hz
<i>L</i>	<i>LINN</i> .
l	litre
LC	Lethal Concentration
LD	Lethal Dose
m	multiplet
ml	mili litre
Me ₂ CO	acetone
MeOH	methanol
m.p.	melting point
MS	Mass Spectrum/Spectra/Spectrometer
NMR	Nuclear Magnetic Resonance
ppm	part per million
s	singlet
t	triplet
TLC	Thin Layer Chromatography



UV **Ultra Violet**

WHO **World Health Organization**

