



UNIVERSITI PUTRA MALAYSIA

**IMMUNOLOGICAL RESPONSE OF SHEEP TO *EPERYTHROZON*
OVIS INFECTION**

SHANKAR GANESH KANABATHY

FPV 2004 16

**IMMUNOLOGICAL RESPONSE OF SHEEP TO *EPERYTHROZON OVIS*
INFECTION**

By

SHANKAR GANESH KANABATHY

**Thesis submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirement for the Degree of Master of
Veterinary Science**

June 2004



DEDICATION

To my parents, family members and beloved wife for their kind support all these while and to the Lotus Feet of Lord Sri Balasubramaniyar, Taman Seri Timah, Balakong



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Degree Masters in Veterinary Science

IMMUNOLOGICAL RESPONSE OF SHEEP TO *EPERYTHROZOON OVIS* INFECTION

By

SHANKAR GANESH KANABATHY

June 2004

**Chairman : Associate Professor Che Teh Fatimah Nachiar
Iskandar, Ph.D.**

Faculty : Veterinary Medicine

The immunity of sheep to *Eperythrozoon ovis* (*E.ovis*) has been investigated through the peripheral blood smears stained with Giemsa. A naturally infected flock monitored for a year revealed the activity of peripheral blood monocytes to be involved in active phagocytosis of infected erythrocytes; a process called erythrophagocytosis. Although neutrophils, lymphocytes and thrombocytes were found to be activated in the initial stage of immune response, the monocytes seemed to predominate the phagocytosis at the later stage of infection during erythrophagocytosis. At all stages and degree of infections, no obvious anaemia, jaundice and emaciation were observed in these well fed sheep flocks. Anaemia was observed in flocks where malnourishment and stress conditions were present with a consistent high degree of parasitaemia. *E.ovis* infection trial in mice



exhibited more lymphocytic activities compared to the sheep , although lymphocytes, neutrophils and thrombocytes were involved in the early enhancement of inflammatory process against *E.ovis* as per in the sheep. These inflammatory processes were observed at day 20 post infection in mice. Similarly, only monocytes were found to be actively involved in erythrophagocytosis at the later stage of infection prior to the disappearance of the organisms from the peripheral circulation. Increased Kupffer cell activity showed liver was also involved in the removal of infected erythrocytes besides the blood peripheral macrophages.

In vitro phagocytosis assay using the Acridine Orange as the flurochrome revealed that peripheral monocytes ingested around eight cells of *E.ovis* per monocyte within 30 minutes upon contact. These cells were also killed within 30 minutes upon ingestion, characterised as red cells within the cytoplasm of monocytes. The Enzyme-linked Immunosorbent Assay was possible for optimization and was not suitable for further development as the Lang's method yields impure antigen from blood lysates. Latex test development was hindered due to the various host and immune serum factors that have resulted in non- specific agglutinations.

The persistence of infection in the flock throughout the one - year period of observation signified that sheep had been constantly infected with *E.ovis* and remained carriers for a very long period. The persistent parasitemia may suggest that the immunity to the parasite has been very complex probably due to highly



diversed antigenic variants, a characteristic exhibited by most rickettsiae in the Order of *Rickettsiales* or as a result of detrimental effects of the organism on the immune mechanism.

Sheep flocks naturally infected with *E.ovis* have remained permanent carriers. The findings from this research suggest that the sheep was unable to confer an effective or protective immune response against the pathogen. Peripheral blood macrophages are the most important first line of defense in removing the *E.ovis* from the peripheral blood.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains Veterinar

**TINDAKBALAS IMUN BEBIRI TERHADAP JANGKITAN
*EPERYTHROZOON OVIS***

Oleh

SHANKAR GANESH KANABATHY

Jun 2004

Pengerusi : Profesor Madya Che Teh Fatimah Nachiar Iskandar, Ph.D.

Fakulti : Perubatan Veterinar

Tindakbalas imun bebiri terhadap jangkitan *Eperythrozoon ovis* (*E.ovis*) telah dikaji melalui saringan darah yang diwarnakan dengan pewarna Giemsa. Kumpulan bebiri yang telah dijangkiti secara semulajadi telah disaring selama setahun. Ianya menunjukkan bahawa monosit darah memainkan peranan yang penting dalam membunuh sel darah merah yang telah dijangkiti *E.ovis*, proses yang dikenali sebagai 'erythrophagocytosis'. Walaupun neutrofil, limfosit dan trombosit dirangsang pada peringkat permulaan inflamasi; monosit merupakan sel yang terus menjalankan proses 'erythrophagocytosis' sehingga akhir proses ini. Pada semua peringkat inflamasi, tiada tanda-tanda kurang darah diperhatikan pada kumpulan bebiri ini. Kekurangan darah telah diperhatikan pada kumpulan bebiri yang tidak mempunyai sumber makanan yang mencukupi. Ujikaji jangkitan *E.*



ovis dalam tikus menunjukkan aktiviti limfosit yang lebih ketara berbanding bebiri walaupun neutrofil dan trombosit turut serta dalam proses inflamasi terhadap *E.ovis*. Proses inflamasi ini diperhatikan berlaku dalam jangkamasa 20 hari dari tempoh jangkitan tikus. Seperti dalam bebiri, monosit merupakan sel yang aktif dalam proses ‘erythrophagocytosis’ atau proses pemakanan sel darah merah. Pertambahan aktiviti sel Kupffer di dalam hati menunjukkan bahawa hati turut serta dalam proses pemusnahan dan penyingkiran sel darah merah yang dijangkiti *E.ovis*.

Ujikaji pemakanan *in vitro* *E .ovis* oleh monosit telah dijalankan dengan menggunakan ‘Acridine Orange’ sebagai fluorokrom. Eksperimen ini menunjukkan bahawa setiap monosit memakan hampir 8 sel *E .ovis* dalam masa 30 minit dari masa pertemuan. Sel *E. ovis* telah dibunuh sepenuhnya dalam masa 30 minit ini, diperhatikan sebagai sel yang telah menjadi merah sepenuhnya dalam sitoplasma monosit. Penghasilan sistem ‘Enzyme-linked Immunosobent Assay’ berjaya ke tahap penentuan faktor-faktor optima sahaja disebabkan kewujudan reaksi tak spesifik berpunca dari antigen yang tak tulin melalui kaedah Lang. Penghasilan ‘Latex Test’ juga tergendala disebabkan reaksi tak spesifik berpunca dari faktor haiwan dan sera bahan ujikaji .

Jangkitan *E. ovis* yang berterusan pada kumpulan bebiri selama setahun menunjukkan bahawa bebiri sentiasa dijangkiti oleh *E. ovis* dan status pembawa telah wujud untuk masa yang berpanjangan. Jangkitan terus menerus ini



menunjukkan bahawa tahap imun kepada parasit ini amat kompleks, mungkin disebabkan ciri-ciri organisma ini yang luas variannya; ciri-ciri yang biasa ditunjukkan oleh patogen – patogen dalam Order ‘*Rickettsiales*’; atau disebabkan oleh kesan buruk patogen ini terhadap sistem imun haiwan.

Kumpulan bebiri yang dijangkiti *E.ovis* secara semulajadi menjadi pembawa penyakit ini. Hasil kajian ini menunjukkan bahawa bebiri gagal mempertahankan dirinya dari segi tindakbalas imuniti terhadap patogen ini. Makrofaj darah merupakan pertahanan badan yang pertama dalam pemusnahan *E.ovis* dalam darah.

ACKNOWLEDGEMENT

I would like to express my heartiest gratitude to Assoc. Prof. Dr. Fatimah for giving me a chance to continue the research in this field. I am also grateful to the Ministry of Science through the IRPA project 01-02-04-0433 for funding the research and financially supporting me for my studies.

Sincere thanks are expressed to Dr. Ungku Chulan and Assoc. Prof. Dr. Abdul Rahman Omar for being patient to deal with the problems raised during the research period. Their kind guidance and encouragement are highly appreciated. Heartiest gratitude is forwarded to Dr. Nadzri Salim for his guidance.

Finally, I'm grateful to the members of faculty especially staff and post graduate students of Biologics Laboratory for assisting and guiding me during the course of research.





This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Veterinary Science. The members of the Supervisory Committee are as follow :

CHE TEH FATIMAH NACHAR ISKANDAR, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

ABDUL RAHMAN OMAR, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

NADZRI SALIM, M.V.S.

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

AINI IDERIS, Ph.D.

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date :



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

SHANKAR GANESH KANABATHY

Date :



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvii
LIST OF FIGURES	xviii
LIST OF PLATES	xix
LIST OF ABBREVIATIONS	xx
 CHAPTER	
1.0 INTRODUCTION	1
2.0 LITERATURE REVIEW	5
2.1 History and Epidemiology of <i>Eperythrozoon ovis</i>	5
2.2 Pathogenesis of <i>Eperythrozoon ovis</i>	6
2.3 Host Immune Response to Rickettsial Infections	10
2.3.1 Role of the peripheral leucocytes against the invading blood parasites	14
2.4 Immunity to <i>Eperythrozoon ovis</i> Infection	15
2.5 Diagnosis of <i>E. ovis</i>	17
2.6 Vaccination against Rickettsial Infections	23
2.7 Relationship of <i>Eperythrozoon</i> spp. to the Wall-less Prokaryotes, <i>Mycoplasma</i> spp.	25
2.7.1 Host Immune Response to <i>Mycoplasma</i> spp.	25



3.0.	HOST IMMUNE RESPONSE TO EPERYTHROZONOSIS	29
3.1	Introduction	29
3.2	Materials and Methods	30
3.2.1	Assessment of <i>in vitro</i> blood leucocyte microbial killing	30
3.2.1.1	<i>E.ovis</i> antigen cell count	30
3.2.1.2	Preopsonization of <i>E.ovis</i> antigen with pooled sheep serum	31
3.2.1.3	Assesment of leucocyte killing of <i>E.ovis</i> using Acridine Orange as a flurochrome	32
3.2.2	Peripheral blood screening of well managed and poorly managed sheep flocks	33
3.2.3	Peripheral blood response of naturally infected sheep flocks to <i>E.ovis</i>	34
3.2.4	Peripheral blood response of experimentally infected mice to <i>E.ovis</i>	34
3.3	Results	36
3.3.1	<i>E.ovis</i> antigen cell count	36
3.3.2	Assessment of <i>in vitro</i> blood leucocyte microbial killing	36
3.3.3	Peripheral blood screening of well managed and poorly managed sheep flocks	37
3.3.4	Blood cellular response of sheep naturally infected with <i>E.ovis</i>	39
3.3.5	Blood cellular response of mice experimentally infected with <i>E.ovis</i>	41
3.4	Discussion	45
4.0	LATEX TEST FOR RAPID DIAGNOSIS OF <i>E. OVIS</i>	47
4.1	Introduction	47
4.2	Materials and Methods	49
4.2.1	Immunolabelling latex particles and optimization of serum dilution	49



4.2.1.1	Latex test	50
4.3	Results	51
4.3.1	Immunolabelling latex particles and optimization of serum dilution	51
4.4	Discussion	53
5.0	ENZYME-LINKED IMMUNOSORBENT ASSAY FOR THE DETECTION OF ANTIBODIES TO <i>E. OVIS</i>	55
5.1	Introduction	55
5.2	Materials and Methods	58
5.2.1	Harvesting <i>E.ovis</i> from sheep erythrocytes	58
5.2.2	Embryonated egg culture of <i>E.ovis</i>	59
5.2.3	Propagation of <i>E.ovis</i> within mice	60
5.2.4	Protein microassay to determine the antigen protein concentration	61
5.2.5	Hyperimmunization of sheep and goat with <i>E.ovis</i>	62
5.2.6	Sucrose gradient centrifugation for antigen purification	62
5.2.7	Enzyme linked immunosorbent assay for the detection of <i>E.ovis</i> antibodies	63
5.2.7.1	Immobilisation of antigen on microtitre plate	63
5.2.7.2	Optimization of antigen, serum, conjugate and reaction time	64
5.2.7.3	Test proper	65
5.3	Results	66
5.3.1	Harvesting <i>E.ovis</i> from infected sheep erythrocytes	66
5.3.2	Embryonated egg culture of <i>E.ovis</i>	66
5.3.3	Propagation of <i>E.ovis</i> within mice	68
5.3.4	Determination of antigen protein concentration	69
5.3.5	Production of hyperimmune serum	70
5.3.6	Sucrose gradient centrifugation	71
5.3.7	Optimization for antigen , serum dilution, conjugate dilution and reaction time	71
5.3.8	Test proper	76



5.4	Discussion	77
6	GENERAL DISCUSSION AND CONCLUSION	80
	BIBLIOGRAPHY	86
	APPENDICES	93
	BIODATA OF THE AUTHOR	96



LIST OF TABLES

Table		Page
2.1	Diagnostic techniques of <i>E.ovis</i>	21
4.1	The agglutination of latex to the <i>E.ovis</i> free and infected erythrocytes	51
5.1	<i>E.ovis</i> in yolk, liver, spleen and kidney of embryonated hen egg at day 15 post incubation	67
5.2	Experimental infection , incubation period and parasitic level attained in mice	69
5.3	Checkerboard titration : Optimization for antigen concentration	72
5.4	Checkerboard titration : Optimization for serum dilution	73
5.5	Checkerboard titration : Optimization for conjugate dilution	74
5.6	Checkerboard titration : Optimization for reaction time	75
5.7	The non-specific OD reading in the blank and test wells.	76



LIST OF FIGURES

Figure		Page
3.1	The level of parasitemia in a sheep in well managed flock and poorly managed flock	38
5.1	The parasitic cycle in the experimentally infected mice.	68
5.2	The regression line of protein concentration of <i>E.ovis</i> harvested from blood of naturally infected sheep on absorbance at 595 nm	70
5.3	<i>E.ovis</i> antigen optimization curve for ELISA.	72
5.4	Serum optimization curve for ELISA.	73
5.5	Conjugate dilution optimization curve for ELISA	74
5.6	Reaction time optimization curve for ELISA	75



LIST OF PLATES

Plate		Page
3.1	Toxic effects of Acridine Orange on <i>E.ovis</i> and phagocytic cells <i>in vitro</i>	37
3.2	Peripheral blood monocyte actively phagocytosing <i>E.ovis</i> infected erythrocytes in sheep	39
3.3	The aggregation of lymphocyte, neutrophil and monocyte at site of inflammation in peripheral blood activated by platelet in sheep	40
3.4	The activity of monocytes , lymphocytes, neutrophils and platelets in the phagocytisation of <i>E.ovis</i> parasitised erythrocytes in mice.	41
3.5	Liver of a control mouse with a normal number and shape of Kupffer cells	42
3.6	Liver of an infected mouse showing increased number and size as well as rounding of the Kupffer cells nuclei	42
3.7	Spleen of a normal control mouse	43
3.8	Spleen of an <i>E.ovis</i> -infected mouse showed increased hemosiderosis	43
3.9	Kidney of a normal mouse showed spaces among glomerular tuft	44
3.10	Kidney of an <i>E.ovis</i> -infected mouse with the proliferative changes in the glomerulus with the compacted appearance of the glomerulus	44
4.1	Non specific agglutination of positive and negative serum labeled latex in blood negative for <i>E.ovis</i>	52



LIST OF ABBREVIATIONS

AO	Acridine Orange
BSA	Bovine Serum Albumin
CFT	Complement Fixation Test
DNA	Deoxyribonucleic acid
EDTA	Ethylene diamine tetra acetic acid
ELISA	Enzyme-linked immunosorbent assay
FCA	Freund's Complete Adjuvant
FIA	Freund's Incomplete Adjuvant
g	Gram
Hb	Hemoglobin
HBSS	Hank's Balanced Salt Solution
IFAT	Immunofluorescent Antibody Test
IFN- γ	Interferon-gamma
IHA	Indirect Hemagglutination Test
LAT	Latex Agglutination Test
mg	Milligram
ml	Millilitre
NK	Natural Killer
OD	Optical Density
PBS	Phosphate Buffered Saline
PVP	Polyvinylpyrrolidone
PBST	Phosphate Buffer Saline Tween
RBC	Red Blood Cell
SPF	Specific Pathogen Free
TMB	Tetramethylbenzidine
μ l	Microlitre



CHAPTER 1

INTRODUCTION

The economic losses due to *Eperythrozoon ovis* (*E.ovis*) infection in the sheep industry are reduced wool and reproduction, poor wool growth (Kreier and Ristic, 1963), increased risk to other diseases in chronically infected animals and mortality (Gulland *et al*, 1987). *Eperythrozoon ovis* has been isolated from sheep in many countries and usually produces mild clinical signs in experimentally inoculated animals. In some circumstances it is associated with a severe disease in young sheep known as ill-thrift. Ill-thrift is restricted to certain geographic situation and has been reported from Australia, New Zealand, France, Norway and South Africa, characterized by a failure of young sheep to thrive when sheep of all other ages appear to be in good health and weight gain (Stewart, 1981). The first report on eperythrozoonosis in Malaysia was by Fatimah *et al*, (1994) from a sheep concurrently suffering from copper toxicity. Mariah *et al*, (1997) reported that the morphology of *E. ovis* in sheep and goats as being coccoid and rod - like in sheep.

The organism is classified based on the species of animals being infected: *Eperythrozoon wenyoni* in cattle, *Eperythrozoon suis* in pigs and *Eperythrozoon ovis* in sheep and goat. The identification of the organism is based on the demonstration of antigen and the antibodies. Prior to 1970, the diagnosis of eperythrozoonosis was based on herd and individual animal histories of



icteroanemia and demonstration of eperythrozoon bodies in blood smears stained by Giemsa. In ovine and bovine eperythrozoonosis, complement fixation test (CFT), enzyme linked immunosorbent assay (ELISA) and passive hemagglutination tests have been developed to detect micro-organisms in the circulating red blood cells (Daddow , 1977; Finerty *et al* ,1969; Kawazu *et al* , 1990; Lang *et al* , 1987). Smith and Rahn (1975) developed an indirect hemagglutination test (IHA) for measuring antibodies to *E.suis*. Smith (1981) reported that IHA negative pigs could be carriers because parasitemia was observed in IHA negative pigs after splenectomy, indicating that the test may lack sensitivity for detecting chronically infected carriers.

The diagnosis of *E.ovis*, based mainly on peripheral blood smear can sometimes be difficult because this test may not be specific and is done on unclotted red blood cells which are frequently lysed in transit. A modified indirect immunofluorescent antibody test (IFAT) was used for easy and specific diagnosis of *E.ovis* from field samples (Ilemobade and Blotkamp, 1978). Antibody levels in the complement fixation test disappeared by 80 days on average and that the test is of limited value in detecting previous infection or exposure (Daddow, 1977). The coating antigen used in ELISA was a crude preparation from infected red blood cell lysates and contained host red blood cell antigens. The presence of host proteins in ELISA coating antigens may interfere with the test results (Hsu *et al*, 1992; Schuller *et al*, 1990). In general, ELISA positive results are significant, but negative results provide no information on the infection status of the animal.



Serological tests for the diagnosis of eperythrozoon have limitation due to marked variability in antibody response, as well as a failure to identify acutely infected sheep. Due to the lack of an efficient test to identify sheep and goats which are chronically or latently infected with *E.ovis*, the true incidence and economic impact of subclinical *E.ovis* infection is unknown.

Molecular techniques such as polymerase chain reaction (PCR), deoxyribonucleic acid (DNA) sequencing and Western blotting offer virtually unlimited opportunities to improve the ability to study and diagnose disease. As diagnostic tools, they have a better sensitivity and specificity compared to most immunological tests (Cox *et al*, 1991; Deacon and Lah, 1989; Peter, 1991). These techniques also offer a means of studying the genetic and pathologic basis of disease at the molecular level.

E.ovis has been detected in Malaysia back in 1980's and lately by Fatimah *et al*, (1994) from a sheep suffering from copper toxicity. Losses due to ovine eperythrozoonosis in Malaysia is unknown and it is therefore difficult to determine losses in an individual flock because of the lack of complete understanding of the nature of the disease and definitive diagnostic aids. However, a diagnostic tool such as the ELISA to detect the antibodies to *E.ovis* is sufficient to enable us to understand the prevalence rate of previous and current exposure. Besides, ELISA too allows the screening of a large number of samples in a day.



The report on erythrophagocytosis in peripheral blood, to date, is yet to be reported and there have been great discrepancies in reports on the role of humoral antibody in protecting sheep from reinfection. As parasitaemia fluctuates in the peripheral blood, a rapid field diagnosis is required for identification of infected animals . Therefore, the hypotheses and objectives of the study were as follow :

The hypotheses were :

- (1) A high antibody titre is correlated with a low level of parasitemia due to *E.ovis*.
- (2) Macrophage plays an important role in phagocytosis and elimination of *E.ovis* in peripheral blood.

The objectives of this study were:

- (1) To demonstrate the role of macrophages in phagocytosing *E.ovis* and elimination of the organisms from the peripheral blood.
- (2) To develop a Latex Test as an alternative for rapid diagnosis of *E.ovis* in the field
- (3) To determine the antibody level at different phases of parasitemia using ELISA.