



**UNIVERSITI PUTRA MALAYSIA**

**EPIDEMIOLOGY AND MORPHOLOGY OF LUNGWORM  
(*DICTYOCAULUS VIVIPARUS*), AND ITS ASSOCIATED LUNG  
PATHOLOGY IN CATTLE AND BUFFALOES IN PENINSULAR  
MALAYSIA**

**LAT LAT HTUN**

**FPV 2007 8**

**EPIDEMIOLOGY AND MORPHOLOGY OF LUNGWORM  
(*DICTYOCAULUS VIVIPARUS*), AND ITS ASSOCIATED LUNG  
PATHOLOGY IN CATTLE AND BUFFALOES IN PENINSULAR  
MALAYSIA**

**LAT LAT HTUN**

**DOCTOR OF PHILOSOPHY  
UNIVERSITI PUTRA MALAYSIA**

**2007**



**EPIDEMIOLOGY AND MORPHOLOGY OF LUNGWORM  
(*DICTYOCAULUS VIVIPARUS*), AND ITS ASSOCIATED LUNG  
PATHOLOGY IN CATTLE AND BUFFALOES IN PENINSULAR  
MALAYSIA**

**By**

**LAT LAT HTUN**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**July 2007**



## **DEDICATION**

**Dedicated with love and gratitude to my parents, U Thein Tun Oo and Daw Maing Saw Waing and gratitude to Dr. Saw Bawm and Moe Zaw Tun**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment  
of the requirement for the degree of Doctor of Philosophy

**EPIDEMIOLOGY AND MORPHOLOGY OF LUNGWORM  
(*DICTYOCAULUS VIVIPARUS*), AND ITS ASSOCIATED LUNG  
PATHOLOGY IN CATTLE AND BUFFALOES IN PENINSULAR  
MALAYSIA**

By

**LAT LAT HTUN**

**July 2007**

**Chairman: Latiffah Hassan, PhD**

**Faculty: Veterinary Medicine**

Bovine dictyocaulosis is an important parasitic disease of cattle and buffaloes and is caused by the lungworm, *Dictyocaulus viviparus*. The parasite is an important cause of lung infection especially in the temperate regions of the world. While the documentation on bovine lungworm is vast in the temperate, it is very sporadic and limited in the tropics. In Malaysia, a tropical country, the occurrence of lungworm infections in cattle and buffaloes has been anecdotal. The present study was carried out to detect the presence of lungworm infections in cattle and buffaloes, to determine the prevalence of lungworm infection in cattle and buffaloes, to identify the risk factors associated with bovine lungworm infection, to compare the morphology of egg, first stage larvae (L1) and adult stage of Malaysian bovine lungworm with those of *D. viviparus* from published reports and Sweden and to compare the histopathological lesions of lungs infected with Malaysian bovine lungworm and those of lungs infected with *D. viviparus*.



A retrospective examination of available records and data was carried out to investigate the presence of lungworm infections in Peninsular Malaysia. Two studies were carried out to address the objective. In the first study, an investigation on lungworm disease outbreak in a beef breeding farm was conducted. It was found that the yearly lungworm-infection mortality rate within the seven-year period was 0.31%. Among the cases, more than half (67%) were male and 33% were females. Seventy-five percent of lungworm infection deaths occurred in calves between the ages of six and 12 months, and 25% occurred in cattle aged 12 to 19 months. Most of the deaths occurred in November (19%) and May (17%). In the second study, data of condemnation of lungs and reasons of condemnation between 1998 and 2004 was collected at the Department of Veterinary Services Headquarters in Kuala Lumpur. Parasitic lung condemnation from all slaughtered animals was 0.11%. The prevalence of parasitic infection in the lungs was found much higher in buffaloes than in cattle ( $t = -3.906$ ,  $p = 0.002$ ).

A cross-sectional study was carried out in four large scale farms (Farms A, B, C and D) and three dairy smallholdings (Farm E) to detect and determine the prevalence of lungworm infection and to identify the risk factors. Blood and faecal sampling on each farm, except Farm E, was performed every two months for a period of seven months. Farm E was sampled only once. Questionnaires on individual animals, farm management and disease occurrence were developed and the data were collected at the time of blood and faecal sampling. Meteorological data was collected from the Climate Division, Malaysian Meteorological Service. The total blood and faecal samples collected from the farms were 602. Baermannisation was performed for parasitological diagnosis and enzyme-link-immunosorbent assay was conducted for

serodiagnosis. The prevalence of lungworm infection based on baermannisation was 4.7%. The highest prevalence was found in Farm E. Using binary logistic regression analysis, gender and the interaction between monthly temperature and monthly rainfall were identified as the statistically significant risk factors for bovine lungworm infection. The likelihood of lungworm infection was about four times greater when the monthly rainfall was >100 mm and the monthly temperature was >27°C to 29.1°C than when the monthly rainfall was <100 mm and when the temperature was <27°C ( $p = 0.002$ ). Female animals were about 2.9 times less likely to be infected than male animals ( $p = 0.01$ ).

Another cross-sectional study was carried out where 11 out of 25 abattoirs in Peninsular Malaysia were visited and slaughtered animals were examined. Animals slaughtered at Universiti Putra Malaysia (UPM) mosque during festivals were also examined. Among the total of 283 lungs from 260 cattle and 23 buffaloes sampled, lungworm was found in three Kedah-Kelantan (KK) cattle (1.1%). The morphological evaluation of egg, L1 and adult worm of the Malaysian bovine lungworm were conducted by comparing with those of *D. viviparus* from published reports and Sweden. Histopathological lesions of infected lungs were also examined. Based on the morphology of the lungworm and the histopathological changes of the affected lungs, the Malaysian bovine lungworm is believed to be most likely *D. viviparus*.

In conclusion, bovine lungworm infection in the Malaysian cattle and buffaloes can be detected and the prevalence is low. The disease occurrence was associated with the gender of the animals, and the climatic conditions. Based on the morphology of

the lungworm and the histopathological changes of the affected lungs, the Malaysian bovine lungworm is believed to be *D. viviparus*.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**EPIDEMIOLOGI DAN MORFOLOGI CACING PEPARU (*Dictyocaulus viviparus*), DAN PATOLOGI PEPARU YANG BERKAITAN PADA TERNAKAN LEMBU DAN KERBAU DI SEMENANJUNG MALAYSIA**

Oleh

**LAT LAT HTUN**

**July 2007**

**Pengerusi: Latiffah Hassan, PhD**

**Fakulti: Perubatan Veterinar**

Bovine dictyocaulosis ialah penyakit parasit yang penting pada ternakan lembu dan kerbau dan penyakit ini disebabkan oleh cacing peparu, *Dictyocaulus viviparus*. Parasit ini adalah penyebab utama jangkitan cacing peparu terutama kawasan dunia yang beriklim sederhana. Menurut laporan, cacing peparu lembu amat banyak terdapat di kawasan iklim sederhana tetapi hanya sekali sekala dan terhad di kawasan tropik. Di Malaysia negara beriklim tropika, kejadian jangkitan cacing peparu pada lembu dan kerbau merupakan anekdot. Kajian ini telah di jalankan untuk mengesan kehadiran jangkitan cacing peparu, menentukan kadar prevalens kejadian jangkitan cacing peparu pada lembu dan kerbau, memahami faktor risiko berkaitan dengan jangkitan cacing peparu, membuat bandingan morfometri dan morfologi peringkat larva dan peringkat dewasa cacing peparu lembu Malaysia dengan cacing *D. viviparus* dan membuat bandingan histopatologi pada lesion peparu yang di jangkiti cacing peparu lembu Malaysia dengan yang di jangkiti cacing peparu, *D. viviparus*.

Pemeriksaan retrospektif pada rekod dan data yang telah dijalankan untuk mengesan kejadian jangkitan cacing peparu lembu di Semenanjung Malaysia. Dua kajian telah dijalankan untuk mencapai objektif. Kajian awal telah dijalankan dipusat pembiakan lembu pedaging untuk menyiasat wabak jangkitan cacing peparu. Di dapati kadar kematian jangkitan cacing peparu tahunan bagi tujuh tahun di ladang tersebut adalah bersamaan 0.31%. Lebih banyak kes di kesan berlaku pada lembu jantan (67%) berbanding lembu betina (33%). Tujuh puluh lima peratus dari kematian akibat cacing peparu berlaku pada lembu yang berumur kurang daripada 12 bulan berbanding lembu berumur 12 hingga 19 bulan. Kebanyakan kematian ternakan lembu berlaku pada bulan November (19%) dan Mei (17%).

Pada kajian kedua, data penghapusan peparu serta penyebab penghapusan diantara tahun 1998 dan tahun 2004 telah di ambil daripada Ibu Pejabat Jabatan Perkhidmatan Haiwan di Kuala Lumpur. Peratusan penghapusan peparu berparasit yang diperolehi daripada semua sembelihan haiwan adalah bersamaan 0.11%. Kadar kelaziman jangkitan parasit peparu lebih tinggi pada kerbau berbanding pada lembu ( $t = -3.906$ ,  $p = 0.002$ ).

Satu kajian keratan rentas telah dijalankan pada empat buah ladang (ladang A, B, C, D) bersama tiga ladang tenusu kecil (ladang E) untuk mengenalpasti dan menentukan kadar prevalens jangkitan cacing peparu serta untuk mengenali faktor risiko. Sampel darah dan tinja najis telah diambil daripada setiap ladang setiap dua bulan selama tempoh lapan bulan. Hanya ladang E, sampel diambil sekali sahaja. Soal-selidik untuk setiap ekor haiwan, pengurusan ladang dan kejadian jangkitan dijalankan pada masa sampel darah dan najis diambil. Maklumat suhu dan hujan di perolehi daripada

Bahagian Kajicuaca, Jabatan Kajicuaca Malaysia. Sejumlah 602 sampel darah dan najis diambil daripada ladang tersebut. Teknik Baermann's digunakan untuk pemeriksaan parasitologi dan *enzyme-link-immunosorbent assay* untuk pemeriksaan serologi. Hasil pemeriksaan menggunakan teknik Baermann's mendapati, kadar prevalens jangkitan cacing peparu adalah bersamaan 4.7%. Ladang E menunjukkan kadar prevalens jangkitan yang paling tinggi. Dengan menggunakan analisis binari logistik regresi, jantina dan interaksi antara suhu bulanan dan taburan hujan bulanan telah menunjukkan keertian statistik sebagai faktor risiko jangkitan cacing peparu pada lembu. Kebarangkalian jangkitan cacing peparu bertambah empat kali ganda pada masa taburan hujan bulanan  $>100$  mm dan suhu bulanan  $>27^{\circ}\text{C} - 29.1^{\circ}\text{C}$  berbanding dengan pada masa taburan hujan bulanan adalah  $<100$  mm dan suhu bulanan  $<27^{\circ}\text{C}$ . Haiwan betina didapati kurang 2.9 kali ganda peluang jangkitan berbanding haiwan jantan.

Satu lagi kajian rentas telah dijalankan keatas 11 dari pada 25 rumah sembelih dan haiwan disembelih telah diperiksa. Haiwan yang disembelih di masjid Universiti Putra Malaysia semasa perayaan juga diperiksa. Cacing peparu telah dijumpai pada tiga lembu Kedah-Kelantan (1.1%), hasil pemeriksaan 283 peparu daripada 260 ekor lembu dan 23 ekor kerbau. Pemerhatian morfometri dan morfologi cacing peparu peringkat awal larva dan peringkat dewasa pada lembu Malaysia telah dibandingkan dengan cacing peparu *D. viviparus* di Sweden dan pada maklumat literasi. Pemeriksaan histopatologi juga dijalankan pada peparu yang dijangkiti. Berdasarkan pemerhatian morfometri dan morfologi cacing peparu dan hasil pemeriksaan histopatologi pada peparu yang dijangkiti adalah di percayai bahawa cacing peparu lembu Malaysia ialah *D. viviparus*.

Kesimpulannya, cacing peparu lembu dan jangkitannya boleh dikesan dan kadar prevalens adalah sangat rendah. Berlakunya jangkitan dihubungkan dengan kejantinaan haiwan dan keadaan cuaca. Berdasarkan morfologi cacing dan histopatologi peparu, cacing peparu Malaysia dipercayai adalah *D. viviparus*.

## ACKNOWLEDGEMENTS

I wish to express my profound appreciation and gratitude to my supervisory committee, Dr. Latiffah Hassan (Chairperson), Assoc. Prof. Dr. Rehana Abdullah Sani and Prof. Dato' Sheikh Omar Abdul Rahman for their invaluable guidance, advice, continuous supervision and encouragement throughout the course of study and in the preparation of this thesis.

I am especially grateful to the MTCP (Malaysian Technical Cooperation Programme) for providing the scholarship to perform this study. My special thanks are extended to His Excellency Agricultural Minister of Malaysia and to His Excellency Livestock and Fisheries Minister of Myanmar, for giving the opportunity to study in Malaysia.

I wish to thank the Department of Veterinary Services Malaysia, farm managers and their staff of the livestock centres, Pusat Ternakan Haiwan (PTH) Ayer Hitam, PTH Batu Arang, PTH Ulu Lepar, PTH Tersat, township veterinary officer of Banting and owners of small holder farms in Banting for granting me permission and helping to collect samples in the farms. I wish to thank the managers of abattoirs in Peninsular Malaysia for their assistance in sampling of lungworms and lungs.

Thanks are extended to all staff in the Public Health Unit, Department of Veterinary Services Malaysia and manager and all the staff of PTH Ulu Lepar, for their kind assistance and cooperation in collecting abattoir and outbreak records. I also wish to express my thanks to the staff from the Climate Division, Malaysian Meteorological



Service for their kind help to select the nearest weather stations to the sampled farms and to collect the meteorological data.

I am also grateful to Dr. Eysker and Dr. Ploeger from The Netherlands for their kind technical guidance during morphological evaluation of larva, Dr. Hoglund from Sweden for his kindly providing Swedish bovine lungworms to compare with Malaysian bovine lungworms and to Prof. P. Dorny from Belgium who first suggested studying the morphology of the Malaysian bovine lungworm. Special thanks to the late Dr. Nadzri Salim from the Faculty of Veterinary Medicine and Dr. Bahaman Abu Samah from the Faculty of Education for their kind and invaluable guidance in statistical analysis.

Sincere appreciation and gratefulness are also extended to Dr. Reuben Sunil Kumar Sharma for helping with the morphology study, Associate Professor Dr. Shaik Mohamed Amin Babjee, Dr. Sumita Sugnaseelan, Dr. Sabri Mohd Yusoff, Mrs. Maizaitul Akmal Moktar, Mr. Abd. Rashid Abdul Rahman, Mr. Mohd Nazari Abd. Hamid, Mr. Ghazali Md Yusoff and Mrs. Latiffah Mohd Hanan for their kind technical guidance and assistance during the laboratory work. Thanks also go to Mr. Krishnan Mariappan for his kind help in solving the problem of computer application.

I would also like especially to thank to Mr. Chamadrae Vengadasamy for his kind transportation to the farms during my sampling period. I wish to express my sincere appreciation to all my colleagues and friends especially, Dr. Khadak Singh Bisht, Dr. Shalina Bt. Mohmad Hal Hajja, Dr. Suriya Kumari A/P Ramiah, Dr. Intan Shameha

Abdul Razak, Dr. Noraniza Mohd Adzahan, Dr. Arul Jothi N and Ms. Siti Sarah Bt. Othman for their kindness and hospitality. My thanks are also in particular to all Myanmar friends and teachers for their warm friendship and kindness.

I would like to acknowledge to U Maung Maung Nyunt, Director General of Livestock Breeding and Veterinary Department, Dr. Myint Thein, Rector, and Pro-Rectors Dr. Khin Ma Ma and Dr. Tin Tin Myaing, University of Veterinary Science, Myanmar, for allowing me to further my study.

I am especially grateful to my retired Rectors Dr Min Soe and Dr. Htun Sein, Pro-Rector Dr. Ni Ni Maw, Professors Dr. Thein Tun, Dr. Sein Lwin and Dr. Yi Yi for their encouragement during my study. Special thanks also go to all my colleagues in the Department of Pharmacology and Parasitology, University of Veterinary Science, Myanmar, for conducting all my duties during my absence.

Deepest thanks are due to my late mother, my father and brother Moe Zaw Tun for their moral support and love. My heartfelt appreciation goes to Dr. Saw Bawm for his invaluable moral support and understanding throughout this study.



I certify that an Examination Committee has met on 6<sup>th</sup> July 2007 to conduct the final examination of Lat Lat Htun on her Doctor of Philosophy thesis entitled “Epidemiology and Morphology of Lungworm (*Dictyocaulus viviparus*), and its Associated Lung Pathology in Cattle and Buffaloes in Peninsular Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Abd. Aziz Saharee, PhD**  
Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

**Saleha Abd. Aziz, PhD**  
Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Internal Examiner)

**Shaik Amin Babjee, PhD**  
Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Internal Examiner)

**Pierre Dorny, PhD**  
Professor  
Faculty of Veterinary Medicine  
Gent University, Belgium  
(External Examiner)

---

**HASANAH MOHD. GHAZALI, PhD**  
Professor / Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 3 AUGUST 2007



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

**Latiffah Hassan, PhD**

Lecturer  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Chairman)

**Rehana Abdullah Sani, PhD**

Associate Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

**Dato' Sheikh Omar Abd. Rahman, PhD**

Professor  
Faculty of Veterinary Medicine  
Universiti Putra Malaysia  
(Member)

---

**AINI IDERIS, PhD**

Professor and Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 9 AUGUST 2007



## **DECLARATION**

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

---

**LAT LAT HTUN**

Date: 23 JULY 2007

## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	ii
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	vii
<b>ACKNOWLEDGEMENTS</b>	xi
<b>APPROVAL</b>	xiv
<b>DECLARATION</b>	xvi
<b>LIST OF TABLES</b>	xxi
<b>LIST OF FIGURES</b>	xxiii
<b>LIST OF ABBREVIATIONS</b>	xxvi
<b>CHAPTER</b>	
1 <b>INTRODUCTION</b>	1
2 <b>LITERATURE REVIEW</b>	7
2.1    The Organism and Disease	7
2.2    Morphology	7
2.3    Life Cycle	9
2.4    Pre-Parasitic Larval Stages	10
2.5    Epidemiology	14
2.5.1    Host Factor	14
2.5.2    Environmental Factor	18
2.6    Economic Importance of Bovine Dictyocaulosis	18
2.7    Diagnosis of Bovine Dictyocaulosis	20
2.7.1    Faecal Detection	20
2.7.2    Serology	22
2.7.3    Pathogenesis and Histopathological Lesions	26
2.8    Control of Bovine Dictyocaulosis	31
2.8.1    Treatment	31
2.8.2    Immunity and Vaccination	33
2.8.3    Grazing Management	37
3 <b>MATERIALS AND METHODS</b>	41
3.1    Target and Study Population	41
3.2    Study Design	41
3.2.1    Sample Size	42
3.2.2    Farm A	43
3.2.3    Farm B	45
3.2.4    Farm C	45
3.2.5    Farm D	46
3.2.6    Farm E	47
3.3    Collection of Sample	48
3.3.1    Faecal Sample	48
3.3.2    Baermann Technique	48
3.3.3    Blood Sample	50
3.3.4    Enzyme-Link Immunosorbent Assay	50
3.3.5    ELISA Test Procedures	51



	3.3.6	Calculation of Percent Positivity Values (PP)	53
	3.3.7	Interpretation of The Percent Positivity	54
4		<b>OUTBREAKS OF LUNGWORM DISEASE IN A BEEF FARM IN MALAYSIA BETWEEN 1994 AND 2000</b>	55
	4.1	Introduction	55
	4.2	Materials and Methods	56
	4.2.1	Farm Management and History	56
	4.2.2	Collection of Data	57
	4.2.3	Analysis of Data	58
	4.3	Results	58
	4.4	Discussion and Conclusion	61
5		<b>CONDEMNATION OF LUNGS IN ABATTOIRS IN PENINSULAR MALAYSIA DUE TO PARASITIC INFECTION FROM 1998 TO 2004</b>	65
	5.1	Introduction	65
	5.2	Materials and Methods	66
	5.2.1	Source of Data	66
	5.2.2	Data Management and Analysis	66
	5.3	Results	66
	5.3.1	Description of Data from Records	66
	5.3.2	Condemnation of Lungs	71
	5.3.3	Condemnation of Lungs due to Parasitic Infection	71
	5.4	Discussion and Conclusion	74
6		<b>PREVALENCE OF LUNGWORM INFECTION IN CATTLE AND BUFFALOES IN PENINSULAR MALAYSIA</b>	78
	6.1	Introduction	78
	6.2	Materials and Methods	80
	6.2.1	Determination of Prevalence Using the Parasitological Method and ELISA	80
	6.2.2	Determination of Prevalence from Examination of Slaughtered Animals	80
	6.2.3	Data Analysis	81
	6.3	Results	81
	6.3.1	Parasitological Prevalence	81
	6.3.2	Serological Prevalence	82
	6.3.3	Prevalence of Adult Lungworm at the Abattoirs	86
	6.4	Discussion and Conclusion	87
7		<b>IDENTIFICATION OF RISK FACTORS ASSOCIATED WITH BOVINE LUNGWORM INFECTION</b>	92
	7.1	Introduction	92

7.2	Materials and Methods	92
7.2.1	Questionnaire	93
7.2.2	Meteorological Data	93
7.2.3	Data Analysis	94
7.3	Results	94
7.3.1	Management of the Farms	94
7.3.2	Prevalence (%) and Meteorological Data from the Weather Stations Nearest to the Sampled Farms	97
7.3.3	Prevalence of Lungworm Infection	99
7.3.4	Univariate Analysis of Hypothesised Risk Factors of Age, Breed, Gender, Monthly Temperature, Monthly Rainfall and Grazing Management and Lungworm Infection	100
7.3.5	Logistic Regression Analysis	100
7.4	Discussion	102
7.4.1	Age	102
7.4.2	Gender	102
7.4.3	Monthly Rainfall and Monthly Temperature	103
7.4.4	Grazing Management	105
7.4.5	Breed	106
7.5	Conclusion	106
8	<b>MORPHOLOGICAL EVALUATION OF THE MALAYSIAN BOVINE LUNGWORM AND THE HISTOPATHOLOGY OF INFECTED LUNG</b>	107
8.1	Introduction	107
8.2	Materials and Methods	108
8.2.1	Sampling from the Abattoir	108
8.2.2	Morphology and Morphometry	109
8.2.3	Comparison of the Morphology of Egg, L1 and Adult of Malaysian Bovine Lungworm with <i>D. viviparus</i>	110
8.2.4	Histopathological Evaluations	110
8.2.5	Statistical Analysis	111
8.3	Results	111
8.3.1	Descriptive Morphology and Morphometry of Adult, L1 and Egg of Malaysian Bovine Lungworms	112
8.3.2	Comparative Morphometry of Adult Male and Female Worms, Egg and L1 between Malaysian Bovine Lungworm and <i>D. viviparus</i> from Published Reports and Sweden	113
8.3.3	Taxonomic Summary	123
8.3.4	Macroscopic and Microscopic Findings of Infected Lungs	124
8.4	Discussion and Conclusion	127
8.4.1	Morphology and Morphometry of Malaysian Bovine Lungworm	127

8.4.2	Pathology of Bovine Lungworm Infection	129
9	<b>SUMMARY, CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE RESEARCH</b>	133
	<b>REFERENCES</b>	141
	<b>APPENDICES</b>	151
	A. Forms for Data Collection from Abattoir Managers	151
	B. Questionnaires on Cross-sectional Study of Bovine Dictyocaulosis	155
	C. Morphometric Definitions	166
	D. Procedure for Staining	167
	<b>BIODATA OF STUDENT</b>	170
	<b>LIST OF PUBLICATIONS</b>	171



## LIST OF TABLES

Table		Page
2.1	Efficacy of current anthelmintics against <i>D. viviparus</i>	34
3.1	Population of cattle and management of five farms during the period of sample collection	44
4.1	Number of deaths in cattle due to lungworm disease between 1994 and 2000 in a beef farm in Peninsular Malaysia	59
5.1	Total number, mean and standard deviation of cattle and buffaloes slaughtered in each of the 25 abattoirs in Peninsular Malaysia from 1998 to 2004	68
5.2	Total number, mean and standard deviation of cattle and buffaloes slaughtered per year in 25 abattoirs in Peninsular Malaysia from 1998 to 2004	69
5.3	Number and rate (%) of lungs condemned due to specific lung lesions from 25 abattoirs in Peninsular Malaysia from 1998 to 2004	70
6.1	Prevalence (%) of bovine lungworm parasitologically positive cases in four large scale farms and one dairy smallholding	83
6.2	Seroprevalence (%) of lungworm in all diagnosed sera from four large scale farms and one dairy smallholding based on three age groups	85
6.3	Cross tabulation between parasitological and serological (weak positive) results	86
6.4	Prevalence (%) of cattle infected with adult lungworm examined at 11 abattoirs and UPM mosque in Peninsular Malaysia	87
7.1	Management of five farms during the period of sample collection	95

7.2	Prevalence (%) of bovine lungworm infection and meteorological data from the nearest weather stations to the sampled farms during the sampling period	98
7.3	Distribution and association of risk factors for the prevalence of bovine lungworm infection in five farms based on parasitological diagnosis	99
7.4	Binary logistic regression of risk factors for bovine lungworm infection based on parasitological diagnosis in five farms	101
8.1	Number of worms found in three Kedah-Kelantan cattle at two abattoirs	111
8.2	Comparative morphometry of male worms between the Malaysian bovine lungworm and <i>D. viviparus</i> from the published reports. All measurements are in mm (unless stated otherwise).	115
8.3	Comparative morphometry of female worms between the Malaysian lungworm and <i>D. viviparus</i> from the published reports and Sweden. All measurements are in mm (unless stated otherwise).	116
A.1	Form for collection of records on condemnation of lungs from abattoirs in Peninsular Malaysia (1994-2004)	148
A.3b	Borang maklumat haiwan ‘Projek cacing peparu didalam lembu dan kerbau	151



## LIST OF FIGURES

<b>Figure</b>		<b>Page</b>
3.1	Friesian-Sahiwal Cross Calf in Farm A	45
3.2	Nelore Breed Cattle in Farm C	46
3.3	Kedah-Kelantan Breed Cattle in Farm D	47
3.4	Table Constructed for the Baermann Technique	49
3.5	ELISA Kit, Ceditest <sup>®</sup> <i>Lungworm</i> (Cedi-Diagnostics B.V., The Netherlands)	51
4.1	The Number of Deaths due to Lungworm Infection between 1994 and 2000	59
4.2	The Number of Deaths due to Lungworm Infection in Different Age Groups between 1994 and 2000	60
4.3	The Number of Cumulative Monthly Deaths due to Lungworm Infection between 1994 and 2000	61
5.1	Species-specific Parasitic Lung Condemnation in 25 Abattoirs from 1998 to 2004	72
5.2	Species-specific Proportional Lung Condemnation due to Parasitic Infection in 25 Abattoirs from 1998 to 2004	73
5.3	Parasitic Lung Condemnation Rate from 1998 to 2004 Based on the Location of Abattoirs	73
5.4	Cumulative Monthly Distribution of Lung Condemnation due to Parasitic Infection in 25 Abattoirs from 1998 to 2004	74
6.1	Pictographs of the First Stage Larvae of Malaysian Bovine Lungworm from Sampled Farms	82