



UNIVERSITI PUTRA MALAYSIA

**DEVELOPMENT OF AN AVIRULENT *PASTEURELLA MULTOCIDA* B:2
BY DISRUPTION OF THE ABA392 DNA FRAGMENT**

DZARIFAH MOHAMED ZULPERI

FPV 2008 2

**DEVELOPMENT OF AN AVIRULENT *PASTEURELLA MULTOCIDA* B:2 BY
DISRUPTION OF THE ABA392 DNA FRAGMENT**

DZARIFAH MOHAMED ZULPERI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfillment of the Requirements for the
Degree of Master of Science**

August 2008



**DEVELOPMENT OF AN AVIRULENT *PASTEURELLA*
MULTOCIDA B:2 BY DISRUPTION OF THE ABA392
DNA FRAGMENT**

DZARIFAH MOHAMED ZULPERI

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2008



Specially dedicated to:

Abah and Mak,

ENCIK MOHAMED ZULPERI ZAKARIA
PUAN SAMIRAH ISMAIL

Adik,

ZARIRAH MOHAMED ZULPERI

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science

DEVELOPMENT OF AN AVIRULENT *PASTEURELLA MULTOCIDA* B:2 BY DISRUPTION OF THE ABA392 DNA FRAGMENT

By

DZARIFAH BINTI MOHAMED ZULPERI

Chairman: Siti Khairani Bejo, PhD

Faculty: Veterinary Medicine

Haemorrhagic septicaemia (HS) is an acute disease infecting cattle and buffaloes caused by *Pasteurella multocida* serotype B:2 which leads to great economic losses in countries with expanded animal industries such as Malaysia. This study was conducted to construct a mutant derived from the 921bp ABA392 virulence gene in order to make avirulent *P. multocida* serotype B:2, thus making this *P. multocida* B:2 mutant a potential candidate for a live-attenuated vaccine against HS.

The detection of a fragment which is related to *P. multocida* B:2 pathogenicity, the 921bp ABA392 virulence gene was carried out using Polymerase Chain Reaction (PCR) assay. Two *P. multocida* isolates, namely PMTB and 3030, isolated from a HS outbreak were used for PCR amplification. The particular 921bp DNA fragment was found to be present in *P. multocida* B:2 genome as 803bp in size. Sequencing of the recombinant plasmids confirmed that the 803bp gene was 98% homologous to the reference sequence, the virulence 921bp ABA392 DNA fragment.



Southern hybridization analysis revealed the 804bp of ABA392 gene was located at approximately 6kb position in the *P. multocida* B:2 genome. PCR performed towards the approximately 6kb DNA fragment produced an 803bp band which was confirmed to be the ABA392 virulence gene. The amplified PCR product was cloned and sequenced. Nucleotide sequences obtained were 98% identical to the reference strain, the 921bp ABA392 virulence gene. Pathogenicity test on the recombinant plasmids proved that the 804bp gene inserted within these plasmids still possessed the virulence properties of the 921bp ABA392 gene which may lead to HS disease in cattle and buffaloes.

In an attempt to produce *P. multocida* B:2 mutants through allelic exchange, the 804bp of ABA392 gene which was disrupted with kanamycin cassette was cloned inside suicide plasmid pAKA19 through shotgun ligation technique. The desired 7kb product was transformed into several different *E. coli* (TOP 10, JM109, AS111Y λ and DH5 α) hosts for preservation. Verification of the 7kb ligation product with digestion by *Pst*I, *Hind*III and *Xho*I restriction enzymes revealed the exact sizes of kanamycin cassette (1.2kb), pAKA19 (5.0kb) and the disrupted 804bp of ABA392 virulence gene (804bp). An antibiotic sensitivity test on *P. multocida* B:2 and the respective *E. coli* strains were performed in order to select the donor and recipient strains for conjugation process. This test revealed that *E. coli* DH5 α was suitable for the donor strain since it showed low resistance to streptomycin and *P. multocida* B:2 as the recipient strain for its high resistance towards streptomycin. Conjugation between donor and recipient strains was then achieved by plate-mating method. About 20 single colonies of

positive transconjugants were picked and subcultured on BHI blood agar containing kanamycin for 5 days to encourage loss of pAKA19 plasmid and to enhance the allelic exchange between the *ABA392::Km^R* insert with the native *ABA392* gene on the recipient chromosome. No plasmid was observed which indicated the loss of suicide plasmid. Direct colony PCR was performed to detect the changes in the *ABA392* gene of the parent strain, where a 2kb band was observed signifying that allelic exchange has taken place and the organism is now a *P. multocida* B:2 mutants.

Parent strains of *P. multocida* B:2 were highly virulent and killed mice within 24 hours. Mice inoculated with *P. multocida* B:2 mutant survived. Direct smear from mice's heart blood inoculated with the mutant showed the existence of bipolar organism which indicated the presence of *P. multocida* B:2. This result firmly suggests that the mutant, named as PMTBK was greatly attenuated and is thus a potential candidate organism for a live attenuated vaccine against HS.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGHASILAN *PASTEURELLA MULTOCIDA* B:2 AVIRULEN MELALUI
TINDAKAN GANGGUAN TERHADAP FRAGMENTEN DNA ABA392**

Oleh

DZARIFAH BINTI MOHAMED ZULPERI

Pengerusi: Siti Khairani Bejo, PhD

Fakulti: Perubatan Veterinar

Hawar berdarah (HS) merupakan penyakit akut yang menyerang lembu dan kerbau berpunca daripada bakteria *Pasteurella multocida* serotaip B:2 yang menyebabkan kemerosotan ekonomi bagi negara yang mempunyai industri penternakan berkembang maju seperti Malaysia. Kajian ini telah dijalankan untuk menghasilkan mutan avirulen daripada bakteria *P. multocida* B:2 terbitan daripada gen virulen ABA392 bersaiz 921bp, seterusnya menjadikan mutan ini calon yang berpotensi bagi penghasilan vaksin hidup teratenuat terhadap hawar berdarah.

Pengesanan fragmen DNA yang berkait rapat dengan kepatogenesis *P. multocida* B:2 iaitu gen virulen ABA392 bersaiz 921bp telah dijalankan melalui assai amplifikasi Tindakbalas Berantai Polymerase (PCR). Dua isolat *P. multocida* iaitu PMTB dan 3030 yang telah dipencilkan dari kawasan wabak penyakit hawar berdarah digunakan dalam amplifikasi PCR. Fragmen DNA bersaiz 921bp telah disahkan hadir dalam genom *P. multocida* B:2 dengan saiz 803bp. Penjujukan terhadap plasmid rekombinan membuktikan bahawa gen



803bp adalah 98% homologous dengan jujukan rujukan iaitu fragmen DNA ABA392 yang virulen.

Analisis penghibridan Southern menunjukkan gen ABA392 bersaiz 804bp terletak pada anggaran posisi 6kb dalam genom *P. multocida* B:2. Amplifikasi PCR ke atas fragmen DNA tersebut menghasilkan produk bersaiz 803bp lalu mengesahkan kehadiran gen virulen ABA392. Produk PCR ini kemudian diklon dan diujuk di mana jujukan-jujukan nukleotida yang diperoleh adalah 98% seiras dengan jujukan rujukan iaitu gen virulen ABA392 bersaiz 921bp. Ujian kepatogenan ke atas plasmid rekombinan membuktikan gen 804bp yang telah disisipkan ke dalam plasmid tersebut masih mengekalkan ciri-ciri virulen dalam gen ABA392 bersaiz 921bp yang menyebabkan penyakit hawar berdarah pada lembu dan kerbau.

Percubaan untuk menghasilkan mutan *P. multocida* B:2 melalui penukaran allele dijalankan melalui penyisipan kaset kanamycin ke dalam gen ABA392 bersaiz 804bp dan diklon ke dalam plasmid pengorban pAKA19 menerusi teknik ligasi 'shotgun'. Produk bersaiz 7kb yang diperoleh dipindahkan kepada beberapa perumah *E. coli* yang berbeza (TOP 10, JM109, AS11Yλ dan DH5α) untuk tujuan penyimpanan. Verifikasi ke atas produk ligasi bersaiz 7kb melalui tindakan pembatasan oleh enzim-enzim pembatas *Pst*I, *Hind*III dan *Xho*I mengesahkan kehadiran kaset kanamycin (1.2kb), pAKA19 (5.0kb) dan gen virulen gangguan ABA392 (804bp). Ujian sensitiviti terhadap antibiotik telah dijalankan ke atas *P. multocida* B: 2 dan strain-strain *E. coli* dalam usaha



memilih penderma dan penerima yang sesuai bagi proses konjugatan. *E. coli* DH5 α telah didapati sesuai sebagai penderma kerana mempunyai kerintangan rendah terhadap streptomycin manakala *P. multocida* B:2 sebagai penerima kerana kerintangan tinggi terhadap streptomycin. Proses konjugatan antara penderma dan penerima dijalankan melalui kaedah plat pengawanan. Kira-kira 20 koloni tunggal transkonjugan positif telah dipilih untuk disubkultur selama 5 hari berturut-turut pada agar darah BHI bagi menggalakkan pelenyapan plasmid pAKA19, di samping menggalakkan penukaran allele di antara gen sisipan *ABA392::Km^R* dan gen induk *ABA392* pada kromosom penerima. Ketidakhadiran plasmid pAKA19 melalui elektroforesis agarose gel membuktikan plasmid tersebut telah berjaya dilenyapkan daripada gen gangguan 804bp. Aplikasi PCR secara langsung bagi mengesan perubahan dalam gen induk strain *ABA392* menghasilkan jalur bersaiz 2kb menandakan berlakunya perubahan allele dan organisma tersebut kini adalah mutan *P. multocida* B:2.

Strain induk *P. multocida* B:2 didapati amat virulen kerana strain ini membawa maut kepada tikus dalam tempoh 24 jam. Walau bagaimanapun, tikus yang telah diinokulat dengan mutan *P. multocida* B:2 didapati hidup. Saput terus daripada jantung tikus yang diinokulat dengan mutan menunjukkan kewujudan organisma dwikutub yang menandakan kehadiran *P. multocida* B:2. Keputusan ini dengan kuat mencadangkan bahawa mutan iaitu PMTBK adalah avirulen dan merupakan calon yang berpotensi untuk penghasilan vaksin teratenuat hidup terhadap hawar berdarah.

ACKNOWLEDGEMENTS

Bismillahirrahmanirrahim

I would like to express sincere gratitude to everybody who helped me to get through this three and halves years. This feeling of grateful and blessed goes to my supervisor, Dr. Siti Khairani Bejo for her enthusiasm towards my work and for giving fruitful suggestions and ideas. Many thanks go to my co-supervisors, Dr. Zunita Zakaria for the optimistic views on my work when things get tougher and Prof. Dr. Raha Abdul Rahim for useful discussion on matters involving Southern hybridization analysis. Not to forget Prof. Dr. Abdul Rani Bahaman, Prof. Madya Dr. Abdul Rahman Omar and Dr. Salmah Ismail for their invaluable advice on my research.

Special thanks go to the staff of Bacteriology Laboratory, Molecular Biology Laboratory, Biologic Laboratory and Histopathology Laboratory for putting up with me for all these years. My heartiest appreciation also goes to all my dearest friends, Ina Salwany, Murni Marlina, Ernie Zuraida, Dr. Ani, Idah and Shafarin for their unconditional friendship and love.

To my family (Abah, Mak and Adik Zary) who I owed the most for their patience, understanding and support throughout the long and winding road of my master study.

This work was sponsored by MOSTI, whom I would like to express my gratitude for the financial support.



I certify that an Examination Committee has met on 11 August 2008 to conduct the final examination of Dzarifah binti Mohamed Zulperi on her Master of Science thesis entitled “Development of Avirulent *Pasteurella multocida* B:2 by Disruption of the ABA392 DNA Fragment” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the degree of Master of Science.

Members of the Examination Committee were as follows:

Abdul Aziz Saharee, PhD

Professor
Centre of External Education
Universiti Putra Malaysia
(Chairman)

Abdul Rani Bahaman, PhD

Professor
Department of Pathology and Veterinary Microbiology
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Siti Suri Arshad, PhD

Associate Professor
Department of Pathology and Veterinary Microbiology
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Ian Robertson, PhD

Associate Professor
Department of Veterinary Clinical Sciences
Murdoch University
Australia
(External Examiner)

Hasanah Mohd. Ghazali, PhD
Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 26 September 2008



DECLARATION

I hereby declare that the thesis is based on my original work for quotation and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any degree at UPM or other institutions.

DZARIFAH MOHAMED ZULPERI

Date: 26 September 2008



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENT	ix
APPROVAL	x
DECLARATION	xii
LIST OF FIGURES	xiii
LIST OF TABLES	xv
LIST OF APPENDIXES	xvi
LIST OF ABBREVIATIONS	xvii
TABLE OF CONTENTS	xx
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	5
2.1 Haemorrhagic septicaemia	5
2.2 Epidemiology of haemorrhagic septicaemia	6
2.3 Pathogenesis of haemorrhagic septicaemia	8
2.4 Treatment and control of haemorrhagic septicaemia	10
2.4.1 Treatment of haemorrhagic septicaemia	10
2.4.2 Vaccination for prevention and controlling Of haemorrhagic septicaemia	10
2.5 <i>Pasteurella multocida</i>	14
2.6 Pathogenicity of <i>Pasteurella multocida</i>	18
2.7 Live-attenuated vaccine against haemorrhagic septicaemia	21
2.7.1 The ABA392 gene	21
2.7.2 Suicide plasmid	23
2.7.3 Conjugation	25
2.7.4 Allelic exchange	27
3 DETECTION OF ABA392 VIRULENCE GENE AND ITS LOCATION IN <i>PASTEURELLA MULTOCIDA</i> B:2 GENOME	30
3.1 Introduction	30
3.2 Materials and methods	34
3.2.1 Bacterial strains	34
3.2.2 DNA extraction	34
3.2.3 Qualification and quantification of DNA	35
3.2.4 Amplification of ABA392 virulence gene from <i>Pasteurella</i> <i>multocida</i> B:2 genome	36
3.2.5 Purification of PCR product	38



3.2.6	Cloning of ABA392 gene into <i>E. coli</i> host	38
3.2.7	Analysis of positive clones	40
3.2.8	Development of a non-radioactive labeled probe	43
3.2.9	Southern hybridization of ABA392 virulence gene in <i>Pasteurella multocida</i> B:2 genome	44
3.2.10	Verification of ABA392 virulence gene location in <i>Pasteurella multocida</i> B:2 genome	47
3.2.11	Virulence test of pMTB and p3030 recombinant plasmids of <i>Pasteurella multocida</i> B:2 in mice	50
3.3	Results	52
3.3.1	Amplification of ABA392 gene from <i>Pasteurella multocida</i> B:2 genome	52
3.3.2	Cloning of ABA392 gene into <i>E. coli</i> host	52
3.3.3	Analysis of plasmid	52
3.3.4	Southern hybridization of ABA392 virulence gene in <i>Pasteurella multocida</i> B:2 genome	54
3.3.5	Verification of ABA392 virulence gene location in <i>Pasteurella multocida</i> B:2 genome	55
3.3.6	Virulence test of pMTB and p3030 recombinant plasmids of <i>Pasteurella multocida</i> B:2 in mice	56
3.4	Discussion	72
4	DEVELOPMENT OF AVIRULENT <i>PASTEURELLA MULTOCIDA</i> B:2 MUTANT	84
4.1	Introduction	84
4.2	Materials and methods	86
4.2.1	Disruption of ABA392 virulence gene with kanamycin cassette in suicide plasmid	86
4.2.2	Construction of avirulent <i>Pasteurella multocida</i> B:2 via allelic exchange	92
4.3	Results	97
4.3.1	Disruption of ABA392 virulence gene with kanamycin cassette in suicide plasmid	97
4.3.2	Construction of avirulent <i>Pasteurella multocida</i> B:2 via allelic exchange	99
4.4	Discussion	113
5	PATHOGENICITY AND STABILITY TESTS OF <i>PASTEURELLA MULTOCIDA</i> B:2 MUTANT	121
5.1	Introduction	121
5.2	Materials and methods	123
5.2.1	<i>In vivo</i> pathogenicity test of <i>Pasteurella multocida</i> B:2 mutant	123
5.2.2	<i>In vitro</i> stability test of <i>Pasteurella multocida</i> B:2 mutant	123
5.3	Results	125
5.3.1	Pathogenicity and stability test of <i>Pasteurella multocida</i>	

	B:2 mutant	125
5.4	Discussion	131
6	GENERAL DISCUSSION AND CONCLUSION	133
	REFERENCES	138
	APPENDIXES	149
	BIODATA OF STUDENT	160
	LIST OF PUBLICATIONS AND AWARD	161



This thesis was submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirements for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Siti Kharani Bejo, PhD

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Zunita Zakaria, PhD

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia

Raha Abdul Rahim, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia

AINI IDERIS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 26 September 2008



LIST OF ABBREVIATIONS

%	- percent
°C	- degree celcius
µg	- microgram
µl	- microliter
µm	- micronmeter
µM	- micromolar
Amp ^R	- ampicillin resistance
BHI	- brain heart infusion
bp	- base pair
BRENDA	- Bacterial Restriction Endonuclease Analysis
CaCl ₂	- calcium chloride
CIAP	- calf intestine alkaline phosphatase
DIG	- digoxigenin
DNA	- deoxyribonucleic acid
dNTP	- deoxynucleotide triphosphate
ds	- double-stranded
<i>E. coli</i>	- <i>Escherichia coli</i>
EDTA	- ethylene-diamine-tetraacetic acid
ELISA	- enzyme linked immunosorbent assay
F	- fertility
G	- gram
HCl	- hydrochloric acid
H ₂ O	- water
HS	- haemorrhagic septicaemia
i.p.	- intraperitoneal
IPTG	- isopropyl-β-D-thiogalacosidase
kb	- kilobase pair
Km ^R	- kanamycin resistance
LB	- Luria-Bertani
LPS	- lipopolysaccharide
L	- liter
M	- molar
MgCl ₂	- magnesium chloride
min	- minutes
ml	- milliliter
MM	- milimeter
mM	- milimolar
<i>mob</i>	- mobilization
NaCl	- natrium chloride
NADH	- nicotinamide adenine dinucleotide hydrogenase
NaOH	- natrium hydroxide
NBT/BCIP	- nitro blue tetrazolium/5-bromo-4-Chloro-3'-indolyphosphate p-toluidine salt
ng	- nanogram

nm	- nanometer
OD	- optical density
OMP	- outer membrane protein
ORF	- open reading frame
<i>ori</i>	- origin
<i>P. multocida</i>	- <i>Pasteurella multocida</i>
PBS	- phosphate buffer saline
PCR	- polymerase chain reaction
rpm	- rotation per minute
R	- resistance
RE	- restriction enzyme
RNA	- ribonucleic acid
s	- seconds
SDS	- sodium dodecyl sulphate
SSC	- sodium chloride-sodium citrate
Str ^R	- streptomycin resistance
TBE	- tris-boric EDTA
<i>Taq</i>	- <i>Thermus aquaticus</i>
T _M	- melting temperature
U	- unit
UV	- ultra-violet
V	- voltan/volt
v/v	- volume per volume
w/v	- weight per volume



Amino Acid	Single/Three Letter	Amino Acid Code
Alanine	A	Ala
Arginine	R	Arg
Asparagine	N	Asn
Aspartic Acid	D	Asp
Glutamine	Q	Gln
Glutamic acid	E	Glu
Glycine	G	Gly
Isoleucine	I	Ile
Leucine	L	Leu
Lycine	K	Lys
Methionine	M	Met
Phenylalanine	F	Phe
Proline	P	Pro
Serine	S	Ser
Threonine	T	Thr
Tryptophan	W	Trp
Valine	V	Val



LIST OF FIGURES

Figure		Page
3.1	The amplification of ABA392 virulence gene on <i>P. multocida</i> serotype B:2 produces 803bp product using lfaF2/lfaR1 primers.	58
3.2	Plasmids digestion of white colonies of the recombinants cultures with <i>EcoRI</i> at 37°C for 3 h.	59
3.3	Alignment of nucleotide sequences of ABA392 gene from <i>P. multocida</i> serotype B: 2 isolates, PMTB and 3030.	60
3.4	Alignment of protein sequences from <i>P. multocida</i> serotype B: 2 isolates, PMTB.	61
3.5	Alignment of protein sequences from <i>P. multocida</i> serotype B: 2 isolates, 3030.	62
3.6	DNA fragment patterns of <i>Pasteurella multocida</i> serotype B: 2 isolates digested by <i>HpaII</i> restriction enzyme	63
3.7	Southern hybridization of <i>Pasteurella multocida</i> serotype B: 2 genome using 803bp ABA392 gene as probe.	64
3.8	Purified ~6kb band of digested <i>Pasteurella multocida</i> serotype B: 2 genome using <i>HpaII</i> restriction enzyme.	65
3.9	PCR amplification of PCR product from ~6kb DNA template using lfaF2 and lfaR1 primers combination.	66
3.10	Plasmids digestion of white colonies with <i>EcoRI</i> at 37°C for 3 h.	67
3.11	Alignment of pPMTB and p3030 nucleotide sequences from <i>P. multocida</i> serotype B: 2 isolates, PMTB and 3030, respectively.	69
3.12	Alignment of pPMTB protein sequences from <i>P. multocida</i> serotype B: 2 isolates, PMTB.	70
3.13	Alignment of p3030 protein sequences from <i>P. multocida</i> serotype B: 2 isolates 3030.	71
4.1	Purified suicide plasmid, pAKA19.	103
4.2	Digestion of plasmid pUC4K with restriction enzyme <i>PstI</i> .	104
4.3	Purification of kanamycin cassette from plasmid pUC4K.	105
4.4	Agarose gel electrophoresis of shotgun ligation product.	106
4.5	The disruption process of the ABA392 virulence gene which was illustrated in DNA and plasmid circles.	107
4.6	Digestion of recombinant plasmid (pPMTBK19) with restriction enzymes <i>PstI</i> , <i>XhoI</i> and <i>HindIII</i> .	108
4.7	Antibiotic resistance test of <i>E. coli</i> and <i>Pasteurella multocida</i> B:2 strains.	109
4.8	Direct colony PCR of <i>Pasteurella multocida</i> B:2 mutant strains (PMTBK) and wild type <i>Pasteurella multocida</i> B:2.	111
4.9	Purification of pAKA19 suicide plasmid.	112
5.1	Direct smear from mice's heart blood after Wright's staining viewed under 100x light microscopy. Slide A and B from the group of mice which were infected with <i>P. multocida</i> B:2 mutant strains	126



- 5.2 (PMTBK) via i.p route. Direct smear from mice's heart blood after Wright's staining viewed under 100x light microscopy. Slide A and B from the group of mice which were infected with the wild type strain of *P. multocida* B:2 (PMTB) via i.p route. 127



LIST OF TABLES

Table		Page
3.1	Groups of mice were inoculated with 10 ⁸ CFU (colony forming unit) pPMTB, p3030, <i>E.coli</i> TOP 10 and pUC19 strains by i.p (intraperitoneal) route.	68
4.1	Antibiotic resistance test of <i>Pasteurella multocida</i> B:2 and <i>E. coli</i> strains.	110
5.1	<i>In vivo</i> pathogenicity test of <i>Pasteurella multocida</i> B:2 mutants.	125
5.2	<i>In vitro</i> stability test of <i>Pasteurella multocida</i> B:2 mutants.	128



CHAPTER 1

INTRODUCTION

Haemorrhagic septicaemia (HS) is an acute disease, mainly infecting cattle and water buffaloes. This disease is caused by two major serotypes of *Pasteurella multocida*, serotype B: 2 and E: 2 according to Carter-Heddleston classification, or serotype 6: B or 6: E based on Namioka-Carter classification. The first detailed report of an acute pasteurellosis, which affected deer, cattle and swine, has been attributed to Bollinger in 1878 who then reproduced the disease in cattle (Carter and de Alwis, 1989).

The outbreaks of this disease are recorded in many countries and killed many cattle and buffaloes every year (Bain *et al.*, 1982). This disease occurred in near and Middle East countries and in several parts in some regions of Africa (Carter and de Alwis, 1989). Moreover, it is also considered to be the most economically important disease of livestock in South East Asia and causes significant economic losses in India and Africa (Chandrasekaran *et al.*, 1981). The latest occurrence of HS in Malaysia was reported in Perak in the year 2005 where the disease has been confirmed to be caused by *Pasteurella multocida* serotype B:2 (Kamarudin, 2005).



HS has emerged as a disease of great economic importance. The annual losses of North American cattle industry due to HS was reported being approximately \$US800 million (Verma and Jaiswal, 1997). In Malaysia, the production of quality livestock is always a major problem since the animals are constantly fraught with diseases, especially HS which threatens the quality of products (Saharee, 2005). This situation is hardening the government's objective for livestock production to reach approximately RM 8 billion by 2010 (Mohd Nordin, 2000).

Vaccination has become the principal method of controlling HS in many countries. In Malaysia, vaccination is considered the most common and cheaper way of controlling outbreaks of HS (Zamri, 2005). Most commonly used vaccines in Malaysia are the alum-precipitated vaccine and the oil adjuvant vaccine. These vaccines did make an impact of increasing the immunity of animals against HS but still HS outbreaks and deaths remain (Dawkins *et al.*, 1990). This may be due to the facts that those vaccines provide only short-term immunity and requires annual administration for effectiveness (Chandrasekaran *et al.*, 1994).

A lot of research has been done in order to produce a better vaccine for this disease. It is widely recognized that an ideal vaccine should possess certain characteristics, such as; easy and economical to produce, stable for use in the tropics, easy to handle in the field with consistency that make it easy to administer, no adverse reactions, high level of immunity with a minimum delay