



UNIVERSITI PUTRA MALAYSIA

**ECTO-AND ENDO-PARASITES IN RODENTS AND SHREWS FROM
FOUR HABITATS IN SELANGOR AND NEGERI SEMBILAN, MALAYSIA**

PARAMASVARAN SITHAMBARAM

FPV 2009 12



**ECTO-AND ENDO-PARASITES IN RODENTS
AND SHREWS FROM FOUR HABITATS IN
SELANGOR AND NEGERI SEMBILAN,
MALAYSIA**

PARAMASVARAN SITHAMBARAM

**MASTERS OF SCIENCE
UNIVERSITI PUTRA MALAYSIA
2009**





**ECTO- AND ENDO-PARASITES IN RODENTS AND SHREWS FROM
FOUR HABITATS IN SELANGOR AND NEGERI SEMBILAN, MALAYSIA**

By

PARAMASVARAN SITHAMBARAM

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of Masters of Science**

Jan 2009

iii





Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Masters of Science.

**ECTO- AND ENDO-PARASITES IN RODENTS AND SHREWS FROM
FOUR HABITATS IN SELANGOR AND NEGERI SEMBILAN, MALAYSIA**

By

PARAMASVARAN SITHAMBARAM

Jan 2009

Chairman : Associate Professor Dr. Rehana Abdullah Sani, D.V.M., Ph.D.

Faculty : Veterinary Medicine

The objectives of this study were to determine the rodent host species, to identify the parasites they harbor and to record those parasites that are of zoonotic importance. A total of 204 rodents were examined from four selected habitats in the states of Selangor and Negeri Sembilan, Malaysia. The rodents were captured alive from five urban sites in Kuala Lumpur, rice fields in Tanjung Karang, a coastal area in Port Dickson and a Primary Forest Reserve in Gombak, Selangor. Standard laboratory parasitological and statistical analyses were used. Fourteen different rodent species belonging to three different mammalian groups were identified. *Rattus rattus diardii* was the predominant rodent species (67%) trapped and examined. The study revealed 53.9 % of the rodents examined had at least one species of helminth. A total of 21 helminth species (7 cestodes, 12 nematodes, 1 trematode and 1 acanthocephala) were recovered. *Gongylonema neoplasticum*, *Nippostrongylus braziliensis*, *Hepatojarkus malayae*, *Heterakis supomosa*, *Capillaria hepatica*,



Raillietina sp. *Hymenolepis nana*, *H. diminuta*, *Hymenolepis* sp. and *Taenia taeniaeformis* were the predominant helminth species identified. The helminths of zoonotic importance: *Capillaria hepatica*, *Rictularia tani*, *Angiostrongylus* sp., *Taenia taeniaeformis*, *Hymenolepis nana*, *H. diminuta*, *Raillietina* sp, *Moniliformis dubius*, *Armillifer moniliformis* were also found. *Trypanosoma* sp. was the only blood parasite found in the 204 rodents examined. Forty seven percent of the rodents were found to be infested with at least one of the 20 species of ecto-parasites recovered. The common ecto-parasites recovered were *Laelaps nuttali*, *L. echidnanus*, *L. sculpturata*, *Polyplax spinulosa*, *Hoplopleura pacifica*, *Ixodes granulatus*, *Dermacentor* sp. and *Haemaphysalis* sp. The ecto-parasites that are of human medical importance are *Laelaps* sp., *Gahrlepiea* sp., *Walchiella* sp., *Ornthonyssus bacoti*, *Haemaphysalis* sp., *Dermacentor* sp., *Ambloyomma* sp. and *Ixodes* sp. and *Xenopsylla cheopis*. A total of 147 ecto-parasites (ticks, n=15, fleas, n= 31, lice, n=30 and mites, n=71) were recovered from the rodents. *Rattus rattus diardii* was found to harbour a wide range of ecto and endo-parasites some of which have been identified as zoonotic. There was no significant relationship between the sex of rodents and parasitic infestation. However, significant association was seen between ecto and endo-parasitic infection and habitats. Rodents from the urban habitats comparatively were found to harbour a wider range of parasites, some of which are of human medical importance.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia, sebagai memenuhi keperluan untuk ijazah Sarjana Sains

EKTO DAN ENDOPARASIT RODENSIA DAN TUPAI DARI EMPAT HABITAT DI SELANGOR DAN NEGERI SEMBILAN, MALAYSIA

Oleh

PARAMASVARAN SITHAMBARAM

Jan 2009

Pengerusi: Associate Professor Rehana Abdullah Sani, D.V. M., Ph.D.

Fakulti: Perubatan Veterinar

Objektif kajian ini adalah untuk menentukan spesis rodensia perumah dan parasit yang dibawa olehnya dan seterusnya mengenalpasti parasit yang berkepentingan zoonotik. Sejumlah 204 rodensia dari empat habitat berlainan yang terpilih di Selangor dan Negeri Sembilan telah diperiksa. Rodensia ini telah ditangkap hidup dari lima lokasi bandar di sekitar Kuala Lumpur, sawah padi di Tanjung Karang, persisiran pantai di Port Dickson dan di Hutan Simpan Primer di Gombak, Selangor. Analisis makmal dikendalikan mengikut prosidur kajian parasitologi dan program statistik yang piawai. Empat belas spesis rodensia yang berlainan dari tiga kumpulan mamalia yang berbeza telah dikenalpasti. *Rattus rattus diardii* adalah spesis rodensia yang paling banyak diperangkap dan diperiksa (67%). Kajian ini mendapati 53.9% rodensia yang diperiksa dijangkiti oleh sekurang-kurangnya satu spesis helmin. Sejumlah 21 spesis helmin (7 sestod, 12 nematod, 1 trematod dan 1 akantosefala) telah ditemui. *Gongylonema neoplasticum*, *Nippostrongylus braziliensis*, *Hepatojarkus malayae*, *Heterakis spumosa*, *Capillaria hepatica*, *Raillietina* sp. *Hymenolepis nana*, *H. diminuta*, *Hymenolepis* sp. dan *Taenia taeniaeformis* adalah



spesis helmin yang paling dominan. Helmin yang penting dari segi perubatan: *Capillaria hepatica*, *Rictularia tani*, *Angiostrongylus* sp., *Taenia taeniaeformis*, *Hymenolepis nana*, *Hymenolepis diminuta*, *Taenia taniaeformis* juga ditemui. *Raillietina* sp, *Moniliformis dubius*, *Armillifer Moniliformis*. *Trypanosoma* sp. adalah satu-satunya parasit darah yang ditemui pada 204 rodensia yang diperiksa. Empat puluh tujuh peratus rodensia didapati diinfestasi oleh sekurang-kurangnya satu spesies dari 20 spesies ektoparasit yang ditemui. Ektoparasit akari yang paling banyak ditemui adalah *Laelaps nuttali*, *L. echidnanus*, *L. sculpturata*, *Polyplax spinulosa*, *Hoplopleura pacifica*, *Ixodes granulatus* and *Dermacentor* sp. dan *Haemaphysalis* sp. Ektoparasit yang penting dari sudut perubatan pula adalah *Laelaps* sp., *Gahrlepi* sp., *Walchiella* sp., *Ornithonyssus bacoti*, *Haemaphysalis* sp., *Dermacentor* sp., *Amblyomma* sp. and *Ixodes* sp. dan *Xenopsylla cheopis*. *Rattus rattus diardii* didapati membawa pelbagai ekto dan endoparasit, sesetengahnya zoonotik. Analisis statistik tidak menunjukkan hubungan yang berkeertian di antara jantina rodensia dan infestasi ekto dan endoparasit tetapi hubungan berkeertian didapati diantara habitat dan infestasi parasit. Rodensia dari habitat bandar didapati dijangkiti oleh pelbagai jenis species ekto dan endoparasit yang sebahagiannya adalah zoonotik berbanding dengan habitat lain.

ACKNOWLEDGEMENTS

I am deeply grateful to my supervisory committee, Associate Professor Dr. Rehana Abdullah Sani (Chairperson), Dr. Latiffah Hassan and Dr. Hanjeet Kaur Sandhu for the invaluable supervision, advice, encouragement and guidance.

I would like to express my thanks to the director of IMR and the head of Epidemiology Unit, Dr. Amal Nasir Mustaffa for their constant encouragement, support and for the allocation of the research grant. Thanks are also due to the Public Services Department (JPA) for sponsoring my studies.

I am deeply indebted and grateful to Dr. M. Krishnasamy and Mr. John Jeffery, for their constant encouragement, guidance and support throughout the study period. I also like to thank the following people: Dr. Lye Mun San, Dr. Ng Kok Han, Dr. Lee Han Lim and Dr. Indra Vythilingam for their support, encouragement and motivation.

I would like to thank the Director and staff of the Pest Control Division of Dewan Bandaraya Kuala Lumpur for the use of their facilities in Setapak and for the assistance in trapping rats in the wet markets.



Special thanks to Ms Sumarni Mohd Ghazali, Mr. Santhana Raj, Mr. Lim Kuang Hock and Mr. Palanisamy for their technical assistance.

My heartiest thanks go to my wife, son and other family members for their moral support and encouragement.

Above all my thanks are due to my spiritual guru Sri Satya Sai Baba for HIS grace and blessings.



I certify that a Thesis Examination Committee has met on 11th November 2008 to conduct the final examination of Paramasvaran s/o Sithambaram on his thesis entitled “Ecto and Endo-Parasites In Rodents From Four Habitats In The States Of Selangor And Negeri Sembilan” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia (P.U.A. (A) 106) 15 March 1998. The Committee recommends that the student be awarded the Masters of Science degree.

Members of the Thesis Examination Committee were as follows:

Dr. Saleha Abdul Aziz, Ph.D.

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Dr. Shaik Mohamed Amin S.M.Babjee, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Dr. Reuben Sunil Kumar Sharma, Ph.D.

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Dr. G. Suresh Kumar, Ph.D.

Professor
Department of Medical Parasitology, Faculty of Medicine
University Malaya
Malaysia
(External Examiner)

Dr. Bujang Kim Huat, Ph.D.

Professor and Deputy Dean
School of Graduate Studies
University Putra Malaysia



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Masters of Science. The members of the Supervisory Committee were as follows:

Rehana Abdullah Sani, D.V. M., Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Dr. Hanjeet Kaur Sandhu, M.B.B.S, M.P.H, DAP & E

Consultant Epidemiologist
Institute for Medical Research
Kuala Lumpur
(Member)

Dr. Latiffah Hassan, D. V. M., Ph.D.

Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
University Putra Malaysia

Date:



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or at any other institution.

PARAMASVARAN S/O SITHAMBARAM

Date: 12th FEB. 2009

TABLE OF CONTENTS

ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL	viii
LIST OF TABLES	xiv
LIST OF FIGURES	xv
LIST OF ABBREVIATIONS	xvi
CHAPTER	
1 GENERAL INTRODUCTION	1
2 LITERATURE REVIEW	4
2.1 INTRODUCTION	4
2.2 NEMATODES	4
2.3 CESTODES	8
2.4 TREMATODES	9
2.5 ACANTHOCEPHALAN	10
2.6 PENTASTOMID	10
2.7 ECTO-PARASITES OF RATS	10
2.8 BLOOD PARASITES OF RATS	11
2.9 BIOLOGY AND DESCRIPTION OF RATS (Muridae)	12
2.9.1 Brown Spiny Rat: <i>Rattus rajah</i> (Thomas)	12
2.9.2 Mueller's Rat: <i>Rattus muelleri</i>	13
2.9.3 Bower's rat: <i>Rattus bowersi</i> (Anderson)	14
2.9.4 Long Tail Giant Rat: <i>Rattus sabanus</i> (Thomas)	14
2.9.5 Whitehead's Rat: <i>Rattus whiteheadi</i> (Thomas)	16
2.9.6 Malaysian Woodrat: <i>Rattus tiomanicus</i> (Miller)	17
2.9.7 Polynesian Rat or Little Burmese Rat: <i>Rattus exulans</i>	18
2.9.8 House Rat: <i>Rattus rattus diardii</i> (Linnaeus)	19
2.9.9 Port-Rat: <i>Rattus norvegicus</i> (Berkenhout)	20
2.9.10 <i>Lariscus insignis</i> (Cuvier)	21
2.9.11 Slender squirrel: <i>Sundasciurus tenuis</i> (Horsfield)	21
2.9.12 Golden-Back Squirrel: <i>Callosciurus caniceps</i> (Grey)	22
2.9.13 Common Treeshrew: <i>Tupaia glis</i> (Diard)	23
2.9.14 House Shrew: <i>Suncus murinus</i> (Linnaeus)	24
2.10 PARASITIC ZOONOSES ASSOCIATED WITH RODENTS	25
3 DISTRIBUTION OF WILD AND URBAN RODENTS FROM FOUR HABITATS IN THE STATES OF SELANGOR AND NEGERI SEMBILAN	26
3.1 INTRODUCTION	26
3.2 MATERIALS AND METHODS	28
3.2.1 Identification of Species	28
3.2.2 Sex and Age	29
3.2.3 Trapping of Forest Rodents	30
3.2.4 Trapping of Urban Rodents	30
3.2.5 Trapping of Rice Field Rodents	30



3.2.6 Trapping of Coastal Rodents	31
3.3 RESULTS AND DISCUSSION	31
3.3.1 Urban Habitat	31
3.3.2 Forest Habitat	32
3.3.3 Coastal Habitat	33
3.3.4 Rice Field Habitat	33
3.4 CONCLUSION	34
4 ENDO-PARASITE FAUNA IN RODENTS AND SHREWS FROM FOUR HABITATS IN THE STATES OF SELANGOR AND NEGERI SEMBILAN	40
4.1 INTRODUCTION	40
4.2 MATERIALS AND METHODS	41
4.2.1 Trapping Sites	41
4.2.2 Collection of Samples	41
4.2.3 Data Analysis	43
4.3 RESULTS AND DISCUSSION	43
4.3.1 Urban Habitat	43
4.3.2 Forest Habitat	48
4.3.3 Rice Field Habitat	51
4.3.4 Coastal Habitat	52
4.3.5 Blood Parasites	53
4.4 CONCLUSION	54
5 ECTO-PARASITE FAUNA IN RODENTS AND SHREWS FROM FOUR HABITATS IN THE STATES OF SELANGOR AND NEGERI SEMBILAN	70
5.1 INTRODUCTION	70
5.2 MATERIALS AND METHODS	71
5.2.1 Examination of Ecto-Parasites	71
5.2.2 Data Analysis	72
5.3 RESULTS AND DISCUSSION	72
5.3.1 Urban Habitat	73
5.3.2 Forest Habitat	75
5.3.3 Rice Field Habitat	77
5.3.4 Coastal Habitat	77
5.4 CONCLUSION	78
6 SUMMARY, CONCLUSION AND RECOMMENDATIONS	85
BIBLIOGRAPHY	91
APPENDICES	102
Appendix 1	102
Appendix 2	107
Appendix 3	111
Appendix 4	112
Appendix 5	114
Appendix 6	115
Appendix 7	116
BIODATA OF STUDENT	117



**PUBLICATIONS AND PRESENTATIONS BASED ON
STUDIES RELATED TO THIS THESIS**

119



LIST OF TABLES

Table		Page
3.1	Distribution of species of rodents trapped from four habitats	35
3.2	Morphological measurements of rodents trapped in forest, rice-field, coastal areas and urban	36
4.1	The number of helminth infections in urban rats	55
4.2	Endo-parasite of rats in five urban locations sampled based on sex of rodents and infection status	56
4.3	Total number of rodents positive for helminth infections from Ulu Gombak Forest Reserve	57
4.4	Helminth infections in rodents from Ulu Gombak Forest Reserve based on species of rodents sampled	58
4.5	Helminth from <i>Rattus rattus diardii</i> from rice fields	60
4.6	Helminths from rodents trapped at the coastal areas of Port Dickson.	60
4.7	Total number of worms recovered from the rodents trapped in Port Dickson	61
4.8	Distribution of endo-parasite infection based on sex of rodents	62
4.9	Distribution of endo-parasite infection based on the habitats	62
5.1	Ecto-parasite found on rats caught in the four habitats	79
5.2	Ecto-parasite recovered from wild (forest) rodents	80
5.3	Ecto-parasites recovered from three rat host species from urban, coastal and rice field habitat	81
5.4	Flea (<i>Xenopsylla cheopis</i>) index for five urban locations	81
5.5	Distribution of ecto-parasite based on sex of rodents	82
5.6	Distribution of ecto-parasite based on the habitats	82



LIST OF FIGURES

Figure		Page
3.1	Rodent trapping locations	37
3.2	Rectangular rat trap used throughout the study	38
3.3	Rat transferred into cloth bag for anesthesia	38
3.4	Gombak Forest Reserve- one of the location for trapping rodents for the study	39
3.5	Trapped forest rodents were gathered before transferred to the Lab in IMR, KL	39
4.1	Collection of rat blood via cardiac puncture method	63
4.2	Thin and thick blood smears	63
4.3	Dissection of rats: removal of viscera organs and intestines	64
4.4	Gastro intestinal tract examination for endo-parasites	64
4.5	A: Ova of <i>Hymenolepis diminuta</i> B: Ova of <i>Hymenolepis nana</i>	65
4.6	A : <i>Armillifer moniliformis</i> (Pentastomid) B: <i>Trypanosoma</i> sp.	66
4.7	Composite picture showing external features of head, body and tail of some identified helminths Note: A= <i>Taenia taeniaeformis</i> , B= <i>Raillietina</i> sp., C= <i>Angiostrongylus</i> sp. Magnification at 20X.	67
4.8	Composite picture of the external features of head, body and tail of some helminths (x 20 Magnification) Note: A= <i>Rictularia tani</i> , B= <i>Gongylonema</i> sp., C= <i>Nippostrongylus brasiliensis</i> . Magnification at 20X.	68
4.9	Composite picture of some preserved whole worms Note: A= <i>Moniliformis dubious</i> , B= <i>Mastophorus muris</i> , C= <i>Hymenolepis</i> sp., D= <i>Gongylonema neoplasticum</i>	69
5.1	Rats combed to dislodge ecto-parasites	83
5.2	Ecto-parasite collection	83
5.3	Composite picture of a selected number of ecto-parasites (mites and chiggers) recovered from the rodents	84



LIST OF ABBREVIATIONS

cm	Centimeter
gm	Gram
kg	Kilogram
km	Kilometer
mL	Millilitre
mm	Millimeter

CHAPTER 1

GENERAL INTRODUCTION

Rodents are a key mammalian group and found in many environments throughout the world. They constitute more than 42% of the known mammalian species (Singleton, *et al.*, 2003). Rodents are a major agricultural, urban and social pest across many of the developed and developing world. In Asia alone the amount of grain eaten by rodents in the rice fields each year would provide enough to feed 200 million Asians for a year (Singleton, *et al.*, 2003). Many of these rodent species are also reservoirs of organisms that cause debilitating diseases in human and livestock (Singleton, *et al.*, 2003). Nearly four million rats are born every day in the developing nations and they are known to spread worldwide more than 60 diseases that can infect humans (Lye Hinds, 2003). Rodent-borne diseases are spread directly to humans through bite wounds, consuming food and water contaminated with rodent faeces and urine. Diseases from rodents are also spread indirectly to humans by way of ticks, mites and fleas (Nadchatram *et al.* 1967; Salleh *et al.*, 2003).

The epidemiology of many of the rodent-borne diseases in Asia and in particular Malaysia is poorly understood. For instance little is known about which species of rodents are the major reservoirs of diseases and how long does the infective stages of helminths and protozoa which originates from rodents persists in domestic and rural environments and how these agents are transmitted under natural conditions.

There is a rising concern that rodents may pose health risk to humans because of the increased movement of people between rural and urban areas and between



countries. The increased clearance of natural habitats for development further promotes rodent-human contact thus providing an ideal situation for disease transmission. The opening up of mountainous region for agricultural activities in north-eastern Tanzania brought about rodent-human interaction and the spread of plague (Makundi, *et al.*, 2003).

The aim of this study is to determine the prevalence of ecto and endo-parasitic infections in rodents found in four distinct habitats (urban, rice field, coastal and forest) in the states of Selangor and Negeri Sembilan in Peninsular Malaysia to assess the potential risk of diseases that are harmful to man and domestic animals.

The four habitats were selected for their unique ecological settings. The rice field (agriculture) urban sites represent habitats in which rodents tend to live in close proximity to human activities. The coastal site of Port Dickson was selected for its popularity for picnics, camping and other water sports recreational activities which generates food waste that attracts rodents to this habitat. This closeness provides ample opportunity for rodent to contaminate the environment thus facilitating transmission of zoonotic pathogens and parasites to humans directly or indirectly via their arthropods.

The Gombak forest reserve site was chosen for its close proximity to an Orang Asli village where the increased frequency of human activities into this forest for hunting, illegal logging and other illegal activities could have disturbed the natural habitat of the wild rodents. This study will reveal whether there has been any marked changes in the forest rodent population and their parasite diversity due to

the pressure of human activities. The information obtained from this study can also be used to assess the risk of zoonotic parasite transmission to humans and animals and also to facilitate future efforts in protecting the integrity of the primary forest's unique ecosystem and animal diversity.

The specific objectives of the study are to:

- 1) To determine the species of rodents found in four different habitats.
- 2) To identify the ecto and endo-parasites from the caught rodents and to determine the parasite species distribution.
- 3) To determine the blood parasites in the rodents.

|

Comment [P1]: deleted

These objectives will be addressed using the following hypothesis:

- 1) Rodent species diversity is dependent on habitat.
- 2) Ecto and endo-parasites found in the rodents is influenced by the location where the rodent host habituate.

CHAPTER 2

LITERATURE REVIEW

2.1 INTRODUCTION

Rodents particularly those belonging to the Muridae family in Malaysia have been well studied because of their medical and economic importance, yet comparatively, very little is known in Malaysia about the epidemiology of the ecto and endo-parasites associated with them; far less on the types of terrain in which the rodents are prevalent.

Parasites in rats (Murids) particularly helminths belonging to the four major groups; Nematoda, Cestoda, Trematoda and Acanthocephala have not been completely described or studied in Malaysia.

Comment [P2]: has deleted

2.2 NEMATODES

The earliest reports of nematodes from Malaysian murids and other animals were provided by Alessandrini (1905) and Adams (1933), who described mostly parasites from domestic animals. The nematode parasites identified from the rodent host were *Syphacia obvelata*, *Cyclodontostomum purivisi* and *Heterakis* sp. from the large intestine of rats caught in Raub. They also recorded another species of *Ancylostoma malayenum* (Alessandrini, 1905) from a rat in Seremban and an oxyurid (*Oxyurid syphacia obvelata*) in rats from Taiping and Pahang. There was a long lapse of interest following which a batch of worms from murids from Pulau

Jarak were identified by Buckely (quoted in Audy *et al.*, 1950) to contain a strongyloid. This worm was subsequently described by Yeh (1955) as *Hepatojarkus malayae*, representing a new species and genus. The nematodes *Capillaria hepatica*, *Rictularia tani* and *Nippostrongylus* sp. were also recognized in the same collection. A larger collection of parasites were examined from other parts of Malaysia around this time and it included worms from murids. These murid worms were examined by Sandosham (1954 and 1957) under the four major helminth groups in which the incidence of infection was recorded. Subsequently, nematode parasites from unidentified house rats were identified by Hall (1916). They were *Srongyloides ratti*, *Angiostrongylus cantonensis*, *Nippostrongylus muris* and *Gongylonema neoplasticum*. Four other undetermined species of *Capillaria*, *Syphacia*, *Protospirura* and *Mastophorus* were also mentioned.

A number of unnamed filarial worms were reported from rats. The first was said to be a new genus of filarial heart worm belonging to the Subfamily Splendiodfilarinae which was isolated from *Rattus sabanus*, *Rattus whitheadi* and *Rattus muelleri* by Sivanandam, *et al.* (1965). They also found another filarial worm from the heart, lung and the liver of *Rattus sabanus*. The fourth species was recovered by Sandosham and Sivanandam (1966) also from the heart of *Rattus sabanus*.

In 1963, Balasingham redescribed the hookworms of *Cyclodontostomum purvisi* from Malayan giant rats. Very few people have described the major groups of helminths in Malaysia during this time. Miyazaki and Dunn (1965) reported *Gnathostoma malaysiae* from *Rattus surifer*. First Sandosham (1957) introduced to a