



UNIVERSITI PUTRA MALAYSIA

**ASSESSMENT OF DEGENERATIVE CHANGES IN SUPERFICIAL
DIGITAL FLEXOR TENDON IN CLINICAL NORMAL HORSES**

SAMER K. TMUMEN

FPV 2005 7

**ASSESSMENT OF DEGENERATIVE CHANGES IN SUPERFICIAL DIGITAL
FLEXOR TENDON IN CLINICAL NORMAL HORSES**

By

SAMER K. TMUMEN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
In Fulfilment of the Requirement for the Degree of Doctor of Philosophy**

June 2005



DEDICATION

To my parents, Father K. Tmumen, my Mother Allah Yarhamha and my family members who encouraged me to pursue a profession I would enjoy for a lifetime.

To my wife Laila, Nouri Eteriki and my son Mohamed Samer, Taha Samer, Nagham Samer and Gufhran Samer



Abstract of the thesis submitted to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Doctor of Philosophy

ASSESSMENT OF DEGENERATIVE CHANGES IN SUPERFICIAL DIGITAL FLEXOR TENDON IN CLINICALLY NORMAL HORSES

By

SAMER K. TMUMEN

June 2005

Chairman : Professor Rasedee Abdullah, PhD

Faculty : Veterinary Medicine

The equine tendon is a highly specialized cord that ensures optimal transmission of movements from muscles to bones. The tendon is elastic, capable of easily resuming its original shape after being stretched and resistant to stretching under physiological condition. Flexor tendons show a high incidence of partial central core rupture preceded by degeneration. Thus, injuries of superficial digital flexor tendon (SDFT) are common in athletic horses.

Equine tendons were isolated from the forelimbs of 45 normal thoroughbreds consisting of 13 young, 22 middle and 10 old aged horses. The sex distribution was not taken into account. The horses were euthanised and forelimb tendon samples obtained. Gross examination showed tendon discolouration suggesting



presence of degeneration in the apparently normal tendon.

Tendon samples were obtained from the central core of the mid-metacarpal region of the forelimb SDFT. The fixed SDFT samples were processed and embedded in wax using a technique modified for the study and stained with haematoxylin and eosin (H&E) stain. All sections were examined under light, confocal laser microscopic and image analysis techniques.

When the degenerative changes in the horse forelimb SDFT were compared, the left SDFT showed more degenerative changes than the right. This reflects the local racing condition which is conducted in anti-clockwise manner. This study also showed that the degree of degenerative changes was higher ($P < 0.05$) in old horses compared to young horse, but not higher than the middle aged horses.

Tendon thickness was compared using the ultrasonography and caliper methods. The mean values for the thickness of the forelimb SDFT obtained using the caliper method (23.3 ± 1.7 mm) were at least 25% greater than those obtained using the ultrasound method (17.2 ± 2.1 mm) suggesting an over-estimation of the caliper method. Measurements using the caliper are obviously less accurate because it measures the thickness of the tendon as well as the thickness the skin and the underlying tissues. The width of the SDFT measured by ultrasonography is by transverse scans only.

The rate of healing of SDFT was compared in thoroughbreds and ponies. The SDFTs of five thoroughbreds and five ponies were injured surgically. The healing of each SDFT was monitored for six months. After six months, SDFT in ponies showed a complete healing, whereas in the thoroughbreds, granulation tissue was still clearly evident. The results showed that SDFT of ponies healed faster than that of thoroughbreds.

The tendon and muscle samples of thoroughbreds and ponies were analysed for lactic acid (LA). The tendon of thoroughbreds has a higher resting LA concentration (3.04 ± 0.42 mmol/L) than the ponies (1.45 ± 0.12 mmol/L) and increased to (7.45 ± 0.71 mmol/L) and (3.42 ± 0.89 mmol/L) in thoroughbreds and ponies. Similarly, the muscle of thoroughbreds had a higher resting LA concentration (13.63 ± 1.33 mmol/L) than the ponies (1.82 ± 0.09 mmol/L) and increased to 27 ± 3.94 mmol/L and 18.91 ± 5.35 mmol/L in thoroughbreds and ponies respectively after exercise. The increase of activity in muscle is reflected in a similar increase in the LA content of SDFT suggesting that the accumulation of LA in the tendon was the result of increased muscle LA production from increased activity. The accumulation of LA may render the tendon prone to injury.

This study suggested that tendon degeneration showed a greater intensity in the left than the right SDFT. While thoroughbreds had a greater resting tendon and

muscle LA concentration than ponies. This may be related to the fact that thoroughbreds have a greater muscle mass and higher physical activity than ponies. Plasma creatine kinase (CK) and plasma aspartate aminotransferase (AST) concentrations in horses are not good indicators of the increase in muscle activity.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENILAIAN PERUBAHAN NYAHJANA PADA TENDON FLEKSOR DIGIT
SUPERFISIAL KUDA NORMAL KLINIKALK**

Oleh

SAMER K. TMUMEN

Jun 2005

Pengerusi: Profesor Rasedee Abdullah, PhD

Fakulti: Perubatan Veterinar

Tendon ekuin merupakan suatu korda terkhusus yang memastikan pemindahan optimum pergerakan daripada otot kepada tulang. Tendon mestilah anjal, berupaya untuk kembali kepada bentuk asal dan selepas ditarik dan tahan ditarik di bawah keadaan fisiologi. Apa yang telah dicadangkan ialah tendon fleksor menunjukkan insidens tinggi untuk pecah separa pada teras pusatnya dan ini didahului dengan degenerasi. Oleh itu, kecederaan pada tendon fleksor digit superfisial (SDFT) biasa berlaku dalam kuda atletik. Lesi ini lebih kerap berlaku di kawasan metacarpus tengah pada kaki depan.

Tendon ekuin telah diperolehi daripada kaki depan 45 ekor kuda torugbred terdiri daripada 13 muda, 22 pertengahan umur, dan 10 kuda tua. Taburan jantannya tidak diambil kira. Kuda dikorbankan dan sample tendon kaki



depan diambil. Pemeriksaan kasar menunjukkan tendon bertukar warna menunjukkan wujudnya degenerasi dalam tendon yang nampaknya normal.

Sampel tendon diperolehi daripada teras pusat pada kawasan metakarpus tengah pada kaki depan SDFT. Sampel SDFT yang ditetapkan, diproses dan dibenamkan dalam lilin dengan mengguna teknik yang telah diubahsuai untuk kajian ini kemudian diwarnakan dengan pewarna hematoksilin dan eosin (H&E). Kesemua irisan telah diperiksa melalui mikroskop cahaya dan konfokal dan teknik analisis imej.

Apabila perubahan degenerasi pada SDFT kaki depan kuda dibandingkan, SDFT kiri menunjukkan lebih banyak perubahan degenerasi daripada kanan. Ini mencerminkan keadaan perlumbaan kuda tempatan di mana ianya dijalankan secara lawan arah jam. Kajian ini juga menunjukkan yang tahap perubahan degenerasi adalah lebih tinggi ($P < 0.05$) pada kuda tua berbanding kuda muda, tetapi tidak lebih tinggi berbanding kuda separuh umur.

Sukatan ketebalan tendon telah dibandingkan dengan mengguna kaedah ultrasonografi dan kaliper. Nilai min untuk ketebalan SDFT kaki depan yang diperolehi melalui kaedah kaliper (23.3 ± 1.7 mm) adalah sekurang-kurangnya 25% lebih tinggi daripada yang diperolehi melalui kaedah ultrasonografi (17.2 ± 2.1 mm) menyaranakan yang kaedah kaliper membawa kepada terlebih anggara/berlebihan. Sukatan mengguna kaliper jelas kurang tepat berbanding

ultrasonografi sebab kaliper menyukat ketebalan tendon termasuk kulit dan tisu di bawahnya sekali. kelebaran SDFT seperti diukur menggunakan ultrasonografi hanyalan pada imbasan transverse sahaja.

Kadar penyembuhan SDFT telah dibandingkan dalam kuda torugbred dan padi. Dalam kajian ini, SDFT daripada lima ekor kuda torugbred dan lima ekor kuda padi dcederakan secara surgeri. Penyembuhan SDFT ini dimantau selama 6 bulan. Selepas 6 bulan SDFT kuda padi sembuh sepenuhnya, sambil dalam kuda torugbred tisu penggranulan masih lagi jelas. Hasil kajian ini menunjukkan yang SDFT kuda padi sembuh lebih cepat daripada kuda torugbred.

Sampel tendon dan otot kuda torugbred dan padi telah dianalisiskan untuk kandungan asid laktiknya. Otot kuda torugbred mempunyai kepekatan asid laktik rihaat (13.63 ± 1.33 mmol/L) yang lebih tinggi daripada kuda padi (1.82 ± 0.09 mmol/L) dan meningkat kepada 27.06 ± 3.94 mmol/L dan 18.91 ± 5.35 mmol/L masing-masing dalam kuda torugbred dan kuda padi selepas latihan. Peningkatan asid laktik dalam otot mencerminkan sebagai peningkatan pada masa sama kandungan asid laktik SDFT, menyarankan yang pengumpulan asid laktik dalam tendon merupakan hasil daripada peningkatan asid laktik otot kerana peningkatan aktiviti. Pengumpulan asid laktik mungkin menyebabkan tendon lebih mudah cedera.

Kajian ini meyarankan bahawa degenerasi tendon menunjukkan intensity yang lebih pada SDFT kiri daripada SDFT kanan. Kuda torugbred mempunyai kepekatan asid laktik riha yang lebih tinggi daripada kuda padi. Ini mungkin berhubungkait dengan hakikat bahawa kuda torugbred mempunyai kepadatan otot dan aktiviti fizikal yang lebih tinggi daripada kuda padi. Kepekatan creatine kinase (CK) dan aspartate aminotransferase (AST) bukanlah petunjuk yang baik untuk peningkatan aktiviti otot pada kuda.



ACKNOWLEDGMENTS

I am grateful to my supervisory committee chairman, Professor Dr. Rasedee Abdullah for his valuable advice and guidance throughout this study. My sincere thanks to Professor Dr. Mohd Zamri Saad, for his assistance in preparation of this thesis, Dr. Mohd Zuki Abu Baker for his guidance in ultrasonography, and Dr Nadzri Salim for his assistance in the statistical analysis. My deepest appreciation goes to Dr. Bashir Ahmed for providing the facility to conduct the experiments.

My deepest thanks to Professor Dato Dr. Sheikh Omar Abdul Rahman for his good explanation and helpful. Associate Prof. Dr. Mohd Hair Bejo for his kindness and helpful, also like to express my gratitude and appreciation to the following people who have been of a great help in my study: Mrs. Sairah Binti Abdul Karim. Dr. Sabri, Mohd Yusoff. Mr. Salehuddin Abd Rahman, Mr. Zain, and Mrs. Noraniza Mohd. Adzahan of the UPM Equine Center; Mr. Mohamed Halmi Othman and Mr. Abdullah Misron of the Haematology and Clinical Biochemistry Laboratory; Mr. Mohd Noh, Mr. Jamil Abdul Samad of Histopathology Laboratory; Mrs Fareda , Ms. Azilah Abul Jalil, Mr. Ho Oi Kuan of the Imaging Unit, Institute of Bioscience.

All members of the Faculty of Veterinary Medicine for sharing their technical skills and their ever available assistances.



Last but not least is my gratitude to the ministry of education in Libya for providing the opportunity and scholarship to pursue the degree. This study is dedicated to my wife Laila N. Eteriki and my juniors Mohamed Samer, Taha Samer, Nagham Samer and Gofhran Samer for being understanding and supportive throughout my study.



I certify that an Examination Committee met on 17th June 2005 to conduct the final examination of Samer K. Tmumen on his Doctor of Philosophy thesis entitle "Assessment of Degenerative Changes in Superficial Digital Flexor Tendonin Clinically Normal Horses" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Tengku Azmi Tengku Ibrahim, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Dato' Sheikh Omar Abdul Rahman, PhD


Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Internal Examiner)

Dato' Abdul Salam Abdullah, PhD

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Internal Examiner)

Takashi Umemura , PhD

Professor
Hokkaido University
Japan
(External Examiner)



GULAM RUSUL/RAHMAT ALI, PhD

Professor/ Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 Jul 2005



The thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as following:

Rasedee Abdullah, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Mohd. Zamri Saad, PhD


Professor
Faculty of Veterinary Medicine
University Putra Malaysia
(Member)

Mohd Zuki Abu Baker, PhD

Lecturer
Faculty of Veterinary Medicine
University Putra Malaysia
(Member)

Nadzri Salim , MPVM

Lecturer
Faculty of Veterinary Medicine
University Putra Malaysia
(Member)



AINI IDERIS, PhD
Professor/ Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **11 AUG 2005**



DECLARATION

I hereby declare that the thesis is based on my original work except for quotation and citation, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



SAMER K. TMUMEN

Date: 20/7/2005

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xiii
DECLARATION	xv
LIST OF FIGURES	xx
LIST OF TABLES	xxi
LIST OF PLATES	xxii
LIST OF ABBREVIATIONS AND SYMBOLS	xxvi
 CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	6
Anatomy	6
Structure of Tendon	6
Collagen fibers	9
Blood supply to the tendon	11
Tendon function	12
Physiological loading of tendon	13
Tendon Injuries	14
Degenerative change of tendon	15
Equine Tendinitis	18
Definition	18
Etiology of Tendinitis	18
Pathogenesis of Equine Tendinitis	18
Clinical Signs	21
Diagnosis of Tendinitis	21
Treatment of Tendinitis	22
Equine Lameness	24
Lameness	24



	Etiology of Lameness	24
	Classification of Lameness	25
	Epidemiology of Lameness	27
	Diagnosis of Lameness	28
III	DEGENERATIVE CHANGES OF THE SUPERFICIAL DIGITAL FLEXOR TENDON IN CLINICALLY NORMAL HORSE	29
	Introduction	29
	Material and Methods	32
	Horses	32
	Tissue sampling	32
	Gross Pathological examination	32
	Sampling processing	33
	Water content	33
	Tissue fixing	33
	Tissue sectioning and staining	33
	Modification of fixation and processing of tendon	34
	Microscopic examination	34
	Image Analysis	34
	Determination of degenerative area in the tendon	35
	Scanning Electron Microscopy (SEM) Techniques	35
	Confocal Microscopy	36
	Statistical Analysis	36
	Results	37
	Water content	37
	Gross Pathology	37
	Tendon morphological characteristic	42
	Light Microscopy	42
	Electron Microscopy (SEM)	44
	Confocal Microscopy	48
	Image analysis	50
	Light Microscopy.	53
	Extent of SDFT degeneration in different age groups	54
	Extent of total SDFT degeneration in all horses	55
	Image analysis	55
	Extent of SDFT degeneration in different age groups	55
	Extent of total SDFT degeneration in all horses	56
	Comparison between light microscopy and image analysis in the estimation of SDFT degeneration	56



Discussion.	58
Conclusion	64
IV ULTRASONOGRAPHIC MEASUREMENTS OF THE THICKNESS OF EQUINE SUPERFICIAL DIGITAL FLEXOR TENDON AND EVALUATION OF THE HEALING PROCESS OF SURGICALLY-INDUCED TENDON INJURY	65
Introduction	65
Materials and Methods	71
Horses	71
Animal preparation	71
Caliper measurement of SDFT width	71
Ultrasonographic examination of the SDFT	72
Surgical procedure to induce tendon injury	74
Ultrasonography examination of injured site	75
Post-operative care	75
Statistical Analysis	76
Results	77
Ultrasonographic characteristics of normal SDFT	77
Assessment of SDFT thickness	80
Ultrasonographic examination of the surgical site	81
Ultrasonographic of the healing process in surgery-induced Tendon injury	81
Discussion	92
Conclusion	96
V LEVEL OF LACTIC ACID, CREATINE KINASE AND ASPARTATE AMINOTRANSFERASE IN THE SUPERFICIAL DIGITAL FLEXOR TENDON IN CLINICALLY NORMAL THOROUGHBREDS AND PONIES	97
Introduction	97
Material and Methods	101
Horses	101
Training Program	101
Blood Samples	101
Tendon biopsy	102
Muscle biopsy	103
Biochemical analysis	103
Post-operative management	103
Statistical analysis	104



Results	105
Plasma lactate activity in thoroughbreds and ponies	105
Plasma CK activity in Thoroughbred and ponies in exercise	106
Plasma AST activities in Thoroughbred and ponies in Exercise	107
Lactic acid concentration in SDFT of horses	109
Lactic acid evaluation of muscle of horses	110
Discussion	111
Conclusion	116
VI GENERAL DISCCUSION AND CONCLUSION	117
SUMMARY	124
BIBLOGRAPHY	125
BIODATA OF THE AUTHOR	146



LIST OF FIGURES

Figures		Page
1	The hierarchical organization of tendon structure: from collagen fibrils to the entire tendon	8
2	Possible pathways for tendon degeneration.	17
3	Showing the stages of superficial digital flexor tendon ruptures or damages	20

LIST OF TABLES

Tables		Page
1	Total number and group of horses used in the study	32
2	The extent of SDFT degeneration by light microscopy and image analysis	53
3	Statistical composition of tendon degeneration as determined by light microscopy and image analysis	57
4	The SDFT thickness determined by the caliper measurement and ultrasonography	80
5	Concentrations of plasma lactic acid in the exercised thoroughbreds and ponies	106
6	Concentrations of plasma creatine kinase in the exercised thoroughbreds and ponies.	107
7	Concentrations of plasma Aspartate aminotransverase in exercise thoroughbreds and ponies	108
8	Concentrations of lactic acid in the SDFT of exercised thoroughbreds and ponies	109
9	Concentrations of lactic acid in the gluteal muscle of exercised thoroughbreds and ponies	110



LIST OF PLATES

Plates		Page
1	Transverse section of clinically normal equine SDFT Showing reddish discoloration (arrow) in the central core	38
2	Transverse section of normal equine SDFT Showing homogenous structures with straw-like colouration (arrows)	38
3	Abnormal structure of SDFT of transverse section with calcified materials (arrows) at the periphery.	39
4	The SDFT of the left forelimb showing the adherence of SDFT to the other collagenous structures, DDFT and inferior chick ligament forming a single structure.	39
5	Longitudinal section of SDFT (1) adhering to the DDFT (2) and Inferior check ligament ICL (3) as one structure. The facia and paratenon were also adhesive (arrows)	40
6	The SDFT (1), DDFT (2) and ICL (3) of the supporting forelimb showing the adhesive structures (arrows)	40
7	The SDFT (1) and DDFT (2) showing the thickened structures (arrows).	41
8	SDFT (1) and DDFT (2) showing thickened structures and atrophy (arrows).	
9	Histological section of normal SDFT showing the arrangement of fibre bundles (arrows) carrying with tenocytes in-between the fibres (X100) (H&E).	43
10	Histological section of the normal SDFT showing homogeneous structure (arrow), absence of tenocytes and destruction of collagen bundles (x40) (H&E).	43
11	Cross-section area of the normal SDFT showing internal tendon structures showing small and large bundles under scanning electron microscopy (arrows).	45
12	Longitudinal section of the normal SDFT showing irregularly arranged bundle of fibres under scanning electron microscopy	45
13	Transverse section of SDFT showing the internal blood supply of the tendon (arrows) x200 under scanning electron microscopy.	46

14	Cross-section of normal SDFT showing the wave pattern which appears as zig.zag conformation (arrow A) the cracks are artifacts (arrow B) x300 under scanning electron microscopy	46
15	Scanning electron microscopic image of SDFT showing degenerative area characterized as a black zone with smooth homogenous surface. x95 (arrow).	47
16	Scanning electron microscopic image of the equine SDFT showing degenerative area characterized as a black homogenous area. x160	47
17	Normal SDFT engorged with blood supply (arrow) under confocal microscopy 10 μ , H&E.	49
18	Confocal macrograph of extensive homogeneous area of degeneration changes of SDFT under confocal microscope 10 μ H&E(arrows).	49
19	SDFT showing an excessive area of degenerative change (arrows) under image analysis technique (x100 H&E).	50
20	Degenerative changes in SDFT shown by image analysis (arrows) (x100) (H&E).	51
21	Normal SDFT showing a blood vessel (arrows) under image analysis techniques x100 μ m	51
22	A section of SDFT showing degenerative change represented by extensive homogenous area (arrows)	52
23	Degenerative change in Plate 22 after shadowing with red colouration (arrows).	52
24	The caliper used for the measurement of SDFT thickness	72
25	The ultrasound machine (TOSHIBA just vision 200) used in the study.	73
26	The forelimb of the horse showing the surgical site and the exposed SDFT	75
27	Longitudinal (A) and transverse (B) scans of the normal SDFT at mid-metacarpal region using 8.0MHz transducer. The longitudinal scan of SDFT shows a multiple, regularly arrange parallel echoes, while transverse scan shows an homogenous echogenicity scattered throughout the tendon parenchyma.	78

28	Longitudinal (A) and transverse scan (B) of normal SDFT using an 8MHz transducer showing the thickness of the tendon and the boundary demarcating the tendon. Note that the tendon structures are similar for SDFT, DDFT and ICL	79
29.A	Transverse scans of the pony of the left SDFT week 1 post-surgery, showing disruption in the fiber pattern and inflammatory changes was presented (arrow).	84
29.B	Transverse scans of the thoroughbred of the left SDFT week 1 post-surgery, showing a decrease in echogenicity. Note that the pattern of SDFT was ruptured due to injury. (arrow).	84
30.A	Transverse scans of pony of left SDFT week 2 and 3 post-surgery, showing hypoechogenic area and disruption of tendon fiber (arrow).	85
30.B	Transverse scans of thoroughbred SDFT in week 2 and 3 post-surgery, showing a typical echogenic spots in the region (arrow).	85
31.A	Transverse scans of pony SDFT in week 4 and 5 post-surgery, showing decrease of echogenicity in the area (arrow).	86
31.B	Transverse scans of thoroughbred SDFT in week 4 and 5 post-surgery, showing the fiber pattern was disrupted and poor healing process (arrows)	86
32.A	Transverse scans pony SDFT in week 6 post-surgery, showing the granulation tissue was beginning to fill the gap in tendon (arrow).	87
32.B	Transverse scans of thoroughbred SDFT in week 6 post-surgery, showing poor formation of granulation tissue (arrow).	87
33.A	Transverse scans of pony SDFT in week 7 and 8 post-surgery, showing decrease in homogeneity of the injured area of tendon.(arrow).	89
33.B	Transverse scans of thoroughbred SDFT in week 7 and 8 post-surgery, showing a slight regeneration of tendon fiber (arrows).	89
34.A	Transverse scans of pony of the left SDFT in week 9 and 10 post-surgery, showing formation of granulation tissue. The tendon was healing rapidly (arrows).	90