



UNIVERSITI PUTRA MALAYSIA

FUNCTIONAL CHARACTERIZATION OF YIRO10W (DSN1) AND YOR228C GENES IN SACCHAROMYCES CEREVISIAE

YIAP BEOW CHIN.

FSMB 2004 15



FUNCTIONAL CHARACTERIZATION OF YIR010W (DSN1) AND YOR228C GENES IN Saccharomyces cerevisiae.

Disertai CD-ROM / disket yang boleh diperolehi di Bahagian Media dan Arkib (Accompanying CD-ROM / disk available at the Media and Archives Division)

By

YIAP BEOW CHIN

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in Fulfilment of Requirement for the Degree of Doctor of Philosophy.

December 2004



To my parents, wife, son and daughter.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

FUNCTIONAL CHARACTERIZATION OF YIR010W (DSN1) AND YOR228C GENES IN Saccharomyces cerevisiae.

By

YIAP BEOW CHIN

December 2004

Chairman: Associate Professor Raha Abdul Rahim, Ph.D.

Faculty: Food Science and Biotechnology

The aim of this project was to functionally decipher two novel genes of *Saccharomyces cerevisiae* (YOR228C and YIR010W). Four levels of study were undertaken to achieve the aims of this study including bioinformatics data mining and prediction, gene deletion study, phenotypic analysis and protein profiling. Sequence analysis of YOR228C predicted the expression of a 34 kDa transmembrane protein of low abundance with the potential of forming a homodimer. It could be a stress-responsive gene that also regulates cell growth in yeast but it was not an essential component of the cell. On the other hand, Yir010wp has a size of 66 kDa and would be a heterodimer in a complex. This low abundant protein may be related to cell division cycle as it was found to be incorporated within the yeast spindle pole body. The necessity for its exact stoichiometry was a sign of gene dosage sensitivity as demonstrated by the reduction in heterologous fitness. Deletion of a single copy of *YIR010W* caused chromosomal segregation error leading to aneuploidy. Furthermore, double knockout of the gene from the genome was



lethal, implying its essentiality to the yeast cell. As a whole, this study has successfully elucidated the general functions of the two genes under investigation, namely the relationship between *YOR228C* to growth/stress-response and the link between *YIR010W* to the cell division cycle.



PENCIRIAN FUNGSI GEN-GEN YIR010W (DSN1) DAN YOR228C DALAM Saccharomyces cerevisiae.

Oleh

YIAP BEOW CHIN

DISEMBER 2004

Pengerusi: Profesor Madya Raha Abdul Rahim, Ph.D.

Fakulti: Sains Makanan dan Bioteknologi.

Tujuan projek ini adalah untuk mengetahui fungsi dua gen yis Saccharomyces baru jaitu YOR228C dan YIR010W. Empat tahap cerevisiae yang penyelidikan telah dijalankan untuk mencapai matlamat projek ini, termasuk pencarian data dan ramalan menggunakan bioinformatik, penyelidikan pemadaman gen, analisis fenotip dan pemprofilan protein. Analisis jujukan YOR228C diramalkan untuk menghasilkan satu protein transmembran bersaiz 34 kDa dan mempunyai jumlah yang sedikit serta berpotensi menjadi homodimer. Ia berkemungkinan adalah suatu gen yang bertindakbalas terhadap tekanan dan juga regulasi pertumbuhan sel yis, tetapi ia adalah komponen yang tidak penting untuk sel. Sebaliknya, Yir010wp mempunyai saiz 66 kDa dan boleh menjadi heterodimer dalam suatu kompleks. Protein berjumlah kecil ini sememangnya berkaitan dengan proses penduaan sel apabila ia didapati melekat kepada badan kutub spindal (spindle pole body) yis. Keperluannya dalam stoikiometri yang betul merupakan suatu tanda kepekaan terhadap dos gen sebagaimana ditunjukan oleh kekurangan



kecergasan heterologus. Permadaman satu gen *YIR010W* menyebabkan kesilapan dalam pembahagian kromosom dan seterusnya menghasilkan aneuploidi. Tambahan pula, pemadaman kedua-dua gen tersebut boleh membinasakan sel yis; menunjukkan kepentingannya terhadap sel tersebut. Secara keseluruhannya, projek ini telah berjaya menerangkan secara ringkas fungsi kedua-dua gen yang dikaji, di mana *YOR228C* merupakan gen yang berkaitan dengan tindakbalas terhadap tekanan dan pertumbuhan manakala *YIR010W* berhubungan dengan proses penduaan sel.



ACKNOWLEDGEMENTS

I hereby wished to take this opportunity to express my sincere appreciation to all the people who have helped me in my pursuit of Ph.D.

First and foremost is my supervisor Assoc. Prof. Dr. Raha Abdul Rahim whose supervision and support was essential for the successful completion of this work. I would like to present my greatest thanks to Dr. Hirzun Mohd. Yusuf for providing me with a chance to be involved in this study. I would also like to thank Dr. Clemente Michael Wong Vui Ling for his constant guidance and advice, not forgetting also Prof. Dr. Son Radu for his precious comments.

Also many thanks to my fellow lab-mates Varma, Hossein, Prema, Ernie, Chyan Leong, Li Lung, Hooi Ling, Yanti, Boon Hooi, Sabrina, Azlina, Mastura, Sharul, Baktiar. Genetic Lab.- Assoc. Prof. Dr. Harikrisna, Assoc. Prof. Dr. Jenni Harikrisna, Dr. Tan Siang Hee, Dr. Ho Chai Ling, Mr. Ong, Chong, Weng Wah, Pick Kuen, Yang Ping and all the friends in that lab. ATCL – Prof. Dr. Manaf, Boon Keat, Anthony Ho, Yang Mooi, Suhaimi and Cheng Leng. Virology Lab. – Mr Wong, Majjit. FTU – Assoc. Prof. Dr. Arbakariaya, Dr. Rosfarizan, Rizal, Sobri and all the friends in the unit. Also to my friends Fei Ling, Ja, Pey Juin, Zul, Fadli, Richard and Ah Miao.

Finally, I would like to devote my utmost gratitude to my parents and family members for their unfailing love and constant support. And to my wife, Wai Ying, my son, Vincent and my daughter Vivian, I love you all.



vii

My Ph.D studentship was funded by National Science Fellowship (NSF) from Ministry of Science, Technology and Innovation (MOSTI), Malaysia.



I certify that an Examination Committee met on 3rd September 2004 to conduct the final examination of Yiap Beow Chin on his Doctor of Philosophy thesis entitled "Functional Characterization of YIR010W (DSN1) and YOR228C Genes in *Saccharomyces cerevisiae*" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Ho Chai Ling, Ph.D.

Faculty of Biotechnology and Biomolecular Sciences Universiti Putra Malaysia (Chairman)

Norihan Mohd. Saleh, Ph.D.

Associate Professor Faculty of Biotechnology and Biomolecular Sciences Universiti Putra Malaysia (Member)

Foo Hooi Ling, Ph.D.

Associate Professor Faculty of Biotechnology and Biomolecular Sciences Universiti Putra Malaysia (Member)

Akihiko Kondo, Ph.D.

Professor University of Kobe (Independent Examiner)

GULAM RUSUL RAHMAT ALI, Ph.D. Professor/Deputy Dean School of Graduate Studies Universiti Putra Malaysia

Date: 30 NOV 2004



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The Members of the Supervisory Committee are as follow:

RAHA ABDUL RAHIM, Ph.D

Associate Professor Faculty of Biotechnonology and Biomolecular Sciences Universiti Putra Malaysia. (Chairman)

SON RADU, Ph.D Professor Faculty of Food Sciences Universiti Putra Malaysia. (Member)

CLEMENTE MICHAEL WONG VUI LING, Ph.D Universiti Malaysia Sabah. (Member)

AINI IDERIS, Ph.D, Professor/ Dean School of Graduate Studies Universiti Putra Malaysia

Date: 10 DEC 2004



I hereby declare that the thesis is based on my original work except for the quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

YIAP BEOW CHIN

Date: 8 12 2004



TABLE OF CONTENTS

		Page
DED	ICATION	ii
ABS	TRACT	iii
ABS	TRAK	v
ACK	NOWLEDGEMENTS	vii
APP	ROVAL	ix
DEC	LARATION	xi
LIST	OF TABLES	xvi
LIST	OF FIGURES	xx
LIST	OF ABBREVIATIONS	xxvi
СНА	PTER	
1	INTRODUCTION	
	1.1 Introduction	1
	1.2 Objectives	3
2	LITERATURE REVIEW	
	2.1 Yeasts	4
	2.1.2 Saccharomyces cerevisiae	5
	2.2 Phenotypic study through mutagenesis	8
	2.3 Bioinformatics	10
	2.4 Yeast Enhanced Green Fluorescent Protein (yEGFP)	12
	2.5 Light Microscopy Application For Yeast	14
	2.5.1 Autofluorescence	16
	2.6 Dosage Sensitivity and Balance Hypothesis	19
	2.7 YOR228C	
	2.7.1 Gene Information	21
	2.7.2 Protein Information	23

2.8 YIR010W

2.8.1 Gene information

2.8.2 Protein information	32
BIOINFORMATICS	
3.1 Introduction	42
3.2 Materials & Methods	43
3.3 Results	
3.3.1 mRNA expression level of YOR228C under	er 44
different conditions 3.3.2 mRNA expression level of YIR010W unde	• •
different conditions 3.4 Discussion	52 59
3.4 DISCUSSION	55
GENE DELETION STUDIES	
4.1 Introduction	65
4.2 Materials & Methods	
4.2.1 Polymerase Chain Reaction (PCR)	65
4.2.2 Transformation of Saccharomyces cerevisiae	66
4.2.3 Diploid Formation	67
4.2.4 Sporulation	67
4.2.5 Tetrad Dissection	68
4.3 Results	
4.3.1 Deletion of YOR228C gene	68
4.3.2 Deletion of YIR010W gene	74
4.3.3 Confirmation of deletion via PCR reaction	79
4.3.4 Tetrad Analysis	89
4.4 Discussion	95
GROWTH PERFORMANCE STUDY OF DELETANTS	
5.1 Introduction	99
5.2 Materials & Methods	
5.2.1 Solid media	100

- 5.2.2 Liquid culture
- 5.3 Results

	5.3.1 Growth study on different media and temperatures	101
	5.4 Discussion	127
6	LOCALIZATION AND EXPRESSION STUDY OF GFP- YIR010W (DSN1) FUSION PROTEINS IN VIVO 6.1 Introduction	135
	6.2 Materials & Methods	100
	6.2.1 Preparation of <i>S.cerevisiae</i> Genomic DNA	136
	6.2.2 Plasmid DNA isolation	137
	6.2.3 Polymerase Chain Reaction (PCR)	138
	6.2.4 Restriction endonuclease digestion of DNA	139
	6.2.5 Purification of digested samples and PCR product	139
	6.2.6 DNA ligation	139
	6.2.7 Sequencing	140
	6.2.8 DNA electrophoresis	140
	6.2.9 SDS – PAGE protein analysis	141
	6.2.10 Western blotting	142
	6.2.11 Preparation of <i>E.coli</i> competent cells	143
	6.2.12 Heat shock transformation method	143
	6.3 Results	
	6.3.1 Cloning strategy	144
	6.3.2 Confirmation of successful cloning	152
	6.3.3 Expression study of YIR010W-GFP fusion protein	158
	6.3.4 Growth assay	166
	6.4 Discussion	171
7	MICROSCOPIC STUDY OF CELLULAR MORPHOLOGIES IN HOSTS CARRYING DIFFERENT COPY NUMBERS OF GENES OF INTEREST 7.1 Introduction	181
	7.2 Materials & Methods	101
		182
	7.2.1 Microscopy sample preparation	102

- 7.2.2 Nuclear staining with DAPI 183
- 7.3 Results



	7.3.1 YOR228C	183
	7.3.2 YIR010W	198
	7.4 Discussion	224
8	GENERAL DISCUSSION	
	8.1 YOR228C	230
	8.2 YIR010W	232
	8.3 Conclusion	234
BIBLIOGRAPHY 23		236
APPENDIX		
Α	Composition of media (per litre) and antibiotic stock solutions	244
В	Solutions for protein electrophoresis of SDS – PAGE	246
С	Yeast Strains	247
D	Plasmids	252
Е	Primers	253
F	Chromatograms	255
G	Bioinfomatic reference results of Chapter 2 And Chapter 3	257
BIOD	ATA OF THE AUTHOR	347



Table		Page
2.1	YOR228C protein information obtained from various	26
	databases.	
2.2	Summary of predictions on compartmentalization, target sites	
	for post translational modification, protein domain and	
	secondary structure of YOR228C by different databases.	27
2.3	Summary of multi-level comparative proteomic for YOR228C	30
	protein.	
2.4	YIR010W protein information obtained from various databases.	33
2.5	Summary of protein interaction search result for YIR010W from	
	The GRID Matching.	35
2.6	Summary of predictions on compartmentalization, targets sites	
	for post translational modification, protein domain and	
	secondary structure of YIR010W by different databases.	36
2.7	Summary of multi-level comparative proteomic for YIR010W	
	protein.	40
3.1	mRNA expression levels for YOR228C.	46
3.2	Summary (from appendix 3.10) of microarray data of	
	YOR228C mRNA level in the presence of different DNA	
	damaging agents.	50
3.3	Summary (from Appendix 3.11a-d) of microarray data of	
	YOR228C mRNA level under different environmental	
	conditions.	51
3.4	mRNA expression levels for YIR010W.	53



xvi

- 3.5 Summary (from appendix 3.14(I), (II)) of microarray data of *YIR010W* mRNA level in the presence of different DNA damaging agents.
- 3.6 Summary (from appendix 3.15(I) (II)) of microarray data of YIR010W mRNA level under different environmental conditions.
- 4.1 Primer pair combinations from the 5 different primers giving different additional number of nucleotides in PCR products and their expected sizes.
- 5.1 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in YPD agar plates at 15°C, 30°C and 37°C.
 104
- 5.2 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in YPG agar plates at 15°C, 30°C and 37°C.
 106
- 5.3 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in SD minimal agar plates at 15°C, 30°C and 37°C.
 107
- 5.4 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in YPD agar plates with various Calcium concentrations at 30°C and 37°C.
 108
- 5.5 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in YPD agar plates with various Magnesium concentrations at 30°C and 37°C.
 110



56

58

- 5.6 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in YPD agar plates with various Sodium concentrations at 30°C and 37°C.
 111
- 5.7 Comparison of wildtypes and disruptants of YOR228C in different genetic backgrounds growing in YPD agar plates with Amphotericin B (100 ng/ml) and Lovastatin (40 μg/ml) at 30°C.
- 5.8 Comparison of wildtypes and disruptants of YIR010W in different genetic backgrounds growing in YPD, YPG and SD minimal agar plates at 15°C, 30°C and 37°C.
 123
- 5.9 Comparison of wildtypes and disruptants of *YIR010W* in different genetic backgrounds growing in YPD agar plates with various Calcium, Magnesium and Sodium concentrations at 30°C and 37°C.
- 5.10 Comparison of wildtypes and disruptants of YIR010W in different genetic backgrounds growing in YPD agar plates with Amphotericin B (100 ng/ml) and Lovastatin (40 μg/ml) at 30°C 126
- 7.1 The average cell size of different host from both wild types and disruptants of *YOR228C*.185
- 7.2 The budding index of different host from both wild types and disruptants of YOR228C.195
- 7.3 The percentage of golden autofluorescence cell (from the overall population) of different host from both wild types and disruptants of YOR228C.
 200
- 7.4 The average cell size of different host from both wild types and



124

xviii

	disruptants of YIR010W.	204
7.5	The budding index of different host from both wild types and	
	disruptants of YIR010W.	211
7.6	The percentage of golden autofluorescence cell (from the	
	overall population) of different host from both wild types and	
	disruptants of YIR010W.	217



LIST OF FIGURES

Figure		Page
4.1	Sequence of the long chimeric primers designed for the	
	deletion of gene YOR228C.	69
4.2	PCR reaction using the long chimeric primers to amplify the	
	KanMX4 deletion cassette from the plasmid pFA6a-KanMX4.	71
4.3	Gel photo showing PCR product (1.6 kb) of the deletion	
	construct in lane L1.	72
4.4	Diagrammatic presentation of homologous recombination of	
	the deletion construct (KanMX4 module) replacing one copy	
	of the gene (YOR228C) in a diploid genome upon	
	transformation into the cell.	73
4.5	Production of double PCR products by primer set F339W &	
	R170W.	75
4.6	PCR product of deletant and wild type.	76
4.7	Diagrammatic presentation of homologous recombination of	
	the deletion construct (URA3 module) replacing one copy of	
	the genes (YIR010W) in a diploid genome upon	
	transformation into the cell.	78
4.8	PCR verifications utilizing primer pairs.	80
4.9	PCR product of yor228c deletant and wildtype control.	81
4.10	Primer pair FC228C/RC228C was used for checking the	
	integrity of the insertion site/ flanking homologous regions	
	and primer pair F450C/RK500C was used to check the	



	integrity upstream and downstream.	83
4.11	PCR verification of single and double deletion in both haploid	
	and diploid from non – deletion control.	84
4.12	The 5 primers (3 forwards & 2 reverses) that made up the 6	
	primer pairs being used for confirmation of insertion of	
	deletion construct into the YIR010W locus.	86
4.13	Comparison of different sizes of PCR products of yir010w	
	single deletant and wild type control.	87
4.14	Yeast tetrad upon sporulation.	90
4.15	Tetrad dissection of yor228c deletant.	91
4.16	Drawing showing testing of the mating type of an unknown	
	haploid zygote.	93
4.17	Tetrad dissection of yir010w deletant.	94
5.1	Qualitative scoring of yeast growth performance.	103
5.2a	Graphs showing the comparisons of growth rate using OD_{600}	
	reading between wildtype control yeast cell against	
	YOR228C disruption mutants in YPD at 30°C.	115
5.2b	Graphs showing the comparisons of growth rate using OD_{600}	
	reading between wildtype control yeast cell against	
	YOR228C disruption mutants in YPD at 37°C.	116
5.3a	Graphs showing the comparisons of growth rate using OD ₆₀₀	
	reading between wildtype control yeast cell against	
	YOR228C disruption mutants in SD Minimum at 30°C.	117
5.3b	Graphs showing the comparisons of growth rate using OD ₆₀₀	
	reading between wildtype control yeast cell against	



YOR228C disruption mutants in SD Minimum at 37°C. 118

- 5.4a Graphs showing the comparisons of growth rate using OD₆₀₀
 reading between wildtype control yeast cell against
 YOR228C disruption mutants in YPG at 30°C.
- 5.4b Graphs showing the comparisons of growth rate using OD₆₀₀
 reading between wildtype control yeast cell against
 YOR228C disruption mutants in YPG at 37°C.
 120
- 5.5a Graphs showing the comparisons of growth rate using OD₆₀₀
 reading between wildtype control yeast cell against *YIR010W*disruption mutants in YPD at 30°C and 37°C.
- 5.5b Graphs showing the comparisons of growth rate using OD₆₀₀
 reading between wildtype control yeast cell against *YIR010W*disruption mutants in YPG at 30°C and 37°C.
- 6.1 PCR product of YIR010W-GFP cloning primers (both Cterminal and N-terminal were the same) showing size of 1749 bp.
- 6.2 Gene orientation for YIR010W and EGFP3 in both C-terminal and N-terminal fusion together with the cutting sites for restriction enzymes (Smal and EcoRI) and Factor Xa.
- 6.3 Diagrammatic presentation of a plasmid map for a C-terminal GFP fusion vector pUG23.
- 6.4 Diagrammatic presentation of a plasmid map for a N-terminalGFP fusion vector pUG34. 148
- 6.5 Diagrammatic presentation of a plasmid map for cloning of



145

	YIR010W into C-terminal GFP fusion vector pUG23.	150
6.6	Diagrammatic presentation of a plasmid map for cloning of	
	YIR010W into N-terminal GFP fusion vector pUG34.	151
6.7	Restriction enzyme digestion verification of YIR010W	153
	cloning.	
6.8	PCR verification of YIR010W cloning.	154
6.9	Forward sequencing result of pUG23 (YIR010W-GFP)	
	showing correct in-frame insertion.	156
6.10	Reverse sequencing result of pUG23 (YIR010W-GFP)	
	showing correct in-frame insertion.	157
6.11	SDS-PAGE result of the total protein extract (using	
	sonication) of yeast samples containing GFP pUG23/ pUG34	
	(E) and GFP vectors with YIR010W fusions (C and N).	160
6.12	Western Blot result of the total protein extract (using	
	sonication) of yeast samples containing GFP pUG23/ pUG34	
	(E) and GFP vectors with <i>YIR010W</i> fusions (C and N).	162
6.13	Comparison of yeast cells expressing GFP in vivo with non-	
	expressing cells.	163
6.14	In vivo YIR010W-GFP fusion protein expressions by pUG23	
	(YIR010W-EGFP3) and pUG34 (EGFP3-YIR010W).	165
6.15	Graph showing the (a) OD_{600} reading and (b) viability count	
	of yeast cells with different copies of YIR010W gene over	
	time at 0mM of Methionine.	167
6.16 a	(a) Graph showing the OD_{600} reading of wildtype yeast cells	
	with GFP pUG23/ pUG34 [2 copies of YIR010W gene] over	



time at different concentrations of Methionine. (b) Graph showing the OD₆₀₀ reading of wildtype yeast cells with pUG34 (N-Fus GFP-YIR010W) [3 copies of YIR010W] over time at different concentrations of Methionine.

- 6.16b (c) Graph showing the OD₆₀₀ reading of yeast cells with different copies of YIR010W gene over time at 1 mM of Methionine.
- 6.17 Comparison of OD₆₀₀ readings of wildtype carrying GFP pUG23/ pUG34 against wildtype with pUG34 (N-Fus GFP-YIR010W) [3 copies of YIR010W] over time at different Methionine concentrations.
- 7.1 Direct comparison of microscopic appearance of *Mat a* wild type (panel A) and *YOR228C* disruptant (panel B) of the same mating type.
- 7.2 Direct comparison of microscopic appearance of *Mat* α wild type (panel A) and *YOR228C* disruptant (panel B) of the same mating type.
- 7.3 Size anomaly in certain cells (long arrows) in YOR228C deletants of *Mat a* (panel A) and *Mat alpha* (panel B) that were bigger than the others (arrow heads) in the same population.
- 7.4 Direct comparison of microscopic appearance of diploid wild type (panel A) and YOR228C single or double disruptants (panel B or C respectively), which shows inconsistence in cell size in the population.



169

170

172

186

187

188