



**UNIVERSITI PUTRA MALAYSIA**

**SHELF LIFE AND QUALITY ATTRIBUTES OF FRESH BEEF  
INFUSED WITH ORGANIC ACIDS**

**MOHAMED ABD ELGADIR MOHAMED.**

**FSTM 2005 4**

**SHELF LIFE AND QUALITY ATTRIBUTES OF FRESH BEEF INFUSED  
WITH ORGANIC ACIDS**

**By**

**MOHAMED ABD ELGADIR MOHAMED**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia  
in Fulfillment of the Requirement for the Degree of Master of Science**

**March 2005**



***To my beloved parents Abd Elgadir and Fatima, who love me a lot,  
to my wife Nedal the candle of my life,  
to my bothers and sisters who bring the happiness to me***



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfillment of the requirement of the degree of Master of Science

**SHELF LIFE AND QUALITY ATTRIBUTES OF FRESH BEEF INFUSED WITH  
ORGANIC ACIDS**

**By**

**MOHAMED ABD ELGADIR MOHAMED**

**March 2005**

**Chairman : Professor Jamilah Bakar, PhD**

**Faculty : Food Science and Technology**

Fresh beef is a highly perishable food. It has a shelf life of one day at ambient temperature and a few days at refrigerated temperature. This study was conducted with the objective of extending the shelf life of fresh beef by infusing organic acids such as citric, tartaric, acetic and lactic acids, and combination of the organic acids with sodium chloride. Fresh beef (*longissimus dorsi*) purchased from the local market were sliced and were infused with citric, acetic, lactic and tartaric acids in concentration of 0.5%, 0.75% and 1%, and combination of 1.00% of citric and acetic acids with and without sodium chloride by placing samples in vacuum desiccators and pulling the vacuum to 29.5 in. Hg. for 20 min.,. All samples were packed in vacuum packs (22 (L) x 18 (w) cm. and stored



at 5°C for 28 days. pH,  $A_w$ , Total Plate Count (TPC), Thiobarbaturic acid values, Hunter colour values, instrumental texture, proximate composition were determined. The pH values of treated samples dropped from the initial pH of 5.30 (untreated) to 4.20 - 4.47 and upon storage, the pH values of all samples increased gradually. The TPC values were lower than  $10^7$  CFU/g on day 16, 20 and 28 in samples treated with 0.5%, 0.75% and 1.00% acids, respectively. The proximate composition of treated samples was affected by infusion process. The instrumental texture of fresh beef was harder upon treatment. The maximum shelf life of treated beef was 12 - 24 days for samples treated with 0.5% of all acids, 16 - 24 days for samples treated with 0.75% and 20 - 28 for samples treated with 1.00%. Citric acid in concentration of 1.00% gave the best effect which was followed by acetic acid.

In citric and acetic acids and citric and acetic acids with sodium chloride combinations, the later in the ratio of 2:1 was more effective in decreasing the initial pH of fresh beef immediately after infusion. Increasing concentration of NaCl in the infusion solution resulted in the smaller decrease in the pH values. The TPC value was observed in samples treated with 2:1 citric: acetic acids. The growth of *S.aureus* and *E.coli* O157:H7 were significantly ( $P<0.05$ ) decreased by 0.85  $\log_{10}$  and 0.73  $\log_{10}$ , respectively. The initial thiobarbituric acid value in untreated fresh beef was 0.735 mg MDA/kg which significantly ( $p< 0.05$ ) decreased



in all treated samples. At the end of storage study, lowest TBA values were obtained in samples treated with 2:1 citric and acetic. The increase in the addition of NaCl caused a parallel increased in TBA values. For colour, Hunter 'b' and 'L' values increased with storage time while 'a' decreased significantly ( $P<0.05$ ). The processed beef burger during chilled storage had a storage life of 8 days.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

## **SHELF LIFE AND QUALITY ATTRIBUTES OF FRESH BEEF INFUSED WITH ORGANIC ACIDS**

Oleh

**MOHAMED ABD ELGADIR MOHAMED**

**Mac 2005**

**Penyelia : Profesor Jamilah Bakar, PhD**

**Fakulti : Sains dan Teknologi Makanan**

Daging lembu segar adalah makanan yang sangat mudah rosak. Ia mempunyai jangka hayat selama sehari pada suhu persekitaran dan beberapa hari pada suhu dingin. Dengan itu, Kajian ini dijalankan dengan objektif untuk memanjangkan jangka hayat daging lembu segar dengan memasukkan asid organik seperti asid sitrik, tartarik, asetik, laktik, dan kombinasi asid – asid organik dengan sodium klorida (NaCl). Daging lembu segar pada bahagian (*longissimus dorsi*) telah dibeli dari pasar tempatan, dipotong dan dimasukkan dengan asid organik (sitrik, asetik, laktik dan tartaric) pada kepekatan 0.5%, 0.75% dan 1.00%, dan kombinasi 1.00% asid sitrik dan asetik dengan (1 hingga 3% NaCl dan tanpa NaCl. Ini dilakukan dengan meletakkan sampel – sampel ke dalam balang pengering vakum dan mengeluarkan gas 29.5 in. Hg. Selama



20 minit. Nilai pH sampel telah menurun dari pH asal 5.30 (tanpa rawatan) kepada 4.20 – 4.47 selepas rawatan; semasa penstoran, nilai pH pagi semua sampel yang meningkat secara perlahan – lahan. Nilai TPC bagi sampel – sampel yang telah dirawat dengan 0.5%, 0.75% dan 1.00% asid adalah kurang dari padah  $10^7$  CFU/g pada hari ke – 16, 20 dan 28. masing – masing telah menjadi komposisi terdekat bagi sampel telah dipengaruhi oleh rawatan memasukkan asid – asid, manakala tekstur daging lembu segar menjadi keras selepas rawatan. Jangka hayat bagi daging lembu yang telah dirawat dengan 0.5% asid adalah di antara 12 – 24 hari, sampel yang dirawat dengan 0.75% asid mempunyai jangka hayat antara 16 – 24 hari dan jangka hayat antara 20 – 28 hari bagi bagi sampel yang dirawat dengan 1.00% asid. Kemasukan asid sitrik pada 1.00% telah memberikan keputusan yang terbaik di ikuti dengan asid sitrik.

Dalam kajian kombinasi campuran asid sitrik dan asetik dengan dan tanpa NaCl telah digunakan. Campuran asid sitrik dan asetik pada nisbah 2:1 tanpa NaCl adalah lebih berkesan dalam menurunkan pH asal daging lembu segar sebaik selepas dirawat. Peningkatan kepekatan NaCl dalam larutan rawatan menyebabkan sedikit penurunan pada nilai pH. Nilai TPC yang terendah dapat diperhatikan dalam sampel yang telah dirawat dengan 2:1 asid sitrik: asetik. Pertumbuhan *S. aureus* dan *E. coli* O157:H7 telah menurun secara bermakna ( $p < 0.05$ ) sebanyak  $0.85 \log_{10}$  dan  $0.73 \log_{10}$  bagi sampel – sampel yang telah dirawat. Nilai TBA asal bagi daging lembu segar tanpa rawatan adalah 0.735





mg MDA/ kg dan telah menurun secara bermakna ( $p < 0.05$ ) dalam semua sampel yang telah dirawat. Pada penghujung kajian penstoran, nilai TBA yang terendah dapat diperolehi dalam sampel yang telah dirawat dengan 2:1 sitrik dan asetik. Peningkatan dalam nilai TBA. Bagi warna, nilai Hunter 'b' dan 'L' meningkat dengan masa perstoran manakala nilai Hunter 'a' menurun secara bermakna ( $p < 0.05$ ). Burger daging lembu mempunyai jangka hayat penstoran selama 8 hari pada suhu 4° C.



## ACKNOWLEDGEMENT

First of all, praises and thanks for Almighty Allah who has given me all the best during all stages of my study. My acknowledgement goes to the following:

My supervisor Prof. Dr. Jamilah Bakar, really dear Prof., no words will be able to express my heartfelt gratitude for your kindly strong constant guidance, advice, invaluable discussion and new ideas through the period of this study.

My supervisory committee Prof. Dr. Yaakob Che Man and Prof. Dr. Russly Abdul Rahman for their advises and stimulating discussion especially during the presentation of my work.

All technicians in our faculty, En. Azman, Pn. Jamaliah; En. Abdul Halim; En. Zolkifli; Pn. Noorfaizan and Pn. Rozmawati (members of biochemistry lab.), for the continuously assistance and friendly relationship. My sincere gratitude also goes to En. Mohamed Soib in the engineering lab. for his unique effort with me in texture measurement and brotherhood relationship. Deepest gratitude to En. Zolkifli and Pn. Jamilah (Microbiology lab.) for their kind assistance. I wish to extend my appreciation to En. Razali Othman and Pn. Siti Shahrul Bariah (processing lab.) for their helpfulness during the processing of burger.



My wife for her love, encouragement, assistant and patience through the length of my study.



I certify that an Examination Committee met on 31<sup>st</sup> March 2005 to conduct the final examination of Mohamed Abd Elgadir Mohamed on his Master of Science thesis entitled "Shelf Life and Quality Attributes of Fresh Beef Infused with Organic Acids" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Salmah Yusof, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Mohd Yazid Abdul Manap, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Internal Examiner)

**Roselina Karim, PhD**

Lecturer  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Internal Examiner)

**Abdul Salam Babji, PhD**

Professor  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(External Examiner)



---

**GULAM RUSUL RAHMAT ALI, PhD**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: **21 JUL 2005**



This thesis submitted to the senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Jamilah Bakar, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Chairman)

**Yaakob Che Man, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)

**Russly Abdul Rahman, PhD**

Professor  
Faculty of Food Science and Technology  
Universiti Putra Malaysia  
(Member)



---

**AINI IDRIS, PhD**

Professor/ Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 11 AUG 2005



## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



---

**MOHAMED ABD ELGADIR**

Date: **18 JULY** 2005

## TABLE OF CONTENT

	DEDECATION	ii
	ABSTRACT	iii
	ABSTRAK	vi
	ACKNOWLEDGEMENT	ix
	APPROVAL SHEETS	xi
	DECLARATION	xiii
	TABLE OF CONTENT	xiv
	LIST OF TABLES	xvii
	LIST OF FIGURES	xix
	LIST OF PLATES	xxi
	LIST OF ABBREVIATIONS	xxii
<b>CHAPTER</b>		
<b>I</b>	<b>INTRODUCTION</b>	<b>1</b>
<b>II</b>	<b>LITERATURE REVIEW</b>	<b>4</b>
	Nutritional and chemical composition of fresh beef	4
	Physico - chemical properties of beef	5
	pH and Water activity	5
	Colour	5
	Texture	8
	Microbial characteristics of fresh beef	10
	Spoilage microorganisms	10
	Gram negative (oxidase positive) rod – shape	
	Bacteria	12
	Coliform bacteria	13
	Gram positive spore – forming bacteria	14
	Lactic acid bacteria.	14
	Pathogenic microorganisms	15
	Packaging and storage of chilled beef products	16
	Spoilage of fresh beef	19
	Microbial	19
	Lipid oxidation in beef and its products	21
	Change in beef flavour during storage	25
	The shelf life of fresh beef	25
	Organic acids as preservatives to prolong the shelf life of fresh beef	27
	Citric acid	28
	Acetic acid	29
	Lactic acid	30
	Tartaric acid	30



	Regulatory status of food grade organic acids	31
	Effect of sodium chloride on fresh beef preservation	32
	Beef burger processing	33
	Storage stability of beef burger	34
	Quality of beef burger	35
	Sensory attributes of beef products	36
<b>III</b>	<b>EFFECT OF ORGANIC ACIDS INFUSION ON SHELF LIFE EXTENSION AND PHYSICO - CHEMICAL PROPERTIES OF FRESH BEEF</b>	<b>38</b>
	Introduction	38
	Materials and methods	41
	Material	41
	Methods	41
	Preparation of organic acid	41
	Preparation of peptone water	41
	Preparation plates of Plate Count Agar	42
	Infusion processing	42
	Storage studies	42
	Analysis	43
	PH	43
	Water activity	43
	Total Plate Count	44
	Colour	44
	Texture	45
	Proximate composition	46
	Thiobarbituric acid value	46
	Statistical Analysis	47
	Results and discussions	47
	PH	47
	Water activity	48
	Total Plate Count	51
	Colour	52
	Texture	58
	Proximate composition	60
	Thiobarbituric acid value	62
<b>IV</b>	<b>SYNERGISTIC EFFECTS OF ORGANIC ACIDS AND SODIUM CHLORIDE COMBINATION ON PHYSICO- CHEMICAL AND MICROBIAL QUALITY OF FRESH BEEF</b>	<b>64</b>
	Introduction	64
	Material and methods	65
	Materials	65
	Analytical methods	65
	pH measurement	65





	Total Plate Count assessment	66
	<i>Staphylococcus aureus</i> numeration	66
	<i>E.coli</i> O157:H7 assessment	66
	TBA measurement	67
	Colour measurement	67
	Statistical analysis	68
	Results and discussions	68
	pH measurement	68
	Total Plate Count assessment	71
	<i>S. aureus</i> and <i>E.coli</i> O157:H7assessment	72
	TBA measurement	76
	Colour measurement	78
	Conclusion	80
<b>V</b>	<b>QUALITY ATTRIBUTES OF BURGER PROCESSED FROM CITRIC ACID TREATED BEEF</b>	
	Introduction	81
	Preparation of beef burger	82
	Storage study	83
	pH measurement of the burger	83
	Water activity measurement of the burger	83
	Total Plate Count analysis of the burger	84
	TBA measurement of the burger	84
	Sensory analysis	84
	Statistical analysis	85
	Results and discussions	85
	pH measurement of the burger	85
	Water activity measurement of the burger	86
	Total Plate Count analysis of the burger	86
	TBA measurement of the burger	88
	Sensory analysis	89
	Conclusion	91
<b>VI</b>	<b>Conclusion and Recommendations</b>	93
	Bibliography	95
	APPENDICES A	111
	APPENDICES B	113
	APPENDICES C	116
	APPENDICES D	119
	APPENDICES E	122
	APPENDICES F	125
	BIODATA OF THE AUTHOR	128



## LIST OF TABLES

Table		Page
1	pH of control and acids treated samples during storage at 5°C for 28 days	49
2	Water activity of control and acids treated samples during storage at 5°C for 28 days	50
3	TPC of control and acids treated samples during storage at 5°C for 28 days	54
4	Hunter 'L' 'a' and 'b' values for control and 0.5% acids treated samples during storage at 5°C for 28 days	55
5	Hunter 'L' 'a' and 'b' values for control and 0.75% acids treated samples during storage at 5°C for 28 days	56
6	Hunter 'L' 'a' and 'b' values for control and 1.00% acids treated samples during storage at 5°C for 28 days	57
7	Texture of control and 0.5% acids treated samples during storage at 5°C for 28 days	59
8	Proximate composition of control acids treated samples during storage at 5°C for 28 days	61
9	TBA values of control acids treated samples during storage at 5°C for 28 days	63
10	pH of control and acids combination treated samples during storage at 5°C for 28 days	70
11	Total Plate counts (TPC) of control and acids combination treated samples during storage at 5°C for 28 days	73
12	<i>S. aureus</i> of control and acids combination treated samples during storage at 5°C for 28 days	74



13	<i>E. coli</i> O157: H7 of control and acids combination treated samples during storage at 5°C for 28 days	75
14	TBA values of control and acids combination treated samples during storage at 5°C for 28 days	77
15	Hunter colour 'a' values of control treated samples during storage at 5°C for 28 days	79
16	Values of analysis of burger during chilled storage (4° C) for 8 days	88
17	Sensory scores for burger chilled stored at 4°C for 8 days	91



## LIST OF FIGURES

Figure		Page
1	Reaction of lipase with fatty acids in meat.	23
2	pH of control and 0.5% acids treated samples of fresh beef during storage at 5°C for 28 days	113
3	pH of control and 0.75% acids treated samples of fresh beef during storage at 5°C for 28 days	114
4	pH of control and 1.00% acids treated samples of fresh beef during storage at 5°C for 28 days	115
5	Water activity of fresh beef (control) and 0.5% acids treated samples during storage at 5°C for 28 days	116
6	Water activity of fresh beef (control) and 0.75% acids treated samples during storage at 5°C for 28 days	117
7	Water activity of fresh beef (control) and 1.00% acids treated samples during storage at 5°C for 28 days	118
8	TPC of fresh beef (control) and 0.5% acids treated sample during storage at 5°C for 28 days	119
9	TPC of fresh beef (control) and 0.75% acids treated sample during storage at 5°C for 28 days	120
10	TPC of fresh beef (control) and 1.00% acids treated sample during storage at 5°C for 28 days	121
11	Hunter 'a' values for fresh beef (control) and 0.5% acids treated sample during storage at 5°C for 28 days	122
12	Hunter 'a' values for fresh beef (control) and 0.75% acids treated sample during storage at 5°C for 28 days	123
13	Hunter 'a' values for fresh beef (control) and 1.00% acids treated sample during storage at 5°C for 28 days	124



14	Texture of fresh beef (control) and 0.5% acids treated sample during storage at 5°C for 28 days	125
15	Texture of fresh beef (control) and 0.75% acids treated sample during storage at 5°C for 28 days	126
16	Texture of fresh beef (control) and 1.00% acids treated sample during storage at 5°C for 28 days	127



## LIST OF ABBREVIATIONS

DFD	Dark Firm Dry meat
PSE	Pale Soft Exudative meat
Mb	myoglobin
MbO <sub>2</sub>	oximyoglobin
MMb	met myoglobin
A <sub>w</sub>	water activity
ERH	Equilibrium Relative Humidity
ISO	International Organization for Standardization
EHEC	Enterohaemorrhagic Escherichia Coli
MAP	modified atmosphere packaging
VSP	vacuum skin packaging
AP	active packaging
VP	vacuum packaging
PE	polyethylene
PP	polypropylene
PS	polystyrene
ABS	acrylonitrile - butadiene – styrene
PC	poly carbonate
EVA	ethylene vinyl acetate
pK <sub>a</sub>	Organic acid dissociation constant
FSIS	Food Safety and Inspection Service
USDA	United States Department of Agriculture



CFU/g	Colony forming unit per gram
CFU/ cm <sup>2</sup>	Colony forming unit per square centimeter
SPI	Soy protein isolate
TPC	Total Plate Count
TBA	Thiobarbituric acid



## CHAPTER I

### INTRODUCTION

Fresh beef is rich in vitamins and minerals and provides an important source of high quality protein. It has a short shelf life of one day or less at ambient temperature (15 – 30°C) and a few days at refrigerated temperature (0 – 10°C) due to microbial spoilage of both pathogenic and non - pathogenic (Dickson and Anderson, 1992) and/or lipid oxidation (Ahn et al., 1992; Shahidi; 1994b; and Morrissey et al., 1998). The maximum shelf life of fresh beef depends on several factors such as pH, water activity, microbial growth and temperature (Farber, 1991).

Many studies have been reported using organic acids to reduce organic spoilage microorganisms in beef such as spraying (Siragusa and Dickson, 1992 and Doores, 1993b), washing (Dorsa et al; 1997; and Cutter and Siragusa, 1994), and dipping (Miller et al., 1993). Food grade additives such as organic acids (citric, acetic, lactic and tartaric) and sodium chloride are reported to prolong the storage life of fresh beef. Organic acids are used as anti-microbial preservatives or acidulant in beef and its products due to their high solubility and low toxicity (Cassens, 1994). Citric acid was investigated for its inhibition effect on bacteria, yeast and molds and was shown to be more inhibitive and effective compared to lactic and citric acids (Sorrells,





1989). Anderson and Marshall (1990) concluded that a 3% concentration of acetic acid was most effective in sanitizing beef muscle inoculated with *Escherichia coli* and *Salmonella*. Spray washing of pork loin with 2% acetic acid before vacuum packaging followed by storage for 28 days at 4°C significantly lowered the aerobic, anaerobic and lactic acid bacteria counts (Anonymous, 1990). Lactic acid sprays have been effective in limiting microbial growth on meat carcasses under a variety of storage conditions at 4 – 55°C (Cutter and Siragusa, 1994). Tartaric acid acts synergistically with antioxidants to prevent rancidity and for miscellaneous and general purpose usage in accordance with good manufacturing practice (Doores, 1993a). However, organic acids have a negative effect on beef colour since they reduce the oxymyoglobin pigment (red colour) to metmyoglobin (brown undesirable colour). These studies were conducted to prolong the storage life of fresh beef.

Processing of beef products causes changes in texture and flavor and also adds variety to the diet. Processing also provides scope to mix the less desirable parts of the carcass with lean meat and in addition is a means of extending meat supplies by including other foodstuffs such as cereal in the product. Beef pattie is one of the examples. The quality attributes of pattie depends largely on the type and quality of lean meat and ingredients used. Hence, the objectives of this study are (1) to determine the effects of the organic acids infusion on shelf life extension and physico - chemical