



UNIVERSITI PUTRA MALAYSIA

**EFFECTS OF NEWCASTLE DISEASE VIRUS STRAINS AF 2240 AND
V4-UPM ON CYTOLYSIS AND APOPTOSIS OF LEUKEMIA CELL LINES**

AIED MOHAMMED AL-ABSI

FBSB 2008 20

**EFFECTS OF NEWCASTLE DISEASE VIRUS STRAINS AF 2240 AND V4-UPM
ON CYTOLYSIS AND APOPTOSIS OF LEUKEMIA CELL LINES**

By

AIED MOHAMMED AL-ABSI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

September 2008



Specially dedicated to

My beloved parents

My loving wife

My daughters

My brothers

My sisters

My country



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the Degree of Doctor of Philosophy

**EFFECTS OF NEWCASTLE DISEASE VIRUS STRAINS AF 2240 AND V4-
UPM ON CYTOLYSIS AND APOPTOSIS OF LEUKEMIA CELL LINES**

By

AIED MOHAMMED AL-ABSI

September 2008

Chairman: Professor Abdul Manaf Ali, PhD

Faculty: Biotechnology and Biomolecular Sciences

Newcastle disease virus (NDV) is a member of the *Paramyxoviridae* that has caused severe economic losses in the poultry industry worldwide. Several strains of NDV were reported to induce cytolysis to cancerous cell lines. In this study, two NDV isolates namely, AF 2240 and V4-UPM were evaluated for their anti-leukemic properties against four leukemic cell lines - HL60 (Promyelocytic leukemia), WEHI 3B (Mouse myelomonocytic leukemia), CEMSS (Human T-lymphobalstic leukemia) and K562 (Erythromyelobalstic leukemia). The cytolytic effects of NDV strains AF 2240 and V4-UPM towards WEHI 3B, HL60, CEMSS and K562 cell lines were determined using microtetrazolium (MTT) assay. The cytolytic dose - fifty percent (CD_{50}) for WEHI 3B, HL60 and CEMSS treated with AF 2240 strain were 2, 25, 16 HAU, respectively, while the CD_{50} for WEHI 3B, HL60 and CEMSS treated with V4-UPM were 8, 110 and 64 HAU, respectively.

Comparatively, both NDV strains showed very low cytolytic activity against K562 and non-leukemic cell lines namely, 3T3 (mouse fibroblasts), mouse lymphocytes



and human peripheral lymphocytes. Further studies were done to observe the morphological changes in the WEHI 3B treated cells using light, transmission and scanning electron microscopes. The apoptosis and necrosis were examined under fluorescence microscope, where the cells were stained with acridine orange (AO) and propidium iodide (PI). The treated cells with NDV strains AF 2240 and V4-UPM showed apoptotic features such as cell shrinkage, cell blebbing, and formation of apoptotic bodies compared to the control cells that did not express any features for apoptosis and necrosis. The early apoptosis was also observed under fluorescence microscope, where the cells were stained with Annexin V and PI.

The virus effect on cell proliferation was determined by MTT assay and BrdU techniques. Furthermore, at molecular level, both NDV strains caused internucleosomal DNA cleavage producing a multiple of 180-200 bp fragments, that were visible as a ladder on the agarose gel. Early apoptosis was also observed using Annexin V flow cytometry method. The percentage of apoptosis for WEHI 3B cells treated with NDV strains AF 2240 and V4-UPM had increased with time. Cell cycle and apoptosis were also determined using flow cytometry PI method. Both NDV strains were not able to arrest WEHI 3B at specific cell cycle phases using flow cytometry PI method.

In this study anti-leukemic activity of both NDV strains was evaluated on BALB/c mice induced leukemia with WEHI 3B cells. A day later they were treated with NDV strains AF 2240 and V4-UPM and arabinocytocine, a commercial drug, as positive control. The mice groups treated with arabinocytocine, NDV strains AF 2240 and V4-UPM showed significant killings ($p < 0.05$) of leukemic cells compared to the



mice group without any treatment. The total white blood cell and percentage of blasts cell in the blood, bone marrow and spleen smears were significantly low ($p < 0.05$) in mice treated with the arabinocytocine, NDV strains AF 2240 and V4-UPM compared to the mice group without treatment, that showed high number of leukemia. Spleen and liver weights were significantly low ($p < 0.05$) in the mice groups treated with arabinocytocine, NDV strains AF 2240 and V4-UPM compared to the mice group without treatment, that showed significant ($p < 0.05$) splenomegaly and hepatomegaly.

Histopathological studies carried out had confirmed haematological results. From the results obtained, the mice groups treated with both NDV stains AF 2240 and V4-UPM showed similar results as arabinocytocine, which is a commercial drug for leukemia. Immunoperoxidase staining, haemagglutination test and real time PCR were carried out to detect NDV in mice organs after treatment with NDV. The results showed no NDV particles were detected in the organs. This study showed that NDV strains AF 2240 and V4-UPM had caused cytolytic effects against WEHI 3B leukemia cell line *in vitro* and *in vivo*.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah.

**KESAN VIRUS PENYAKIT NEWCASTLE STRAIN AF 2240 DAN V4-UPM
TERHADAP SITOTOSIK DAN APOPTOSIS PADA SEL LEUKEMIA**

Oleh

AIED MOHAMMED AL-ABSI

September 2008

Pengerusi: Profesor Abdul Manaf Ali, PhD

Fakulti: Bioteknologi dan Sains Biomolekul

Virus penyakit Newcastle (NDV) merupakan ahli *Paramyxoviridae* yang menyebabkan banyak kes kemelesetan ekonomi di industri penternakan di seluruh dunia. Beberapa strain NDV telah dilaporkan dapat mencetuskan sitolisis terhadap sel barah. Dalam pengajian ini, dua strain NDV yang diasingkan adalah AF2240 dan V4-UPM telah diujikan kesan anti-leukemic terhadap empat jenis sel leukemia yang dinamakan HL60 (Promyelocytic leukaemia), WEHI 3B (myelomonocytic leukaemia tikus), CEMSS (T-lymphoblastic leukaemia manusia) dan K562 (Erythromyeloblastic leukaemia). Kesan sitotosik oleh NDV strain AF2240 dan V4-UPM terhadap sel WEHI 3B, HL60, CEMSS dan K562 telah ditentukan melalui analisa methyl thiazolyl tetrazolium (MTT). Kesan sitotosik lima puluh peratus (CD_{50}) untuk WEHI 3B, HL60 dan CEMSS yang dirawat dengan AF2240 adalah 2, 16, 35 HA unit masing-masing, manakala CD_{50} untuk sel WEHI 3B, HL60 dan CEMSS yang dirawat dengan V4-UPM adalah 8, 32 dan 64 HA unit masing-masing.



Secara perbandingan, kedua-dua strain NDV menunjukkan kesan sitotoksik yang sangat rendah terhadap K562 dan sel bukan leukemia yang dinamakan, 3T3 (fibroblasts tikus), limfosit tikus dan limfosit pinggir manusia. Kajian selanjutnya telah dijalankan untuk memperhatikan perubahan morfologi dalam sel WEHI 3B yang dirawat dengan menggunakan cahaya, pancaran dan imbasan elektron mikroskop. Sifat morfologi, apoptosis dan nekrosis juga diperiksa dengan mikroskop pendarfluor di mana sel diwarnai dengan akridin oren (AO) dan propidium iodida (PI). Sel yang dirawat dengan NDV strain AF 2240 dan V4-UPM menunjukkan sifat apototik seperti kecutan sel, blebbing sel, pembentukan jasad apototik dan sel nekrotik berbanding dengan sel pengawal negatif tidak menunjukkan sebarang sifat apoptosis dan nekrosis. Apoptosis awal juga diperhatikan melalui mikroskop pendarfluor di mana sel telah diwarnakan dengan Annexin V dan PI.

Kesan virus terhadap pembelahan sel telah dikenal pasti dengan teknik MTT dan BrdU. Di samping itu, dari segi molekul, kedua-dua strain virus ini dapat menyebabkan pemotongan DNA internukleus untuk menghasilkan pelbagai serpihan 180-200 bp yang dapat dilihat sebagai tangga pada gel agarosa. Awal apoptosis juga diperhatikan melalui Annexin V kaedah aliran sitometri. Peratus apoptosis untuk sel WEHI 3B yang dirawat dengan NDV strain AF2240 dan V4-UPM telah meningkat dari semasa ke semasa. Kitar sel dan apoptosis juga ditentukan melalui kaedah aliran sitometri PI. Kedua-dua strain NDV tidak dapat menahankan WEHI 3B pada fasa kitar sel yang tertentu dengan menggunakan kaedah aliran sitometri PI.



Dalam kajian ini aktiviti anti-leukemik pada kedua-dua strain NDV telah dikaji dengan tikus BALB/c yang telah dicetuskan leukemik dengan sel WEHI 3B. Pada hari keesokan mereka telah dirawat dengan NDV strains AF2240 dan V4-UPM dan arabinositosin sebagai ubat kormesial yang dijadikan pengawal positif. Kumpulan tikus yang dirawat dengan arabinositosin, NDV strain AF 2240 dan V4-UPM, masing-masing menunjukkan kesan pembunuhan yang ketara ($p < 0.05$) terhadap sel leukemic berbanding dengan tikus untuk kumpulan tanpa sebarang rawatan. Jumlah sel darah putih dan peratus sel blast di sapuan darah, sum-sum tulang dan limpa adalah rendah ($p < 0.05$) di tikus yang dirawat dengan NDV strain AF 2240 dan V4-UPM, masing-masing, berbanding dengan kumpulan tikus tanpa rawatan yang menunjukkan nombor leukimia yang tinggi. Keberatan limpa dan hati adalah rendah secara ketara ($p < 0.05$) di kumpulan tikus yang dirawat dengan arabinocytocine, NDV strain AF 2240 dan V4-UPM, masing-masing, berbanding dengan kumpulan tikus tanpa sebarang rawatan yang menunjukkan splenomegaly dan hepatomegaly yang ketara ($p < 0.05$).

Kajian histopatologikal juga dijalankan dan mengesahkan keputusan hematologikal. Daripada keputusan yang diperolehi, kumpulan tikus yang dirawat dengan kedua-dua NDV strain AF 2240 dan V4-UPM menunjukkan keputusan yang seiras dengan arabinocytocine yang merupakan ubat kormesial untuk leukemia. Pewarna immunoperoxidase, analisa hemagglutinin dan real time PCR telah dijalankan untuk mengesan NDV di organ tikus setelah dirawat dengan NDV. Keputusan telah menunjukkan tiada zarah NDV dikesan di organ tikus. Kajian ini menunjukkan NDV



strain AF 2240 dan V4-UPM menyebabkan kesan sitolitik terhadap sel leukemia WEHI 3B secara *in vitro* dan *in vivo*.



ACKNOWLEDGEMENTS

In the name of Allah, the most gracious, the most merciful

I own foremost my profound gratitude to Allah, the Almighty for providing me the strength and diligence to complete this dissertation despite several obstacles encountered throughout the progress of this study which at times seemed insurmountable.

I would like to express my deepest and heartiest thanks and indebtedness to my supervisor, Prof. Dr. Abdul Manaf Ali, for his guidance and suggestions through the course of this study in the midst of his heavy responsibilities. The same appreciation goes to my co-supervisors; Associate Prof Dr. Abdul Rahman Omar, Prof. Dr. Aini Ideris and Prof. Dr. Mohammed Hair Bejo for their patience and guidance throughout the study.

I would like to thank Dr. Norjahan Mohd. Alitheen for her valuable guidance as well as for helping me to do flow cytometry readings during the practical part of my study and I would like to thank all staff at the Microscope Unit, Institute of Bioscience, Universiti Putra Malaysia.

Last but not least, I would like to express my deepest gratitude to my beloved parents, wife, sisters and brothers for their endless encouragement, patience and sacrifices which had helped me throughout my student life.



I certify that a Examination Committee has met on 22 September 2008 to conduct the final examination of Aied Mohammed Al-Absi on his Doctor of Philosophy thesis entitled “Cytolytic Effects and Apoptosis Induction of Newcastle Disease Virus Strains AF 2240 and V4-UPM on Leukemia Cell Lines In vitro and In vivo” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the Doctor of Philosophy. Members of the examination committee were as following:

Raha Abdul Rahim, PhD

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Fauziah Othman, PhD

Professor

Faculty of Medicine and Health Science

Universiti Putra Malaysia

(Internal Examiner)

Tan Weng Siang, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

Mohd Nizam Hj. Isa, PhD

Professor

Malaysian University of Science and Technology

(External Examiner)

Hasanah Mohd, Ph.D.

Professor/Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the supervisory committee were as follows:

Abdul Manaf Ali, PhD

Professor
Faculty of Biotechnology and Bimolecular Sciences
Universiti Putra Malaysia
(Chairman)

Aini Ideris, PhD

Professor
Faculty of Veterinary Medicine,
Universiti Putra Malaysia.
(Member)

Abdul Rahman Omar, PhD

Lecturer
Institute of Bioscience
Universiti Putra Malaysia
(Member)

Mohd Hair Bejo, PhD

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia.

Date: 19 December 2008



DECLARATION

I declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously and is not concurrently submitted for any other degree at UPM or at any other institutions.

AIED MOHAMMED AL-ABSI

Date: 26 February 2009



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	x
APPROVAL	xii
DECLARATION	xiii
LIST OF TABLES	xx
LIST OF FIGURES	xxiii
LIST OF PLATES	xxvii
LIST OF ABBREVIATIONS	xxix
CHAPTERS	
1 INTRODUCTION	1
2 LITERATURE REVIEW	9
2.1 Cancer	9
2.1.1 Cancer Genes	10
2.1.2 Classification of Tumors	11
2.2 Leukaemia	11
2.2.1 History of Leukaemia	12
2.2.2 Incidence of Leukaemia	13
2.2.3 Classification of Leukaemia	14
2.2.4 Molecular Pathogenicity of Leukemia	15
2.2.5 Diagnosis of Leukaemia	21
2.2.6 Treatment of Leukaemia	29
2.3 Newcastle Disease	32
2.3.1 History	32
2.3.2 Classifications	33
2.3.3 Structures	33
2.3.4 Pathotyping	34
2.3.5 Molecular Basis of NDV	35
2.3.6 NDV strain AF 2240	36
2.3.7 NDV strain V4-UPM	37
2.4 Cancer Therapy	38
2.4.1 Cancer Gene Therapy	38
2.4.2 Cancer Chemotherapy	38
2.4.3 Cancer Virotherapy	39
2.4.4 Virotherapy versus Chemotherapy as Anticancer Agent	41
2.4.5 Virotherapy Strategies of Cancer Treatment	42
2.5 Mechanisms of Oncolytic Viruses to Destruct Tumor Cells	42



2.6 Replication of Oncolytic Virus in Tumor Cells	44
2.7 History of Oncolytic Virus as Anticancer	45
2.8 Newcastle Disease Virus as Cancer Therapy	47
2.8.1 Regime of NDV Administration	49
2.8.2 In vitro Studies of NDV	50
2.8.3 In Vivo Studies of NDV	52
2.8.4 Oncolytic Activity of Local NDV strains	54
2.8.5 Human Clinical Studies of NDV as Anticancer	55
2.9 Cell Death	62
2.9.1 Apoptosis	63
2.9.2 Necrosis	66
2.10 Apoptosis Induction by Oncolytic NDV	68
CYTOLYTIC EFFECT OF NDV STRAINS AF2240 AND V4-UPM ON LEUKAEMIA CELL LINES	72
3.1 INTRODUCTION	72
3.2 MATERIAL AND METHOD	74
3.2.1 Preparation of virus stock	74
3.2.2 Cell Lines Maintenance	78
3.2.3 Preparation of Spleen and Thymus Cell Suspensions	79
3.2.4 Isolation of Peripheral Blood Lymphocyte	79
3.2.5 Cell Counting	81
3.2.6 Cytotoxicity Assay (MTT)	81
3.2.7 Proliferation Assay	84
3.2.8 Statistical Analysis	87
3.3 RESULTS	87
3.3.1 Titration of The Virus	87
3.3.2 Cytotoxicity of NDV Strains AF 2240 and V4-UPM	88
3.3.3 Cell Proliferation Assay	103
3.4 DISCUSSION	110
3.4.1 Titration of The Virus	110
3.4.2 Cytotoxicity of Newcastle Disease Virus Strains AF 2240 and V4-UPM	110
3.4.3 Cell Proliferation Assay	113



CHARACTERIZATION AND MORPHOLOGICAL CHANGES OF WEHI 3B CELL LINE TREATED WITH NDV STRAINS AF 2240 AND V4-UPM	116
4.1 INTRODUCTION	116
4.2 MATERIALS AND METHODS	120
4.2.1 Morphological Assessment	120
4.2.2 DNA Extraction and Laddering	124
4.2.3 Flow cytometry	126
4.2.4 Statistical Analysis	127
4.3 RESULTS	128
4.3.1 Morphological Assessment of Apoptosis	128
4.3.2 DNA Fragmentation Assay	150
4.3.3 Flow cytometry	152
4.4 DISCUSSION	162
4.4.1 Morphological Assessment of Apoptosis	162
4.4.2 DNA Fragmentation Assay	169
4.4.3 Flow cytometry	171
NDV STRAINS AF 2240 AND V4-UPM AS ANTI-LEUKEMIC AGENT IN BALB/C LEUKEMIA INDUCED MICE	174
5.1 INTRODUCTION	174
5.2 MATERIALS AND METHODS	177
5.2.1 Animal Groups and Cells	177
5.2.2 Virus and Arbinocytocine Treatment	178
5.2.3 Mortality	179
5.2.4 Body Weight	179
5.2.5 Spleen and Liver Weights	179
5.2.6 Haematology Study	179
5.2.7 Histology study	182
5.2.8 Real Time PCR	184
5.2.9 Virus detection Using Haeagglutination Test (HA)	186
5.2.10 Toxicity Test	187
5.2.11 Statistical Analysis	187
5.3 RESULTS	188
5.3.1 Mortality	188
5.3.2 Mice Body Weight	189
5.3.3 Hematological Study	191
5.3.4 Spleen and Liver Weight	204



5.3.5 Histopathological Study	209
5.3.6 Real Time PCR	219
5.3.7 Virus detection Using Haeagglutination Test (HA)	221
5.3.8 Toxicity Test	222
5.4 DISCUSSION	229
5.4.1 Mortality	229
5.4.2 Mice Body Weight	230
5.4.3 Hematological Study	230
5.4.4 Spleen and Liver Weight	232
5.4.5 Histopathological Study	233
5.4.6 Real Time PCR	235
5.4.7 Virus detection Using Haeagglutination Test (HA)	235
5.4.8 Toxicity Test	236
6 CONCLUSION	239
7 REFERENCES	246
8 APPENDICES	270
9 BIODATA OF THE STUDENT	289



LIST OF TABLES

Table		Page
2.1	Morphological and biochemical differences between necrosis and apoptosis.	67
3.1	HAU titer of NDV strains AF 2240 and V4-UPM	87
3.2	Cytotoxic effect dose (CD ₅₀) of Newcastle disease virus strains (AF 2240 and V4-UPM) against WEHI 3B cell after 6, 24, 48 and 72 hrs	91
3.3	Cytotoxic effects dose (CD ₅₀) of Newcastle disease virus strains (AF 2240 and V4-UPM), and commercial drugs (doxorubicin and arabinocytocine) against leukemia and normal cell lines at 72 hrs of incubation	92
3.4	Percentage of cell viability of human lymphocytes, mouse spleen lymphocytes and normal mouse fibroblasts treated with 2 and 8 HAU of NDV strains AF 2240 and V4-UPM, respectively that killed 50% of WEHI 3B cell line.	92
3.5	Percentage of cell viability of human lymphocytes, mouse spleen lymphocytes and normal mouse fibroblasts treated with 25 and 110 HAU of NDV strains AF 2240 and V4-UPM, respectively that killed 50% of HL60 cell line.	93
3.6	Percentage of cell viability of human lymphocytes, mouse spleen lymphocytes and normal mouse fibroblasts treated with 16 and 64 HAU of NDV strains AF 2240 and V4-UPM, respectively that killed 50% of CEM-SS cell line.	93
4.1	Percentage of necrotic, apoptotic and viable of WEHI 3B cells in the population after treated with CD 50 of NDV AF2240 virus strain at different times.	136
4.2	Percentage of necrotic, apoptotic and viable of WEHI 3B cells in the population after treated with CD 50 of NDV V4-UPM virus strain at different times.	136
4.3	Cell cycle analysis of WEHI 3B cell population treated with NDV strain AF 2240 at CD ₅₀ value (2 HAU) after staining with propidium iodide.	156



4.4	Cell cycle analysis of WEHI 3B cell population treated with NDV strain V4-UPM at CD ₅₀ value (8 HAU) after staining with propidium iodide.	156
4.5	Effects of NDV strain AF 2240 at CD ₅₀ value (2 HAU) on WEHI 3B cells by using Annexin-V FITC staining flow cytometry.	161
4.6	Effects of NDV strain V4-UPM at CD ₅₀ value (8 HAU) on WEHI 3B cells by using Annexin-V FITC staining flow cytometry.	161
5.1	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on survival time of BALB/c mice inoculated with WEHI-3B leukemia cells.	189
5.2	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on body weight of BALB/c mice inoculated with WEHI-3B leukemia cells.	190
5.3	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the total leukocyte count in the peripheral blood of BALB/c mice inoculated with WEHI-3B leukemia cells	194
5.4	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the percentage of blasts cells in the peripheral blood of BALB/c mice inoculated with WEHI-3B leukemia cells.	195
5.5	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the percentage of blast cells in the bone marrow of BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	198
5.6	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the percentage of blast cells in the spleen of BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	202
5.7	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the spleen weight and length in the BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	206
5.8	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the liver weight in the BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	207
5.9	Ct and Tm values of SYBR Green I real time PCR detecting NDV strains AF 2240 and V4-UPM in Spleen, heart, liver and kidney of BALB/c mice.	221



5.10	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the liver function enzymes of BALB/c mice inoculated with WEHI-3B leukemia cells.	224
5.11	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the kidney function enzymes of BALB/c mice inoculated with WEHI-3B leukemia cells.	226
5.12	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the heart function enzymes of BALB/c mice inoculated with WEHI-3B leukemia cells.	228



LIST OF FIGURES

Figure		Page
2.1	Infection and killing of tumour cells by an oncolytic virus.	45
2.2	Features of the apoptotic and necrotic cell death process	65
3.1	Percentage of cell viability of WEHI 3B cells treated with NDV AF 2240 and V4-UPM virus strains after 6 hrs.	94
3.2	Percentage of cell viability of WEHI 3B treated with NDV AF 2240 virus strain after 24, 48, and 72 hrs.	95
3.3	Percentage of cell viability of WEHI 3B treated with NDV V4-UPM virus strain after 24, 48, and 72 hrs.	96
3.4	Percentage of cell viability of K 562 treated with NDV virus strains AF 2240 and V4-UPM after 72 hrs.	97
3.5	Percentage of cell viability of CEMSS cells treated with NDV AF 2240 virus strain after 72 hrs.	97
3.6	Percentage of cell viability of CEMSS cells treated with NDV V4-UPM virus strain after 72 hrs.	98
3.7	Percentage of cell viability of HL60 cells treated with NDV AF 2240 virus strain after 72 hrs.	98
3.8	Percentage of cell viability of HL60 cells treated with NDV V4-UPM virus strain after 72 hrs.	99
3.9	Percentage of cell viability of 3T3 treated NDV AF 2240 and V4-UPM virus strains after 72 hrs.	99
3.10	Percentage of cell viability of mouse spleen lymphocyte cell treated with NDV AF 2240 and V4-UPM virus strains after 72 hrs.	100
3.11	Percentage of cell viability of peripheral blood cell treated with NDV AF 2240 and V4-UPM virus strains after 72 hrs.	100
3.12	Percentage of cell viability of HL 60, CEMSS, WEHI 3B, 3T3, K562, and human lymphocyte blood cells treated with Doxorubicin after 72 hrs.	101



3.13	Percentage of cell viability of HL 60, CEMSS, WEHI 3B, K562, human lymphocyte blood cell and 3T3 cells treated with arabinocytocine after 72 hrs.	102
3.14	MTT proliferation assay, effect of different concentrations (CD_{50} and CD_{75}) of NDV AF 2240 virus strain on the proliferation of WEHI 3B cell line at 24, 48 and 72 hrs post-inoculation.	106
3.15	The percentage of viable and non-viable WEHI 3B cells in the population after treatment with CD_{75} value of NDV AF2240 virus strain at various time intervals	106
3.16:	The percentage of viable and non-viable WEHI 3B cells in the population after treatment with CD_{50} value of NDV AF2240 virus strain at various time intervals	107
3.17	MTT proliferation assay, effect of different concentrations (CD_{50} and CD_{75}) of NDV V4-UPM virus strain on the proliferation of WEHI 3B cell line at 24, 48 and 72 hrs post-inoculation.	107
3.18	The percentage of viable and non-viable WEHI 3B cells in the population after treatment with CD_{75} value of NDV V4-UPM virus strain at various time intervals	108
3.19	The percentage of viable and non-viable WEHI 3B cells in the population after treatment with CD_{50} value of NDV V4-UPM virus strain at various time intervals	108
3.20	BrdU Proliferation assay, effect of different concentrations (CD_{50} and CD_{75}) of NDV AF 2240 virus strain on the proliferation and viability of WEHI 3B cell line at 24, 48 and 72 hrs post-inoculation.	109
3.21	BrdU Proliferation assay, effect of different concentrations (CD_{50} and CD_{75}) of NDV V4-UPM virus strain on the proliferation and viability of WEHI 3B cell line at 24, 48 and 72 hrs post-inoculation.	109
4.1	Agarose-gel-electrophoretic patterns showing DNA fragmentation of WEHI 3B cells treated with NDV strain AF 2240 at CD_{50} (2 HAU).	151
4.2	Agarose-gel-electrophoretic patterns showing DNA fragmentation of WEHI 3B cells treated with NDV strain V4-UPM at CD_{50} (8 HAU).	151



4.3	DNA fluorescence histograms of WEHI 3B cells treated with NDV strain AF 2240 at CD ₅₀ value (2 HAU) after staining with propidium iodide.	154
4.4	DNA fluorescence histograms of WEHI 3B cells treated with NDV strain V4-UPM at CD ₅₀ value (8 HAU) after staining with propidium iodide.	155
4.5	Density plots showing Annexin-V FITC staining of WEHI 3B cells treated with NDV strain AF 2240 at CD ₅₀ value (2 HAU)	159
4.6	Density plots showing Annexin-V FITC staining of WEHI 3B cells treated with NDV strain V4-UPM at CD ₅₀ value (8 HAU)	160
5.1	Experimental design of in vivo study	178
5.2	Amplification curve of SYBR Green I real time PCR detecting NDV strains AF 2240 and V4-UPM in the spleen, heart, liver and kidney of BALB/c mice.	220
5.3	Melting curve of SYBR Green I real time PCR detecting NDV strains AF 2240 and V4-UPM in spleen, heart, liver and kidney of BALB/c mice.	220
A.1	Percentage of necrotic, apoptotic and viable of WEHI 3B cells in the population after treated with CD ₅₀ of NDV AF2240 virus strain at different times.	270
A.2	Percentage of necrotic, apoptotic and viable of WEHI 3B cells in the population after treated with CD ₅₀ of NDV V4-UPM virus strain at different times	270
A.3	Cell cycle analysis of WEHI 3B cell population treated with NDV strain AF 2240 after staining with propidium iodide.	271
A.4	Cell cycle analysis of WEHI 3B cell population treated with NDV strain V4-UPM after staining with propidium iodide.	271
A.5	Effects of NDV strain AF 2240 on apoptosis of WEHI 3B cells analysed by Annexin-V FITC staining flow cytometry.	272
A.6	Effects of NDV strain V4-UPM on apoptosis of WEHI 3B cells analysed by Annexin-V FITC staining flow cytometry	272
B.1	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on survival time of BALB/c mice inoculated with WEHI-3B leukemia cells.	273



B.2	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on body weight of BALB/c mice inoculated with WEHI-3B leukemia cells	273
B.3	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the total leukocyte count in peripheral blood of BALB/c mice inoculated with WEHI-3B leukemia cells.	274
B.4	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the percentage of Blasts cells in peripheral blood of BALB/c mice inoculated with WEHI-3B leukemia cells.	274
B.5	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the percentage of blast cells in bone marrow of BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	275
B.6	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the percentage of blast cells in spleen of BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	275
B.7	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the spleen weight in the BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	276
B.8	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the spleen length in the BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	276
B.9	The effects of Ara-C, NDV strains AF 2240 and V4-UPM on the liver weight in the BALB/c mice inoculated with WEHI-3B leukemia cells for 30 days	277

