



UNIVERSITI PUTRA MALAYSIA

**PARTIAL CHARACTERISATION AND SOME PROPERTIES
OF A THERMOPHILIC BACTERIUM ISOLATED
FROM A PETROLEUM RESERVOIR
IN SARAWAK, MALAYSIA**

YOU LI LING

FSMB 1999 7

**PARTIAL CHARACTERISATION AND SOME PROPERTIES
OF A THERMOPHILIC BACTERIUM ISOLATED
FROM A PETROLEUM RESERVOIR
IN SARAWAK, MALAYSIA**

By

YOU LI LING

**Thesis Submitted in Fulfilment of the Requirements for
the Degree of Master of Science in the Faculty of
Food Science and Biotechnology
Universiti Putra Malaysia**

July 1999



Specially dedicated to :

My beloved

Daddy, Mummy and Brother,

Husband,

And Friends.



ACKNOWLEDGEMENTS

The author wishes to thank her family members for their love, support, encouragement and for always being there for her.

The author extends her deepest gratitude to the chairman of the supervisory committee, Dr. Raha Abdul Rahim, for her kindness, guidance, suggestions, and her willingness to help. The author also wishes to thank the members of the supervisory committee, Assoc. Prof. Dr. Arbakariya Ariff, Prof. Dr. Mohamed Ismail Abdul Karim and Dr. Mohammed Saifulaman Mohammed Said for their advice, supervision and guidance.

Special thanks are also conveyed to all the members of the Microbial Molecular Biology Laboratory, Animal Tissue Culture Laboratory, and Fermentation Laboratory.

Appreciation also given to Kak Madihah Md. Salleh for her guidance through out my project. I would like to thank En. Rosli and all the staffs in the fermentation Lab for their assistance and help.

I deeply grateful to the staff of the Electron Microscope Unit, namely Mr. Ho Oi Kuan and Cik Azilah Abd. Jalil for their unreserve help during Transmission and Scanning Electron Microscope study.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	vii
LIST OF FIGURES	ix
LIST OF PLATES	xi
LIST OF ABBREVIATIONS	xii
ABSTRACT	xiii
ABSTRAK	xv
CHAPTER	
I GENERAL INTRODUCTION	1
II LITERATURE REVIEW	5
Anaerobic Bacteria Adapted to Environmental Stress.....	5
Thermophiles	7
Microbial Species	10
Adaptation Mechanisms	13
Thermostable Enzyme	13
Membranes and Other Cell Components	15
Cell Envelopes of Archaeobacteria	15
Molecular Biology	16
Microbial Survival in Reservoir Environmental ...	20
Formulation of Liquid Enrichment Media	21
Carbon Sources	22
Nitrogen Sources	22
Inorganic Ions	23
Vitamins	24
Electron Acceptor	25
Solid Media	25
Harvesting of Cell Under Anaerobic Conditions	26
Preservation of Anaerobes	27
Preservation by Subculture	27
Preservation by Freezing	28
Preservation by Freeze-drying	29
Anaerobic Culture Conditions	30



	Anaerobic Jar Technique	31
	Plastic Bag Culture	32
	Indication of Anaerobiosis	32
	Hungate Method	33
	Anaerobic Chamber Method	34
	Application of Thermophilic Microorganisms	35
	Microbial Enhanced Oil Recovery	35
	Industrial Enzymes	37
III	IDENTIFICATION AND CHARACTERISATION OF BACTERIUM MRU5	40
	Introduction	40
	Materials and Methods	44
	Microorganism	44
	Gram Staining	44
	Spore Staining (Schaeffer-Fulton Method)	45
	Phase Contrast Microscopy	46
	Transmission Electron Microscopy	46
	Scanning Electron Microscopy	48
	Determination of End Products	49
	Determination of Organic Acids	49
	Solvent	49
	H ₂ S	49
	Effect of MRU5 Bacterium on Viscosity of Motor Diesel oil	50
	Results and Discussion	51
	End Product of Fermentation Using Different Carbon Sources	57
	Organic Acids	57
	Solvent	59
	H ₂ S Production	59
	Viscosity of the Motor Diesel Oil	60
	Conclusion	62
IV	KINETIC AND MODELLING OF GROWTH OF BACTERIUM MRU5 IN DIFFERENT MEDIUM FORMULATIONS AND CULTURE CONDITIONS ...	64
	Introduction	64
	Materials and Methods	66



Microorganism	66
Medium Preparation	66
Trace Element	66
Vitamin Solution	67
Formulated Enrichment Media	67
Preservation of the Microorganism	68
Measurement of Growth	68
pH, Temperature, and Sodium Chloride Ranges of Growth	69
Utilisation of Carbon Sources	69
Utilisation of Nitrogen Sources	70
Mathematical Method	70
Model Based on Gompertz equation.....	71
Fitting the Data	75
Potential Problems Fitting the Growth Curve ...	75
Results and Discussion	77
Effect of Temperature	77
Effect of pH	82
Effect of Salt Concentration	86
Utilisation of Carbon Sources	88
Utilisation of Nitrogen Sources	92
Conclusion	99
V GENERAL DISCUSSION, SUGGESTION FOR FURTHER WORK	100
REFERENCES	103
APPENDIX	114
1 Preparation of Vitamin Solution.....	115
2 Electron Microscopy	117
3 Chromatograms which Obtained from High Performance Liquid Chromatography	119
4 Plates	124
VITA	127



LIST OF TABLES

Table		Page
1	Historical Development of Thermophilic Microbiology	5
2	Comparison of Requirements for Optimal Growth of Anaerobes Adapted to Extreme Environmental Conditions	6
3	Habitats of Thermophiles	9
4	Taxonomy of Archae-hyperthermophiles	12
5	Enzymes from Anaerobic Hyperthermophiles	14
6	Examples of Redox Indicators	34
7	Examples and Properties of Some Commercially Important Enzymes from Thermophiles	38
8	Organic Acids Produced by Strain MRU5 During Growth on Different Carbon Sources	58
9	The Comparison Between MRU5 and <i>Hyperthermus sp.</i>	63
10	Nomenclature of High-temperature Bacteria	79
11	Growth Parameters of MRU5 Grown at Different Culture temperature as Calculated using Gompertz Model	80
12	Growth Parameter of MRU5 Grown at Different Culture pHs as Calculated Using Gompertz Model	84
13	Bacterial Yield Obtained from Different Carbon Sources	88
14	Growth Parameters of MRU5 Grown in Culture Containing Different Carbon Sources as Calculated According to Gompertz Model	90
15	Bacterial Yield Obtained from Different Nitrogen Sources	95



16	Bacterial Yield Obtained from Different Inorganic Nitrogen Sources	95
17	Comparison of Estimates of Growth Parameters of MRU5 Grown in Culture Containing Different Organic Nitrogen Sources	96



LIST OF FIGURES

Figure		Page
1	Schematic profiles of archaebacterial cell envelopes	18
2	Distribution of cell wall and cell envelope polymers among archae.....	19
3	Effect of bacterium MRU5 on viscosity of motor diesel oil ...	61
4	Effect of temperature on growth profile of strain MRU5	78
5	The fitness of the experimental data of growth of MRU5 cultivated at different temperatures to calculated data according to Gompertz model; (a) 60°C; (b) 70°C; (c) 80°C; (d) 90°C; and (e) 100°C	81
6	Effect of pH concentration on growth of strain MRU5	83
7	The fitness of the experimental data of growth of MRU5 cultivated at different pHs to calculated data according to Gompertz model (a) pH4; (b) pH5; (c) pH6; (d) pH7; (e) pH8; and (f) pH9	85
8	Effect of NaCl concentration on growth of strain MRU5	87
9	Effect of carbon sources on growth profile of strain MRU5 ...	89
10	The fitness of the experimental data of growth of MRU5 cultivated at different pHs to calculated data according to Gompertz model (a) xylose; (b) glucose; (c) fructose; (d) lactose;(e) arabinose; and (f) sucrose	91
11	Effect of organic nitrogen sources on growth profile of strain MRU5	93
12	Effect of inorganic nitrogen sources on growth profile of strain MRU5	94



13	The fitness of the experimental data of growth of MRU5 cultivated at different pHs to calculated data according to Gompertz model (a) yeast extract; (b) casein hydrolysate; (c) peptone; (d) NH_4NO_3 ; (e) $(\text{NH}_4)_2\text{SO}_4$	98
14	Standard lactic acid chromatogram	119
15	Standard oxalic acid chromatogram	119
16	Standard formic acid chromatogram	120
17	Standard acetic, propionic, and butyric acids chromatogram.	120
18	Chromatogram shows that organic acids produced by MRU5 in media containing glucose	121
19	Chromatogram shows that organic acids produced by MRU5 in media containing sucrose	121
20	Chromatogram shows that organic acids produced by MRU5 in media containing xylose	122
21	Chromatogram shows that organic acids produced by MRU5 in media containing fructose	122
22	Chromatogram shows that organic acids produced by MRU5 in media containing lactose	123
23	Chromatogram shows that organic acids produced by MRU5 in media containing arabinose	123



LIST OF PLATES

Plate		Page
1	A micrograph of light microscopy shows that MRU5 bacteria is a gram negative bacteria at 1,000 magnifications ..	52
2	A micrograph of phast contrast microscopy at 1,000 magnifications	52
3	Scanning electron micrograph of bacteria MRU5 at 15,000 magnifications	53
4	Negative stained bacteria MRU5 viewed under transmission electron microscopy at 50,000 magnifications .	53
5	Transmission electron micrograph of a section of MRU5 at 60,000 magnifications	54
6	A micrograph of scanning electron microscopy of MRU5 at 10,000 magnifications.....	54
7	Growth of bacteria in big bottles	124
8	Scanning Electron Microscope (JEOL 6400)	124
9	Transmission Electron Microscopy (Hitachi 7100).....	125
10	High Performance Liquid Chromatography (Shimadzu, SPD-10A).....	125
11	Gas Chromatography (GC-17A).....	126
12	Degassing System.....	126



LIST OF ABBREVIATIONS

OD	-	optical density
N ₂	-	nitrogen
CO ₂	-	carbon dioxide
O ₂	-	oxygen
K	-	potassium
Ca	-	calcium
Mg	-	magnesium
H	-	hydrogen
S	-	sulphur
H ₂ S	-	hydrogen sulphite
Na	-	sodium
Cl	-	chlorine
P	-	phosphorus
nm	-	nanometer
μm	-	micrometer
°C	-	degree celcius
mL	-	mililiter
M	-	molar
Eh	-	reduction potential
E'o	-	standard reduction potential
SEM	-	Scanning Electron Microscopy
TEM	-	Transmission Electron Microscopy
UV	-	ultraviolet
MEOR	-	microbial enhanced oil recovery



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Master of Science.

**PARTIAL CHARACTERISATION AND SOME PROPERTIES OF A
THERMOPHILIC BACTERIUM ISOLATED FROM A PETROLEUM
RESERVOIR IN SARAWAK, MALAYSIA**

By

YOU LILING

July 1999

Chairperson : Raha Abdul Rahim, Ph.D.

Faculty : Food Science and Biotechnology

The hyperthermophilic bacteria (MRU5) was isolated from an oil-producing well in Sarawak Malaysia. MRU5 is strictly anaerobic and was grown in enrichment media which were been degassed with CO₂ : N₂ (80 : 20). By Gram and spore staining, MRU5 was found to be a gram negative, non-spore producing bacteria. Scanning Electron Microscopy (SEM) and Transmission Electron Microscopy (TEM) revealed the shape of the bacteria to be irregular and coccoid which occurs singly, in pairs or clumps. The bacteria measures 0.75 µm in diameter and is surrounded by an outer sheath coating. MRU5 was found to grow best at pH 7 and 10% salinity with the optimal temperature range of between 90°C to 100°C. MRU5 was grown in xylose, glucose, fructose, arabinose, lactose, and sucrose. From the results obtained, xylose was the most efficiently utilised fermentable carbon source. Growth of the microorganism is supported by yeast extract. End products such as organic acids and H₂S were produced by fermentation of carbon sources but none of



the tested solvents were produced. This microorganism is able to reduce oil viscosity by producing organic acids. This means that MRU5 has the potential to be ecologically important for enhancing oil recovery. The modified Gompertz equation was found to be appropriate to model the growth of bacterium MRU5 at different medium compositions and culture conditions.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan bagi mendapat Ijazah Master Sains.

**PENCIRIAN SEPARA DAN SIFAT-SIFAT BAKTERIA THERMOFILIK
YANG DIPENCILKAN DARI TELAGA MINYAK
DI SARAWAK, MALAYSIA**

Oleh

YOU LI LING

Julai 1999

Pengerusi : Raha Abdul Rahim, Ph.D.

Fakulti : Sains Makanan dan Bioteknologi

Bakteria MRU5 merupakan sejenis bakteria hiperthermofilik yang disaring daripada salah sebuah telaga minyak di Sarawak, Malaysia. MRU5 adalah bakteria anaerobik, dan ditumbuhkan dalam media yang telah diperkayakan dan dinyahoksigen dengan menyalurkan campuran gas dalam nisbah CO₂ : N₂ (80 : 20). Penggunaan “Scanning Electron Microscopy” (SEM) dan “Transmission Electron Microscopy” (TEM) telah menunjukkan bakteria ini berbentuk kokus tidak sekata yang diliputi oleh lapisan selaput tebal, berdiameter 0.75 μm, dan ia wujud secara individu, berpasangan atau berkelompok. MRU5 didapati tumbuh pada keadaan optimum pada suhu 90°C, pH 7 dan dengan darjah kemasinan 10%. MRU5 ini boleh tumbuh dalam media yang mengandungi xylosa, glukosa, fruktosa, arabinosa, laktosa dan sukrosa. Keputusan menunjukkan xylosa dan estrak yis masing-masing

adalah sumber karbon dan nitrogen yang paling sesuai. Produk akhir yang dibebaskan daripada proses fermentasi sumber karbon adalah asid organik dan hidrogen sulfida. Asid organik yang dihasilkan berupaya mengurangkan kelikatan minyak. Ini menunjukkan MRU5 mempunyai potensi dalam meningkatkan pengeluaran minyak dari telaga minyak. Persamaan Gompertz yang diubahsuai adalah didapati sesuai digunakan sebagai model pertumbuhan bagi bakteria ini dalam komposisi media dan keadaan pertumbuhan yang berlainan.

CHAPTER I

GENERAL INTRODUCTION

Temperature is one of the most important variable in the environment. The classification of living microorganism based on their relation to temperature has therefore always been considered as one of the most basic elements of systematic biology. Microorganisms have traditionally been divided into three main groups in this respect, e.g., psychrophiles (-3°C to 20°C), mesophiles (13°C to 45°C) and thermophiles (42°C to 100°C or more).

Thermophilic microorganisms, which have an optimum temperature of more than 45°C , are found in many diverse habitats, such as thermal waters, soils and fermenting hay, hot water boilers and thermophilic biogas plant (Brock, 1978). The ranges of thermophiles still need to be divided further, and a relatively simple division is to define thermophilic and extremely thermophilic prokaryotes as those which can grow from $55-60^{\circ}\text{C}$ to $80-85^{\circ}\text{C}$ and hyperthermophiles as those which can grow optimally above $80-85^{\circ}\text{C}$. This division will put all known thermophilic bacteria (except Thermotogales) in the thermophilic range and the majority of the thermophilic archaeobacteria is in the hyperthermophilic range. Thermophiles, which have been of interest for long



time, both to scientists and to the public. This is understandable since these microorganisms can live and thrive under extreme conditions which most living microorganisms cannot tolerate. Thus, these microorganisms may contain extracellular and intracellular compounds such as special enzymes which are thermostable.

Thermophiles are believed to have a great potential in biotechnology and some products are already on the market. Thermostable DNA polymerase, for example, which was isolated from *Thermus aquaticus* are essential for the well-known PCR technology (Edwards, 1990).

Many enzymatic and microbiological industrial processes are carried out at high temperatures and those that are not could, in many cases, benefit from increased heating if not prevented by some unstable component in the process. The main advantages of increased temperature are generally; higher reaction rates, higher solubility of most chemicals, and increased fluidity and diffusion rates. The early notion that high temperatures prevents microbial contamination is only partly correct. It does, in fact, prevent growth of most environmental mesophilic microorganisms and pathogens but thermophilic microorganisms, especially spore formers, are surprisingly common contaminants in many fermentation processes which are run at elevated temperature.

Most of the microorganisms which have an ability to survive in oil-producing well can tolerate extreme environments. The survival of hyperthermophilic and hyperhalophilic microorganisms isolated from an oil-producing well is usually due to maintenance under strict anaerobic conditions. Various types of bacteria such as hydrocarbon-utilising bacteria and methane forming bacteria isolated from an oil producing well have several features with potential in enhancing oil recovery. The potential useful products of Microbial Enhanced Oil Recovery (MEOR) metabolism from these microorganisms are categorised into five general classes : polymers, biomass, organic acids, gases, and biosurfactants and solvents. Among those microorganisms, the most interesting microorganisms which contribute a lot in enhancing oil recovery are the fermentative bacteria. Direct utilisation of the fermentative bacteria on the sulphate and carbonates might promote the release of oil from those material. The fermentative bacteria produced acids which promoted the dissolution of carbonate rock and thus released the droplets of entrapped oil. Some of the bacteria also produced gases, including carbon dioxide, hydrogen and methane which increased the pressure and expel oil from the microtraps in oil reservoir.

Several hyperthermophilic bacteria have been isolated in Malaysia (Madiah *et al.*, 1996). A strain isolated from an oil-producing well in Sarawak was used in this study.

This project is carried out to achieve the following objectives:

- i) To characterise a thermophilic bacterium isolated from an oil-producing well in Malaysia.
- ii) To study the effect of each nutrient component in medium formulation and culture condition on growth of the isolated strain, and production of some metabolites such as organic acids and solvent.
- iii) To study the growth kinetics of the bacterium at different medium formulations and culture conditions.

CHAPTER II

LITERATURE REVIEW

Anaerobic Bacteria Adapted to Environmental Stress

A strong thread of incentive has dominated the search for new thermophilic archaeobacteria, the incentive to define experimentally the maximum temperature capable of supporting life. A steady rise in the maximum temperature for bacterial growth reflects the increased interest of the scientific community together with improvement in laboratory technique and sampling technology. Table 1 shows the historical development of thermophilic microbiology (Cowan, 1992).

Table 1 : Historical Development of Thermophilic Microbiology

Period	Growth temperature range (°C)	Representative microorganisms
Pre-1960s	37-65	<i>Bacillus stearothermophilus</i>
1960s-70s	60-85	<i>Bacillus caldolyticus</i> <i>Thermus</i> spp. <i>Thermoplasma acidophilum</i> <i>Sulfolobus</i> spp.
1980s	85-105	<i>Desulfurococcus</i> spp. <i>Thermoproteus tenax</i> <i>Pyrodictum</i> spp.

Prokaryotic microorganism can differ from eukaryotic cells in part because they have adapted to grow under extreme temperatures ($>100^{\circ}\text{C}$), salinity (saturated NaCl), pH ($<2.0, >10$), and substrate stress (limited chemical free energy or on toxicants). The origin of physiological biochemistry is described as follows; pH, neutral; temperature 37°C ; atmosphere, aerobic; salinity, 1.5%; substrate, glucose. Table 2 shows the comparison of requirements for optimal growth of anaerobes adapted to extreme environmental conditions.

Table 2 : Comparison of Requirements for Optimal Growth of Anaerobes Adapted to Extreme Environmental Conditions

Class	Growth requirement	Species example	Reference
Thermophile	High temperature ($>60^{\circ}\text{C}$)	<i>Thermoanaerobacter brokii</i>	Zeikus <i>et al.</i> , (1979)
Halophile	High salinity ($\geq 10\%$ NaCl)	<i>Haloanaerobium praevalens</i>	Zeikus <i>et al.</i> , (1979)
Acidophile	High acidity (external pH ≥ 2.0 , internal pH < 6.0)	<i>Sarcina ventriculi</i>	Smit, (1933)
Alkaliphile	High alkalinity	<i>Methanohalophilus zhilinae</i>	Mathrani, <i>et al.</i> , (1988)
Syntroph	Derives limited free energy and forms inhibitory catabolites (<10 kJ/mol of substrate consumed; H_2 , HCOOH, acetate formed), grows with a metabolic partner	<i>Syntrophospora bryantii</i>	Zhao, <i>et al.</i> , (1990)
CO utilizer or dehalogenator	Derives energy from substrate detoxification (CO as energy source)	<i>Butyribacterium methylotrophicum</i>	Lynd, <i>et al.</i> , (1982)

The starting materials for synthesising the organic matter under anoxic conditions included H_2S , CO, HCN which are poisonous gases for most aerobic

microorganisms, but these are important metabolites for certain anaerobic bacterial species. An assumption has been made that among microorganisms living today, obligate anaerobic bacteria are the simplest in structure and biochemistry, and are the most closely related to the earliest forms of life. In general, the physiological processes for adaptation to environmental stress in anaerobic bacteria seem to have involved differently from those in aerobic bacteria for two major reasons. First, anaerobes are energy limited during the chemoorganotrophic growth mode because they cannot couple dehydrogenation reactions to oxygen reduction and gain high level of chemical free energy (Thauer *et al.*, 1977). Second, growth of most chemoorganotrophic anaerobes (except for methanogens) is naturally associated with the generation of toxic end product (e.g., organic acids or alcohols, HS^-), which requires anaerobic species to develop certain dynamic adaptation mechanism or tolerance towards their catabolic end products.

Thermophiles

A number of natural habitats of high temperatures exist, ranging from sun-heated soils and littered with temperatures of 60°C to 70°C to erupting volcanoes, reaching 1000°C (Brock, 1967). Liquid-water environments with the highest known naturally occurring temperature is close to sea level, hence had boiling points around 100°C. Over the past two decades, habitats have been found at the



bottom of the ocean with temperature up to 350°C, raising questions about the possibility of life in these environments.

Thermophilic aerobic and anaerobic sporeforming bacteria were isolated from self-heating soil environments before the 1940s (Brock, 1986). About two decades ago, non-sporulating aerobic bacteria were detected in thermal springs of constantly high temperature, and the first thermophilic microorganism *Sulfolobus acidocaldarius*, an archaeobacterium with optimum temperature of above 80°C and upper growth temperature at around 92°C, was isolated (Brock *et al.*, 1972). Another archaeobacterium, *Methanobacterium thermoautotrophicum*, was the first nonsporulating thermophilic anaerobe described (Zeikus *et al.*, 1972).

The diversity and ecology of thermophilic anaerobes in thermal spring environments was first described in 1979 (Zeikus, 1979 ; Zeikus, *et al.*, 1979). Since then, hyperthermophilic anaerobes have been isolated from continental and submarine volcanic areas, such as solfatara fields, geothermal power plants, and geothermally heated sea sediments and hyperthermal vents (Stetter, 1986 ; Stetter and Zillig, 1985). Table 3 outlines the sites from which thermoanaerobes have been isolated. Interestingly, some microorganisms have been isolated from areas with temperature much higher than their maximum growth temperature, e.g., *Hyperthermus butylicus* and *Fervidobacterium islandicum*, which suggests that in these environments the microorganisms may not actively growing. The same