



**UNIVERSITI PUTRA MALAYSIA**

**PREVALENCE OF PATHOGENIC *VIBRIO* SPP AND MOLECULAR  
CHARACTERISATION OF SELECTED *VIBRIO CHOLERA*E  
SEROVARS ISOLATED FROM SEAFOOD IN MALAYSIA**

**NASRELDIN ELHADI HUSSEIN MOHAMED**

**FSMB 2002 6**

**PREVALENCE OF PATHOGENIC *VIBRIO* SPP AND MOLECULAR  
CHARACTERISATION OF SELECTED *VIBRIO CHOLERAE* SEROVARS  
ISOLATED FROM SEAFOOD IN MALAYSIA**

**By**

**NASRELDIN ELHADI HUSSEIN MOHAMED**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

**February 2002**

*Dedicated to my parents,  
my lovely son,  
my brothers  
my sisters  
and for Sitelbenat for being Sitelbenat*

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy.

**PREVALENCE OF PATHOGENIC *VIBRIO* SPP AND MOLECULAR CHARACTERISATION OF SELECTED *VIBRIO CHOLERAE* SEROVARS ISOLATED FROM SEAFOODS IN MALAYSIA**

**BY**

**NASRELDIN ELHADI HUSSEIN MOHAMED**

**February 2002**

**Chairman: Associate Professor Son Radu, Ph.D.**

**Faculty: Food Science and Biotechnology**

The global resurgence of cholera is an important public health challenge as the number of countries affected by this infection continues to increase. Cholera continues to be a major health challenge in Malaysia. To adequately control the infection requires a thorough understanding of its modes of transmission and implementing a comprehensive programme which takes into account not only the health aspect but also social, economic, behavioral and political dimension as well. The food-borne cholera and other pathogenic vibrios have not been thoroughly investigated in food in Malaysia. This study demonstrated the presence of toxigenic and non-toxigenic *Vibrio cholerae* O1, *Vibrio cholerae* O139, *Vibrio cholerae* non-O1/O139 and other vibrios pathogenic to mankind, which were isolated from seafood marketed from wet-markets and supermarkets from different locations in Malaysia from July 1998 to June 1999.

In this prevalence study, a total of 768 seafood samples were examined for the presence of pathogenic *Vibrio* species. A total of 431 of 768 (56.1%) seafood samples

were contaminated with *Vibrios*. The frequencies of *Vibrio* contamination in seafood were: shrimp (*Panaeus indicus*) 58.6%, squids (*Logio sp.*) 44%, squids (*Sepia sp.*) 60%, cockles (*Anadara granosa*) 82%, blue crabs (*Callinectes sapidus*) 62%, Shellfish (*Lithophaga malaccana*) 48% and peel mussels (*Perna viridis*) 24%. *V. damsela* and *V. alginolyticus* was the most frequently isolated species, followed by *V. meschnikovii*, *V. fluvalis*, *V. parahaemolyticus*, *V. vulnificus*, *V. cholerae* and *V. mimicus*. Of the 433 shrimp samples examined, 35.5% (254/433) were found to be positive for *Vibrio*, which yielded 791 isolates of *Vibrios*. Thus, 68.2 (58/85), 62.2 (28/45), 88 (22/25), 96 (24/25), 25 (5/20) and 100% (48/48) of the samples obtained from Selayang, Kajang, Bangi in Selangor; Seremban in Negri Sembilan; Penang (North Peninsula Malaysia); and Kuching in Sarawak (on Borneo island) yielded 158, 74, 62, 68, 14, and 231 isolates of *Vibrios*, respectively. Examination of 185 shrimp samples from Serdang wet market in Selangor over a nine-month period gave a contamination rate of 37.2% with incidence for the months ranged from 15 to 40%.

The incidence of *V. cholerae* in the various shrimp samples obtained from wet markets and supermarkets showed that a total of 14 (3.2%) and 3 (0.69%) of the 433 samples were contaminated with *V. cholerae* non-O1/O139 and O139, respectively. The highest number of *V. cholerae* non-O1/O139 were found in the shrimp samples collected from Kuching in Sarawak supermarkets with contamination rate of 10.4% (5/48) which yielded 20 isolates, followed by Kajang wet market (5.8% or 5/45) which yielded 16 isolates. The lowest rate of *V. cholerae* non-O1/O139 contamination were observed in Selayang wet market (2.2% or 1/45), Serdang wet market (1% or 2/185)

and Penang wet market (10% or 2/20). *V. cholerae* non-O1/O139 was not detected in shrimp samples collected from Bangi wet market and Seremban supermarkets. *V. cholerae* O139 was isolated only from Kuching location with contamination rate of 4.1% (3/48).

A total of 75 squids (*Logio sp*), 60 squids (*Spia sp*), 50 cockles (*Anadara granosa*), 50 blue crabs (*Callinectes sapidus*), 50 shellfish (*Lithophaga malaccana*), 50 peel mussels (*Perna viridis*) samples, were examined for the presence of pathogenic *Vibrios* at different wet markets and supermarkets in Selangor location [Petaling Jaya, Subang Jaya, Serdang, Seri Kembangan, Seri Petaling, and Kajang] in Malaysia. The contamination rate of *Vibrio* in Squids (*Logio sp*) and Squids (*Spia sp*) samples taken from Petaling Jaya and Subang Jaya supermarkets were 37.5% (15/40), 51.4% (18/35), 46.6% (14/30) and 73.3 (22/30) respectively. The cockles (*Anadara granosa*) collected from Petaling Jaya, Subang Jaya, Serdang and Seri Kembangan had contamination rate of 90% (9/10), 80% (8/10), 86.6% (13/15) and 73.3% (11/15) respectively. The frequencies of *Vibrios* contamination in blue crabs which were sampled from four locations; Seri Kembangan supermarket, Seri Petaling, Seri Kembangan wet market were 50% (5/10), 53.3% (8/15), 40% (2/5), 50% (5/10) respectively with the highest contamination observed in the 20 samples taken from Serdang wet market were all positive with a rate of 80% (16/20).

The prevalence of *V. cholerae* O1, O139 and non-O1/O139 from different types of seafood (squids, cockles, blue crabs, shellfish, peel mussels) sampled from different

location in Petaling Jaya, Subang, Selayang, Seri Petaling, Serdang, Seri Kembangan and Kajang from a total of 335 samples, gives contamination rates of 4.47% (15/335) which yielded 33 isolates of *V. cholerae* non-O1/O139 and 0.59% (2/335) which yielded 3 isolates of *V. cholerae* O139. *V. cholerae* O139 was recovered from cockles (*Anadara granosa*) from Serdang wet market and blue crab obtained from Seri Kembangan wet market, with contamination rate of 6.6% (1/15) which yielded 1 isolate, and 10% (1/10) which yielded 2 isolates, respectively. *V. cholerae* O1 serotype Ogawa was recovered from blue crab sample obtained from Serdang wet market with contamination rate of 5% (1/20) which yielded 1 isolate.

In this study growth survival of *V. cholerae* O1 was investigated in soya bean milk, coconut milk, glutinous rice, rice, shrimp and crab at different five storage temperature (4, 10, 15, 25 and 35 °C). The growth of *V. cholerae* O1 in different five selected food was very slow at 4, 10 and 15 °C. Growth of *V. cholerae* O1 was very significant in all food studied at 25 and 35 °C. *V. cholerae* O1 inoculated into crabs meat and shrimp entered the growth phase after 3 hours as compare to soya bean milk, rice, glutinous rice and coconut milk show growth phase after 12 hours.

In this study all strains of *V. cholerae* isolated from seafoods were screened for the antibiotic susceptibility and plasmid profiles. *V. cholerae* O1 and O139 isolates do not harbor plasmid, plasmid were detected only in two isolates of *V. cholerae* non-O1/O139 ranging in size from 3.4 to 4.6 megaDalton. The presence or absence of *ctx* and *tcpA* genes in all 97 isolates of *V. cholerae* were examined by using Multiplex PCR and

DNA colony hybridization test. Shrimp collected from Kuching in Sarawak resulting 20 isolates of *V. cholerae* non-O1/O139 and 11 isolates of *V. cholerae* O139. All isolates of *V. cholerae* O1 and O139 were positive for *ctx* and *tcpA* genes and belong to El Tor biotype. 19.5% of *V. cholerae* isolates were positive, and the result obtained by multiplex PCR is completely agreed with the result of DNA colony hybridization. The results obtained by PCR and DNA colony hybridization of positive strains of *V. cholerae* O1, O139 and non-O1/O139 isolated from seafood provide evidence that there is high priority of contamination of finding toxigenic strains of *V. cholerae* in seafood marketed in wet-markets and supermarkets in Malaysia. The results of this study demonstrated that toxigenic *V. cholerae* was detected in shrimp, cockles and crabs marketed in wet-markets and supermarkets and 19.5% of isolates contain genes encoding virulence factors. However, the significance of the presence of such *V. cholerae* O1, O139 and non-O1/O139 strains in seafood products in relation to potential hazards to public health is questionable. Further studies are required to determine the pathogenic potential and also the ecology and taxonomy of this species in Malaysia.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan ijazah Doktor Falsafah

**PREVALENCE OF PATHOGENIC *VIBRIO* SPP AND MOLECULAR CHARACTERISATION OF SELECTED *VIBRIO CHOLERAE* SEROVARS ISOLATED FROM SEAFOODS IN MALAYSIA**

Oleh

**Nasreldin Hussein Mohamed**

**February 2002**

**Pengerusi: Profesor Madya Dr. Son Radu**

**Fakulti: Sains Makanan dan Bioteknologi**

Kemunculan penyakit kolera secara global adalah cabaran kesihatan yang penting kerana bilangan negara yang dipengaruhi infeksi ini semakin meningkat. Kolera terus menjadi cabaran kesihatan yang utama di Malaysia. Pengertian yang mendalam tentang mod transmisi dan pelaksanaan program komprehensif yang melibatkan aspek kesihatan, social, ekonomi, tabiat dan dimensi politik adalah penting untuk mengawal infeksi ini. Penyakit kolera belum dikaji sepenuhnya di Malaysia. Kajian ini menunjukkan kehadiran *Vibrio cholerae* O1, *Vibrio cholerae* O139, *Vibrio cholerae* non-O1/O139 dan *Vibrio* lain yang patogenik kepada manusia yang dipencilkan daripada makanan laut di pasar malam dan pasaraya dari lokasi yang berbeza di Malaysia dari Julai 1998 hingga Jun 1999.

Di dalam kajian ini, sejumlah 768 sampel makanan laut telah dikaji untuk kehadiran spesis *Vibrio*. Sejumlah 431 daripada 768 (56.1%) sampel makanan laut telah

dikontaminasi dengan *Vibrio*. Frekuensi kontaminasi *Vibrio* di dalam makanan laut adalah: udang (*Panaeus indicus*) 58.6%, sotong (*Logio sp.*) 44%, sotong (*Sepia sp.*) 60%, tiram (*Anadara granosa*) 82%, ketam (*Callinectes sapidus*) 62%, shellfish (*Lithopaga malaccana*) 48% dan peeled mussels (*Perna viridis*) 24%. *Vibrio damsela* dan *V. alginolyticus* adalah sepsis yang paling banyak dipencilkan, diikuti oleh *V. meschnovii*, *V. fluvialis*, *V. parahaemolyticus*, *V. vulnificus*, *V. cholerae* dan *V. mimicus*. Daripada 433 sampel udang yang dikaji 35.5% (254/433) didapati positif untuk *Vibrio*, ini memberi jumlah sebanyak 791 pencilan *Vibrio*. Oleh kerana itu, 68.2 (58/58), 62.2 (28/45), 88 (22/25), 96 (24/25), 25 (5/20) dan 100% (48/48) sample yang diperolehi dari Selayang, Kajang, Bangi di Selangor; Seremban di Negeri Sembilan; Pulau Pinang (Utara Semenanjung Malaysia) dan Kuching di Sarawak (Kepulauan Borneo) memberi 158, 74, 62, 68, 14 dan 231 pencilan *Vibrio* masing-masing. Ujian ke atas 185 sampel udang dari pasar malam di Selangor selama 9 bulan memberi ladar kontaminasi sebanyak 37.2% di mana insiden untuk bulan-bulan tersebut adalah dalam lingkungan 15 hingga 40%.

Insiden *Vibrio* di malam pelbagai sampel udang yang diperolehi dari pasar malam dan pasaraya-pasaraya adalah sebanyak 14 (3.2%) dan 3 (0.69%) daripada 433 sampel yang dikontaminasi dengan *Vibrio cholerae* non-O139 dan O139, masing-masing. Bilangan tertinggi *Vibrio cholerae* telah dijumpai di dalam sampel udang yang diperolehi dari pasaraya di Kuching, Sarawak dengan kadar kontaminasi sebanyak 10.4% (5/48) dan memberi sebanyak 20 pencilan, diikuti dengan pasar malam Kajang (5.8% atau 5/48) yang memberi sebanyak 16 pencilan. Kadar kontaminasi *Vibrio cholerae* yang terendah diperhatikan di pasar malam Selayang (2.2% atau 1/45), pasar malam Serdang (1% atau

2/185) dan pasar malam Pulau Pinang (10% atau 2/20). *Vibrio cholerae* non-O139 tidak dikesan di dalam sample udang yang diperolehi dari pasaraya Bangi dan pasaraya Seremban. *Vibrio cholerae* O139 hanya dipencilkan dari lokasi Kuching dengan kadar kontaminasi sebanyak 4.1%.

Sejumlah 75 sotong (*Logio sp.*), 60 sotong (*Spia sp.*), 50 tiram (*Anadara granosa*), 50 ketam biru (*Callinectes sapidus*), 50 shellfish (*Lithopaga malaccana*), peeled mussels (*Perna viridis*) sampel telah dikaji untuk kehadiran *Vibrio* di pasar malam dan pasaraya yang berbeza di Selangor (Petaling Jaya, Subang, Serdang, Seri Kembangan, Seri Petaling dan Kajang) di Malaysia kadar kontaminasi *Vibrio* di dalam sotong (*Logio sp.*) dan sotong (*Spia sp.*) sampel diperolehi dari Petaling Jaya dan Subang Jaya adalah 37.5% (15/40), 51.4% (18/35), 46.6% (14/30) dan 73.3 (22/30) masing-masing. Tiram (*Anadara granosa*) yang dikumpul dari Petaling Jaya, Subang Jaya, Serdang dan Seri Kembangan mengalami kadar kontaminasi sebanyak 90% (9/10), 80% (8/10), 86.6% (13/15) dan 73.3% (11/15) masing-masing. Frekuensi kadar kontaminasi di dalam ketam biru yang diambil dari 4 lokasi iaitu pasaraya Seri Kembangan, Seri Petaling, pasar malam Seri Kembangan adalah 50%, 53.3%, 40%, 50% masing-masing dengan kadar kontaminasi tertinggi diperhatikan di dalam 20 sampel yang diambil dari pasar malam Serdang di mana kadarnya adalah 40%.

Taburan *Vibrio cholerae* non-O139 dan O139 dari pelbagai sampel makanan laut yang diperolehi dari lokasi yang berbeza di Petaling Jaya, Subang, Selayang, Sri Petaling, Serdang, Seri Kembangan dan Kajang merangkumi sebanyak 335 sampel dan memberi

kadar kontaminasi sebanyak 4.47% dimana 33 pencilan adalah *Vibrio cholerae* non-O139 dan 0.59% memberi 3 pencilan *Vibrio cholerae* O139. *Vibrio cholerae* O139 diperolehi dari tiram di pasar malam Serdang dan ketam biru diperolehi dari pasar malam Seri Kembangan dengan kadar kontaminasi 6.6% dan memberi satu pencilan dan 10% memberi 2 pencilan, masing-masing. *V. cholerae* O1 serotype Ogawa diperolehi daripada sampel ketam biru yang diperolehi dari pasar malam Serdang mengalami kadar kontaminasi 5% dan memberi 1 pencilan.

Kadar pertumbuhan *V. cholerae* O1 telah dikaji di dalam susu kacang soya, susu kelapa, nasi kanji, nasi, udang dan ketam pada lima suhu simpanan yang berbeza. Pertumbuhan *V. cholerae* O1 dalam kelima –lima sampel makanan yang berbeza adalah sangat perlahan pada suhu 4, 10 dan 15 C. Pertumbuhan *V. cholerae* O1 adalah amat ketara di dalam kesemua sampel makanan yang dikaji pada suhu 25 dan 35 C. *V. cholerae* O1 diinokulasi di dalam daging ketam dan udang memasuki fasa pertumbuhan selepas 3 jam berbanding dengan susu soya, nasi kanji dan susu kelapa yang menunjukkan fasa pertumbuhan selepas 12 jam.

Menurut kajian ini semua *V. cholerae* yang dipencilkan dari makanan laut telah diskriminasi untuk kerintangan antibiotik dan profil plasmid. Pencilan *V. cholerae* O1 dan O139 tidak mengandungi plasmid, plasmid hanya dikesan di dalam 2 pencilan *V. cholerae* O1 yang bersaiz 3.4 hingga 4.6 megadalton. Kehadiran atau ketidakhadiran gen CTX dan *TcpA* di dalam kesemua 97 pencilan *V. cholerae* yang dikaji menggunakan multiplex PCR dan ujian hibridasi koloni DNA. Udang yang dikumpul dari Kuching

Sarawak menghasilkan 20 pencilan *V. cholerae* non-O1/O139 dan 11 pencilan *V. cholerae* O139. Kesemua pencilan *V. cholerae* nonO1/O139 adalah positif dan keputusan ini diperolehi menggunakan multiplex PCR yang selari dengan keputusan ujian hibridasi koloni DNA. Keputusan yang diperolehi dari PCR dan hibridasi koloni DNA adalah positif untuk *V. cholerae* O1, O139 dan non-O1/O139 yang dipencilkan daripada makanan laut serta membuktikan bahawa adanya kontaminasi pencilan *V. cholerae* yang tinggi di dalam makanan laut yang dijual di pasar malam di dalam udang, tiram dan ketam yang dijual di pasar malam dan pasaraya dan 19.5% pencilan mengandungi gen faktor virulen. Walau bagaimanapun kehadiran signifikan *V. cholerae* O1, O139 dan non O1/O139 di dalam produk makanan laut yang berkaitan dengan potensi bahaya kepada kesihatan umum adalah satu persoalan. Kajian yang lebih mendalam perlu dilakukan untuk memastikan kesan patogenik, ekologi dan taksonomi spesis ini di Malaysia.

## ACKNOWLEDGMENTS

I would like to express my sincere gratefulness and appreciation to my supervisor, Associate Professor Dr. Son Radu, who has devoted much of his time for invaluable guidance, effort, supervision and support throughout this course study.

My sincere gratitude and appreciation also goes to the Dean of the Faculty of Food Science and Biotechnology, who is also my co-supervisor, Professor Dr. Gulam Rusul Rahmat Ali for his financial support, advice, suggestion and encouragement.

Sincere thanks are also extended to my other co-supervisors, Dr. Zaiton Hassan and Dr. Lum Keng Yeang for their support, encouragement, suggestion throughout the course of this study.

Special thanks also to Professor Dr. Mitsunori Nishibuchi from the Department of Microbiology, Faculty of Medicine, The Center for South East Asian Studies, Kyoto University, Japan, for his support, guidance and confirmation of the *Vibrio cholerae* strains used in this study.

Special thanks also extended to Professor Dr. Toshio Shimada from the National Institute of Infectious Diseases, Tokyo, Japan, for his suggestion and effort to confirm the serotypes of all *Vibrio cholerae* strains in this study.

Sincere gratitude is also extended to all the staff of the Faculty of Food Science and Biotechnology, Universiti Putra Malaysia, who contributed one way or another towards the completion of my study.

Specials thanks are also extended to all the post-graduate students in Laboratory of Food Safety and Molecular Typing, Faculty of Food Science and Biotechnology, Universiti Putra Malaysia: Yousr Abdulhadi Noaman for her help and support, Ahmad Zainuri, Samuel, Sushil Kaur, Abinya, Endang, Yuherman, Sahilah, Nizam, Kqueen, Wai Ling, Yock Ann, Noorlis, Yin Sze, Rozila, Siti Zulaiha and Noorzaleha.

Also, I would like to express my thank to my best friend and brother Fathi Hassan Miligi from Department of Computer Science, Sudan University, Sudan, for his support who accompanied me throughout the research and specially the technical problem during my writing: words can not express my gratitude for his love, support and patience. My thanks also extended to my friends Mr. Omer Ishag El Dai, Imad Eldin Abdo Abo Sham, Dr. Mohamed Sharief and Mohamed Abd Alla (Takinah).

Lastly, heartfelt appreciation to my parents, brothers, sisters and Sitelbenat Khalid Al-Bakhit for their love and support, encouragement who share much of my joy and sorrow.

**NASRELDIN ELHADI HUSSEIN**

I certify that an Examination Committee met on 21<sup>st</sup> February, 2002 to conduct the final examination of Nasreldin Elhadi Hussein Mohamed on his Doctor of Philosophy thesis entitled “Prevalence of Pathogenic *Vibrio* spp and Molecular Characterisation of Selected *Vibrio cholerae* serovars Isolated from Seafood in Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**SON RADU, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia  
(Chairman)

**GULAM RUSUL RAHMAT ALI, Ph.D.**

Professor/Dean  
Faculty of Food Science and Biotechnology,  
Universiti Putra Malaysia  
(Member)

**ZAITON HASSAN, Ph.D.**

Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**ABDUL REEZAL ABDUL LATIF, Ph.D.**


Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Lum Keng Yeang, Ph.D.**

Strategic, Environment and Natural Resource Research  
Malaysian Agricultural Research and Development Institute  
(Member)

**G. BALAKRISH NAIR, Ph.D, FNASc.**

Professor/Associate Director and Head  
Laboratory Sciences Division  
International Centre for Diarrhoeal Diseases Research, Dhaka, Bangladesh  
(External Examiner)



**SHAMSHER MOHAMAD RAMADILI, Ph.D.**

Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: **20 MAR 2002**



The thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for degree of Doctor of Philosophy.



---

**AINI IDERIS, Ph.D.**  
Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 13 JUN 2022

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
**NASRELDIN ELHADI HUSSEIN MOHAMED**

Date: 19-3-2002

## TABLE OF CONTENTS

		Page
DEDICATION.....		ii
ABSTRACT.....		iii
ABSTRAK.....		viii
ACKNOWLEDGEMENTS.....		xiii
APPROVAL.....		xv
DECLARATION.....		xvii
LIST OF TABLES.....		xxii
LIST OF PLATES.....		xxvi
LIST OF FIGURES.....		xxviii
<b>CHAPTER</b>		
<b>I</b>	INTRODUCTION.....	1
	The Objectives of This Study.....	7
<b>II</b>	LITERATURE REVIEW.....	8
	The genus <i>Vibrio</i> .....	8
	Taxonomy.....	9
	Pathogenic Vibrios.....	12
	<i>Vibrio cholerae</i> .....	12
	<i>Vibrio parahaemolyticus</i> .....	15
	<i>Vibrio vulnificus</i> .....	15
	<i>Vibrio alginolyticus</i> .....	16
	<i>Vibrio fluvialis</i> .....	17
	<i>Vibrio hollisae</i> .....	18
	<i>Vibrio mimicus</i> .....	18
	<i>Vibrio metschnikovii</i> .....	19
	Vibrios Mode of Infection.....	20
	Habitat.....	21
	Ecology of <i>Vibrio cholerae</i> .....	21
	Structure, Classification and Antigenic Types of <i>Vibrio cholerae</i> .....	23
	Environmental Survival and Persistence of <i>Vibrio cholerae</i> .....	26
	Enrichment Media for Isolation of <i>Vibrio cholerae</i> .....	28
	Plating Agar Media for <i>V. cholerae</i> and other Vibrios.....	29
	Street Food Vending in Malaysia.....	31
	Epidemiology of Cholera.....	32
	Seventh Pandemics of Cholera.....	33
	Eighth Pandemics of Cholera.....	34
	Epidemiology of Cholera in Malaysia.....	38

Control of Cholera.....	41
Food as a Medium for Survival and Vehicle transmission of <i>Vibrio cholerae</i> .....	43
Fruits and Vegetables.....	43
Seafood.....	44
Dairy Product.....	45
Poultry and Meat.....	45
Antibiotic Resistance Among Toxigenic <i>Vibrio cholerae</i> .....	46
Plasmid and Resistance.....	49
Molecular Epidemiology of <i>V. cholerae</i> .....	50
Virulence Factors Associated Genes in <i>V. cholerae</i> ..	52
TCP Pathogenicity Island.....	53
CTX Genetic Element.....	54
DNA Hybridization.....	56
Polymerase Chain Reaction.....	58
<b>III</b> <b>PREVALENCE OF PATHOGENIC <i>VIBRIO</i> SPECIES IN SEAFOOD.....</b>	<b>60</b>
Introduction.....	60
Materials and Methods.....	63
Sampling Methods....	63
Shrimp ( <i>Penaeus indicus</i> ) Samples.....	63
Squids ( <i>Logio</i> sp) Samples.....	63
Squids ( <i>Spia</i> sp) Samples.....	64
Cockles ( <i>Anadara granosa</i> ) Samples .....	64
Blue crabs ( <i>Callinectes sapidus</i> ) Samples..	64
Shellfish ( <i>Lithophaga malaccana</i> ) Samples.	64
Peel Mussels ( <i>Perna viridis</i> ) samples.....	65
Cultural Procedures.....	65
Identification Procedures for Vibrios.....	66
Biochemical Test Procedures.....	67
Elevated Temperature Enrichment Method for Isolation of <i>V. cholerae</i> .....	73
Principle of this Method.....	73
<i>Vibrio cholerae</i> Recovery.....	73
Preparation of Test Sample.....	73
Isolation procedures for <i>V. cholerae</i> .....	74
Treatment of Typical colonies to be <i>V. cholerae</i> .....	74
Serotyping .....	77
Detection of Protease.....	78
Kanagawa Test.....	79
RESULT.....	80
DISCUSSION.....	115

<b>IV</b>	<b>SURVIVAL AND GROWTH OF <i>VIBRIO CHOLERAE</i> O1 IN FOOD.....</b>	<b>121</b>
	Introduction.....	121
	Materials and Methods.....	124
	Culture.....	124
	Foods.....	124
	Media.....	124
	Preparation of Foods.....	125
	Soya bean milk.....	125
	Coconut Milk.....	125
	Glutinous Rice.....	126
	Rice.....	126
	Shrimp.....	126
	Blue Crab.....	127
	Inoculation and incubation.....	127
	Enumeration .....	128
	Results.....	129
	Growth of <i>Vibrio cholerae</i> O1 in Soya Bean Milk.....	129
	Growth of <i>Vibrio cholerae</i> O1 in Coconut Milk.....	130
	Growth of <i>Vibrio cholerae</i> O1 in Glutinous Rice.....	131
	Growth of <i>Vibrio cholerae</i> O1 in Rice...	131
	Growth of <i>Vibrio cholerae</i> O1 in Shrimp.....	137
	Growth of <i>Vibrio cholerae</i> O1 in Blue crab	137
	Discussion.....	142
<b>V</b>	<b>PLASMID PROFILING AND ANTIBIOTIC SUSCEPTIBILITY OF DIFFERENT SEROVARS OF <i>VIBRIO CHOLERAE</i> NON ISOLATED FROM SEAFOOD IN MALAYSIA.....</b>	<b>148</b>
	Introduction .....	148
	Materials and Methods.....	150
	Bacterial strains, media and Propagation.....	150
	Antibiotic Susceptibility.....	150
	Multiple Antibiotic Resistance (MAR)	
	Indexing of isolates.....	151
	Plasmid DNA Isolation.....	152
	Agarose gel Electrophoresis and Staining....	153
	Detection of Plasmid DNA Bands.....	153
	Molecular Weight determination of Plasmids	154
	Results.....	155
	Antibiotic susceptibility of <i>V. cholerae</i>	155

	isolates from seafood.....	
	Plasmid profiles of <i>Vibrio cholerae</i> .....	157
	Discussion .....	168
<b>VI</b>	<b>APPLICATIONS OF MULTIPLEX POLYMERASE CHAIN REACTION AND DNA COLONY HYBRIDIZATION TO DETECT CHOLERA TOXINS GENES.....</b>	<b>176</b>
	Introduction .....	176
	Material and Methods.....	179
	Bacterial Strains.....	179
	DNA Extraction.....	179
	Detection of Virulence Genes by PCR and Multiplex PCR.....	179
	DNA Hybridization.....	185
	Preparation of DNA Probe.....	185
	Colony Blot and DNA Hybridization.....	186
	Hybridization Method.....	187
	Prehybridization Method.....	187
	Post-hybridization and Detection of ..... Radiolabeled Probe.....	188
	Result.....	190
	Discussion.....	217
<b>VII</b>	<b>GENERAL DISCUSSION AND CONCLUSION...</b>	<b>224</b>
	REFERENCES.....	238
	BIODATA OF THE AUTHOR.....	268

## LIST OF TABLES

Table		Page
1	Vibrio species associated with human diseases.....	11
2	Foods implicated in foodborne vibrio infections.....	22
3	Main enrichment broth for isolation of <i>V. cholerae</i> , <i>V. parahaemolyticus</i> , <i>V. vulnificus</i> and other pathogenic Vibrio species.....	30
4	Cholera cases notified to WHO in 1999.....	35
5	Cholera cases notified to WHO in 1999 in Asia...	36
6	Cholera reported by state, Malaysia, 1995 .....	41
7	Eight key differential tests to differentiate the 12 pathogenic <i>Vibrio</i> species into six group.....	68
8	Differentiation of arginine dihydrolase positive <i>Vibrio</i> species and comparison with <i>Aeromonas</i> species.....	69
9	Differentiation of arginine dihydrolase negative, lysine decarboxylase positive <i>Vibrio</i> species.....	70
10	Biochemical reactions of <i>V. cholerae</i> .....	78
11	Prevalence of <i>Vibrio</i> in seafood from wet-markets and supermarkets.....	86
12	Incidence of Vibrios in shrimp at different wet-markets and supermarkets .....	90
13	Isolation of <i>Vibrio</i> species from shrimps sampled at Serdang wt-market over a 9 months period .....	91
14	Prevalence of <i>V. cholerae</i> non-O1/O139 and O139 in shrimp according to location.....	92
15	Result of serotyping and biotyping of <i>V. cholerae</i>	

	isolated from shrimp according to location.....	93
16	Distribution of <i>Vibrio</i> species isolates from shrimp ( <i>Penaeus indicus</i> ) according to location.....	96
17	Prevalence of <i>V. vulnificus</i> in shrimp according to location.....	97
18	Prevalence of <i>V. parahaemolyticus</i> in shrimp according to location.....	98
19	Prevalence of <i>V. alginolyticus</i> in shrimp according to location.....	99
20	Prevalence of <i>V. metschnikovii</i> in shrimp according to location.....	100
21	Prevalence of <i>V. mimicus</i> and <i>V. damsela</i> in shrimp according to location.....	101
22	Prevalence of <i>V. fluvalis</i> in shrimp according to location.....	102
23	Incidence of Vibrios in seafood at different wet-markets and supermarkets in Selangor, Malaysia...	103
24	Prevalence of <i>V. cholerae</i> non-O1/O139 from different types of seafood according to location...	104
25	Prevalence of <i>V. cholerae</i> O1 from different types of seafood according to location.....	105
26	Prevalence of <i>V. cholerae</i> O139 from different types of seafood according to location.....	106
27	Result of serotyping and biotyping of <i>V. cholerae</i> isolated from cockles and blue crab according to location.....	107
28	Prevalence of <i>V. parahaemolyticus</i> from different types of seafood according to location.....	109
29	Prevalence of <i>V. vulnificus</i> from different types of seafood according to location.....	110
30	Prevalence of <i>V. alginolyticus</i> from different types	111



	of seafood according to location.....	
31	Prevalence of <i>V. fluvalis</i> from different types of seafood according to location.....	112
32	Prevalence of <i>V. metschnikovii</i> from different types of seafood according to location.....	113
33	Prevalence of <i>V. damsela</i> from different types of seafood according to location.....	114
34	pH of food samples.....	127
35	Test of significance of plate count of <i>V. cholerae</i> O1 in soya bean milk.....	130
36	Test of significance of plate count of <i>V. cholerae</i> O1 in coconut milk.....	130
37	Test of significance of plate count of <i>V. cholerae</i> O1 in glutinous rice.....	131
38	Test of significance of plate count of <i>V. cholerae</i> O1 in rice.....	132
39	Test of significance of plate count of <i>V. cholerae</i> O1 in shrimp.....	137
40	Test of significance of plate count of <i>V. cholerae</i> O1 in blue crab.....	138
41	Antimicrobial resistance of <i>V. cholerae</i> O139 serotype Ogawa isolated from seafood.....	158
42	Antimicrobial resistance of <i>V. cholerae</i> non-O1/O139 serotype Ogawa isolated from seafood...	159
43	Plasmid profiles and antibiotic resistance patterns of <i>V. cholerae</i> O1 and O139 isolated from seafood.....	160
44	Plasmid profiles and antibiotic resistance patterns of <i>V. cholerae</i> non-O1/O139 isolated from seafood	160
45	Control strains of <i>V. cholerae</i> .....	181