



**UNIVERSITI PUTRA MALAYSIA**

**RAPID METHODS FOR ANALYSIS OF EDIBLE OILS AND FATS  
BY FOURIER TRANSFORM INFRARED SPECTROSCOPY**

**MOHAMED ELWATHIG SAEED MIRGHANI**

**FSMB 2002 14**

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

**RAPID METHODS FOR ANALYSIS OF EDIBLE OILS AND FATS  
BY FOURIER TRANSFORM INFRARED SPECTROSCOPY**

**MOHAMED ELWATHIG SAEED MIRGHANI**

**DOCTOR OF PHILOSOPHY  
UNIVERSITI PUTRA MALASIA**

**2002**



**RAPID METHODS FOR ANALYSIS OF EDIBLE OILS AND FATS  
BY FOURIER TRANSFORM INFRARED SPECTROSCOPY**

**By**

**MOHAMED ELWATHIG SAEED MIRGHANI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
Fulfilment of the Requirement for the Degree of Doctor of Philosophy**

**October 2002**



***ESPECIALLY DEDICATED TO MY BELOVED FAMILY***



Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirements for the degree of Doctor of Philosophy

**RAPID METHODS FOR ANALYSIS OF EDIBLE OILS AND FATS BY  
FOURIER TRANSFORM INFRARED SPECTROSCOPY**

By

**MOHAMED ELWATHIG SAEED MIRGHANI**

**October 2002**

**Chairman : Professor Dr. Yaakob B. Che Man**

**Faculty : Food Science and Biotechnology**

Analysis using Fourier transform infrared (FTIR) spectroscopy techniques on edible fats and oils extracted from palm fruit, groundnut, sesame seed, cottonseed and animal body fats rendered from cow, chicken, lamb and lard were investigated. The studies included development and applications of rapid FTIR techniques to determine some quality parameters such as moisture content in crude palm oil (CPO), soap and hexane residues in refined palm oil and groundnut oil, malondialdehyde (MDA) as a secondary oxidation product in refined palm oil, minor components such as sesamol and gossypol in sesame and cottonseed oils, and aflatoxins in groundnut and groundnut cake. The detection of lard in different mixtures with other animals' body fats such as cow, chicken and lamb was also investigated.

Different sample handling techniques were used such as transmission cells of NaCl, BF<sub>2</sub>, KBr and attenuated total reflectance (ATR) using internal reflectance element (IRE) of ZnSe. Partial Least Square (PLS) and Principle Component Analysis



(PCA) statistical methods were used to drive calibrations from FTIR versus actual or chemical values. In this study the frequency of 3700-3072  $\text{cm}^{-1}$  was used to determine moisture content in CPO as it indicates the absorption of compounds containing hydroxyl groups (OH). The frequency at 1675-1500  $\text{cm}^{-1}$  was used to determine soap residues in refined edible oils. For the determination of hexane residue in oils, the frequency used included all the data from 2935-2817  $\text{cm}^{-1}$ , 1490-1333  $\text{cm}^{-1}$  and 1200-1000  $\text{cm}^{-1}$  for  $-\text{CH}_3$  and  $-\text{CH}_2$ , and in-plane  $-\text{CH}$  bending.

In the determination of MDA as a secondary oxidation product, the correlation and variance spectra were used to select the best regions (2900-2800 and 1800-1600  $\text{cm}^{-1}$ ) to derive calibration from FTIR versus values obtained by chemical methods with SEC of 1.49. The spectral regions included the data from 3650-3000, 1600-1450 and 1200-900  $\text{cm}^{-1}$  that were used to determine sesamol in sesame seed oil. The study also included a qualitative and semiquantitative determination of palm and groundnut oils as adulterants in sesame seed oil using the spectral regions from 1504- 1503, 1400-1397 and 917-914  $\text{cm}^{-1}$ . The gossypol was also determined as an important quality factor in cottonseed oil and cakes using the spectral regions from 3600-2520 and 1900-800  $\text{cm}^{-1}$ . The study also covered the detection of lard in mixture of body fats of chicken, lamb and cow by using changes in frequency and absorbance of spectral regions 3009-3000, 1418-1417, 1385-1370, 1126-1085 and 966-967  $\text{cm}^{-1}$ . The simple Beer-Lambert law was used to develop equations for the determination of mixtures.

Aflatoxins exhibit characteristic absorption bands at wavelengths of 3004-2969  $\text{cm}^{-1}$  for  $\text{CH}_2$ , aromatic  $=\text{CH}$ ,  $-\text{C}-\text{H}$ ,  $\text{C}=\text{C}$  and phenyls, 1744-1720  $\text{cm}^{-1}$  for  $\text{C}=\text{O}$ , 1364-369  $\text{cm}^{-1}$  for methyl adjacent to epoxy ring, 1217-1220  $\text{cm}^{-1}$  for in plane  $-\text{C}-\text{H}$  bending.



CH bending of phenyl, 1035-1037  $\text{cm}^{-1}$  for symmetric stretching of =C–O–C or symmetric bending of phenyl, and 900-902  $\text{cm}^{-1}$  which may be for isolated H. In this calibration set the spectral regions that showed the highest correlation between concentration information and spectral response were set to include the data from 3000-2932, 1832-1693, 1400-1329 and 1250-1187  $\text{cm}^{-1}$  for aflatoxins B<sub>1</sub>, with standard errors of calibration (SEC) of 1.80 parts per million (ppm).

All of the results were in good correlation and of comparable accuracy to the classical wet chemical methods such as the American Oil Chemists Society (AOCS), Association of Official Analytical Chemists (AOAC) and International Union of Pure and Applied Chemistry (IUPAC) methods. This study represents the use of FTIR spectroscopy as a new rapid analytical technique developed for determination of some quality parameters of fats and oils, together with the detection of adulterants and contaminants. The FTIR spectroscopic technique has the potential to replace the time- and effort-consuming chemical methods for fast analysis of fats and oils. This can also eliminate the use of toxic chemicals that are hazardous to the analysts as well as to the environment in the analysis.





Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KAEDAH PANTAS UNTUK MENGANALISA MINYAK DAN LEMAK  
MASAK DENGAN PENGGUNAAN  
SPEKTROSKOPI FOURIER TRANSFORM INFRARED**

**Oleh**

**MOHAMED ELWATHIG SAEED MIRGHANI**

**Oktober 2002**

**Pengerusi: Profesor Yaakob Bin Che Man, Ph.D.**

**Fakulti: Sains Makanan dan Bioteknologi**

Kajian telah dijalankan untuk menganalisa minyak dan lemak yang diekstrak daripada buah kelapa sawit, kacang tanah, biji bijan, biji kapas dan lemak haiwan dari lembu, ayam, kambing dan khinzir dengan menggunakan Spektroskopi Fourier Transform Infrared (FTIR). Kajian ini merangkumi pembangunan dan penggunaan teknik pantas FTIR untuk menentukan beberapa parameter kualiti seperti kandungan lembapan dalam minyak sawit mentah, sabun, sisa heksana dalam minyak yang ditulin; kandungan aflatoksin dalam kacang tanah dan hampas kacang tanah; malondialdehid (MDA) dari 2900 – 2800 dan 1800 – 1600  $\text{cm}^{-1}$  sebagai produk oksidasi sekunder; dan komponen minor seperti sesamol dan gosipol dalam minyak bijan dan biji kapas. Pengesanan lemak khinzir dalam campuran lemak binatang lain seperti lembu, ayam dan kambing juga telah dijalankan.



Pelbagai teknik pengendalian sampel yang berlainan telah digunakan, contohnya sel-sel telusan NaCl, BF<sub>2</sub>, KBr dan pantulan keseluruhan « attenuated » (ATR) dengan menggunakan elemen pantulan dalaman (IRE) ZnSe. Partial Least Square (PLS) dan kaedah statistik Prinsip Analisis Komponen (PCA) telah digunakan untuk mengukur kalibrasi FTIR berbanding dengan nilai kimia atau nilai sebenar. Dalam kajian ini, frekuensi pada 3700–3072 cm<sup>-1</sup> telah digunakan untuk menentukan kandungan lembapan dalam minyak sawit mentah (CPO) kerana ia mewakili penyerapan gelombang bagi bahan yang mengandungi kumpulan hidroksil (OH) seperti air (H-OH). Frekuensi pada 1675–1500 cm<sup>-1</sup> telah digunakan untuk menentukan sisa sabun dalam minyak makan tulin. Manakala bagi penentuan sisa heksana dalam minyak, frekuensi yang digunakan termasuk semua data dari 2935–2817 cm<sup>-1</sup>, 1490–1333 cm<sup>-1</sup> dan 1200–1000 cm<sup>-1</sup> bagi CH<sub>3</sub> dan –CH<sub>2</sub> dan pembengkakan –CH dalam planar.

Bagi penentuan MDA sebagai produk oksidasi sekunder, korelasi dan variasi spektra telah digunakan untuk memilih kawasan spektra terbaik (2900–2800 dan 1800–1600 cm<sup>-1</sup>) untuk menghasilkan kalibrasi FTIR berlawanan nilai-nilai yang diperoleh dari kaedah kimia dengan SEC 1.49. Kawasan spektra merangkumi data dari 3650–3000, 1600–1450 dan 1200–900 cm<sup>-1</sup> telah digunakan untuk menentukan sesamol dalam minyak bijan. Kajian ini juga termasuk penentuan secara kualitatif dan semi-kuantitatif bagi minyak sawit dan minyak kacang tanah yang telah bercampur dengan minyak bijan (1504 - 1503, 1400 - 1397 dan 917 - 914 cm<sup>-1</sup>). Gosipol dalam minyak biji kapas juga telah ditentukan sebagai satu faktor kualiti penting dalam perdagangan minyak biji kapas dan hampas kapas (3600 - 2520 dan 1900 - 800 cm<sup>-1</sup>). Kajian ini juga meliputi pengesanan lemak khinzir dalam campuran lemak binatang

seperti ayam, kambing dan lembu dengan menggunakan perubahan frekuensi dan penyerapan kawasan spektra 3009-3000, 1418-1417, 1385-1370, 1126-1085 dan 966-967  $\text{cm}^{-1}$ . Kaedah Beer-Lambert telah digunakan untuk menghasilkan formula penentuan campuran.

Aflatoksin menunjukkan ciri rantai serapan pada jarak gelombang 3004–2969  $\text{cm}^{-1}$  bagi  $\text{CH}_2 = \text{CH}$  aromatik,  $-\text{C}-\text{H}$ ,  $\text{C}=\text{C}$  dan kumpulan fenil, 1477–1720  $\text{cm}^{-1}$  bagi  $\text{C}=\text{O}$ , 1364–11369  $\text{cm}^{-1}$  bagi kumpulan metil yang bercantum dengan gelang epoksi, 1217–1220  $\text{cm}^{-1}$  bagi fenil akibat pembengkokan  $-\text{CH}$  dalam planer, 1035 1037  $\text{cm}^{-1}$  bagi getaran simetri  $=\text{C}-\text{O}-\text{C}$  atau pembengkokan simetri fenil, dan 900–902  $\text{cm}^{-1}$  bagi  $-\text{H}$  yang telah diasingkan. Set kalibrasi bagi kawasan spektra maklumat kepakatan. Reaksi spektra telah ditetapkan dan merangkumi semua data dari 3000–2932  $\text{cm}^{-1}$ , 1832–1693  $\text{cm}^{-1}$ , 1400–1329  $\text{cm}^{-1}$  dan 1250–1187  $\text{cm}^{-1}$  bagi aflatoxin B<sub>1</sub>, dengan ralat standard kalibrasi (SEC) 1.80 ppm.

Semua keputusan kajian ini telah menunjukkan korelasi yang baik dan kejituan yang setanding dengan kaedah kimia klasik seperti dalam kaedah American Oil Chemists Society (AOCS), Association of Official Analytical Chemists (AOAC) dan International Union of Pure and Applied Chemistry (IUPAC). Kajian ini membuktikan bahawa spektroskopi FTIR adalah satu kaedah analitikal baru yang pantas bagi penentuan sesetengah parameter kualiti di dalam lemak dan minyak juga pengesanan bahan pencemaran. Teknik spektroskopi FTIR mempunyai potensi untuk menggantikan kaedah kimia yang memakan masa dan tenaga. Ini juga boleh mengelakkan penggunaan bahan kimia bertoksik yang membahayakan juru analisis dan alam sekitar.

## AKNOWLEDGEMENTS

I pray to Almighty ALLAH Subhanahu wa Ta'ala who give me the thoughts, the will, and guided me to complete this work. I pray that ALLAH will bless this work and make it useful for mankind, and that He will forgive us.

My sincere and deepest gratitude to Professor Dr. Yaakob B. Che Man, the chairman of my supervisory committee for his guidance, encouragement, patience and continuous follow up during the course of this study. My appreciation and gratitude is also extended to members of my supervisory committee, Prof. Dr. Jinap Selamat, Assoc. Prof. Dr. Jamilah Bakar and Assoc. Prof. Badlishah Sham Baharin for their advice, punctuate comments and support.

My gratitude is also due to all the staff of the Department of Food Technology, and the Faculty of Food Science and Biotechnology, UPM for their co-operation. My special appreciation is extended to my colleagues Dr. Irwandi Jaswir, Dr. Tan C. Ping, Ms. Gabby Setiowaty, Ms. Wanna Ammawath, Mr. Kambis Shams and Ms. Mariam Abdulatif for their kind help and friendly attitude.

I would like to acknowledge the financial support provided by IRPA fund for this study awarded to Prof. Dr. Yaakob B. Che Man. Acknowledgement is also due to the National Oilseed Processing Research Institute (NOPRI), University of Gezira, Sudan, especially Prof. Dr. Ismail H. Hussein, the Director of the Institute for granting me the opportunity to pursue my PhD studies.



I am gratefully indebted to the one who have lighted the way for me, my mother Haja Waqeya (Um-Aboha), my sisters, brothers and rest of our extended family for their support, encouragement and invaluable assistance.

Finally but first in my thoughts, I owe my sincere gratitude thanks to my beloved wife, Sabah and daughters Sarrah, Faizah and Doaa for their understanding, patience, care and love.



I certify that an Examination Committee met on 4<sup>th</sup> October 2002 to conduct the final examination of Mohamed Elwathig Saeed Mirghani on his Doctor of Philosophy thesis entitled “Rapid Methods for Analysis of Edible Oils and Fats by Fourier Transform Infrared Spectroscopy” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Suhaila Mohamed, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairperson)

**Yaakob B. Che Man, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Jinap Selamat, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Badlishah Sham Baharin**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Jamilah Bte Bakar, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Andrew Proctor, Ph.D.**

Professor  
Department of Food Science  
University of Arkansas, Fayetteville, Arkansas 72704, USA  
(Independent Examiner)

  
**SHAMSHER MOHAMAD RAMADILI, Ph.D.**

Professor/ Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia  
Date: 14 NOV 2002



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

**Yaakob B. Che Man, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Chairperson)

**Jinap Selamat, Ph.D.**

Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Badlishah Sham Baharin**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)

**Jamilah Bte Bakar, Ph.D.**

Associate Professor  
Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia  
(Member)



---

**AINI IDERIS, Ph.D.**

Professor/Dean


School of Graduate Studies

Universiti Putra Malaysia

Date: 9 JAN 2003

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



**Mohamed Elwathig Saeed Mirghani**

Date: Oct. 29th. 2002



## TABLE OF CONTENTS

DEDICATION .....	ii
ABSTRACT .....	iii
ABSTRAK .....	vi
ACKNOWLEDGMENTS .....	ix
APPROVAL SHEET .....	xi
DECLARATION .....	xiii
LIST OF TABLES .....	xix
LIST OF FIGURES .....	xxii
LIST OF ABBREVIATIONS .....	xxix
<b>CHAPTER</b>	
<b>I GENERAL INTRODUCTION .....</b>	<b>1</b>
<b>II LITERATURE REVIEW .....</b>	<b>5</b>
Infrared Spectroscopy .....	5
Vibration of Molecules .....	9
Instrumentation .....	11
History of Interferometers .....	11
Michelson Interferometer .....	11
Fourier Transformation .....	13
Advantages of Fourier Transformed IR Spectroscopy .....	15
Sample Handling .....	16
Transmission Technique .....	17
Attenuated Total Reflectance (ATR) .....	19
Polyethylene Infrared Cards .....	21
Data Handling Techniques .....	23
Detection of Overlapped Bands .....	23
Smoothing and Interpolation .....	23
Baseline Correction .....	23
Peak Intensity Measurements .....	24
Spectral Stripping .....	24
Ratio Method .....	25
Quantitative Analysis .....	25
Beer-Lambert Law .....	26
Classical Least Squares (K – Matrix) .....	27
Inverse Least Squares (P – Matrix) .....	27
Partial Least Square (PLS) .....	28
Principal Component Regression (PCR) .....	29
Validation .....	29
Estimation of Errors .....	30
Some FTIR Spectroscopy Applications .....	31
FTIR Spectroscopy Applications for Food and Lipids .....	32
Edible Fats and Oils .....	35



	Sources of Fats and Oils .....	36
	Properties and Characteristics of Fats and Oils .....	38
	Some Quality Parameters of Fats and Oils .....	38
	Analytical and Quality Control .....	40
	Moisture Content .....	42
	Residual Soap in Oil .....	44
	Hexane in Solvent Extracted Oil .....	45
	Aflatoxins .....	47
	Thiobarbituric Acid (TBA) Test .....	50
	Gossypol in Cottonseed Oil .....	52
	Sesamol in Sesame Seed Oil .....	53
	Multivariate Calibrations .....	55
	Authentication of Vegetable Oils Using FTIR Spectroscopy .....	55
	Detection of Lard Adulteration .....	57
<b>III</b>	<b>DETERMINING MOISTURE CONTENT IN CRUDE PALM OIL BY FOURIER TRANSFORM INFRARED SPECTROSCOPY .....</b>	<b>60</b>
	Introduction .....	60
	Materials and Methods .....	61
	Samples and Sample Preparations .....	61
	Analysis .....	62
	AOCS Vacuum Oven Method .....	62
	IUPAC Distillation Method .....	62
	FTIR Spectroscopy Scanning .....	63
	Validation .....	64
	Results and Discussion .....	65
	Moisture Content Obtained by AOCS Vacuum Oven and IUPAC Distillation Methods .....	65
	Absorption Bands of Water .....	65
	Conclusion .....	79
<b>IV</b>	<b>DETECTION OF SOAP RESIDUES IN REFINED VEGETABLE OILS BY FOURIER TRANSFORM INFRARED SPECTROSCOPY .....</b>	<b>80</b>
	Introduction .....	80
	Materials and Methods .....	82
	Samples and Chemicals .....	82
	Chemical Analysis .....	82
	Instrumental Analysis .....	82
	Statistical Analysis .....	84
	Validation .....	85
	Results and Discussion .....	85
	Chemical and FTIR Predicted Results .....	85
	Spectra .....	88
	Selection of the Optimal Frequency for Prediction .....	91
	Statistical Analysis .....	93
	Conclusion .....	101



<b>V</b>	<b>DETERMINING HEXANE RESIDUE IN EDIBLE OILS BY FTIR SPECTROSCOPY WITH ATTENUATED TOTAL REFLECTANCE</b> .....	102
	Introduction .....	102
	Materials and Methods .....	104
	Samples and Chemicals .....	104
	Spectra Acquisition .....	105
	Mathematical and Statistical Analysis .....	105
	Results and Discussion .....	106
	Development of Calibration Models .....	107
	Conclusion .....	118
<b>VI</b>	<b>DETERMINATION OF AFLATOXINS IN GROUNDNUT AND GROUNDNUT CAKE USING FTIR SPECTROSCOPY WITH ATTENUATED TOTAL REFLECTANCE</b> .....	119
	Introduction .....	119
	Materials and Methods .....	122
	Samples and Chemicals .....	122
	Extraction and Cleanup .....	122
	Thin Layer Chromatography (TLC) .....	123
	FTIR Method .....	125
	Spectra Acquisition .....	125
	Mathematical and Statistical Analysis .....	126
	Results and Discussion .....	126
	Development of Calibration Models .....	129
	Conclusion .....	143
<b>VII</b>	<b>DETERMINING TBARS IN PALM OIL USING MULTIVARIATE CALIBRATION OF FOURIER TRANSFORM INFRARED SPECTRA</b> .....	144
	Introduction .....	144
	Materials and Methods .....	146
	Materials .....	146
	Chemical Analysis .....	146
	Instrumental Analysis .....	146
	Statistical Multivariate Analysis .....	147
	Validation .....	148
	Results and Discussion .....	149
	Chemical Methods .....	149
	Spectra .....	150
	Optimal Frequency Region for Malonaldehyde Prediction .....	152
	Statistical Analysis .....	156
	Conclusion .....	162



<b>VIII</b>	<b>DETERMINING SESAMOL IN SESAME SEED OIL BY FOURIER TRANSFORM INFRARED SPECTROSCOPY .....</b>	<b>163</b>
	Introduction .....	163
	Materials and Methods .....	164
	Materials .....	164
	Samples .....	165
	Fourier Transform Infrared Spectra .....	165
	Results and Discussion .....	167
	Spectra .....	167
	Conclusion .....	175
<b>IX</b>	<b>DETECTION OF PALM AND GROUNDNUT OILS AS ADULTERANTS IN SESAME OIL BY FOURIER TRANSFORM INFRARED SPECTROSCOPY .....</b>	<b>176</b>
	Introduction .....	176
	Materials and Methods .....	177
	Materials .....	177
	Samples .....	178
	HPLC Analysis of TAG .....	178
	FTIR Spectra .....	179
	Results and Discussion .....	179
	FTIR Spectra .....	182
	Conclusion .....	189
<b>X</b>	<b>DETECTION OF GOSSYPOL IN COTTONSEED OIL BY FOURIER TRANSFORM INFRARED SPECTROSCOPY .....</b>	<b>190</b>
	Introduction .....	190
	Materials and Methods .....	192
	Materials .....	192
	Chemical Analysis .....	192
	Instrumental Analysis .....	193
	Statistical Analysis .....	193
	Validation .....	194
	Results and Discussion .....	194
	Chemical Method .....	194
	FTIR Spectra .....	195
	Development of Calibration Models .....	196
	Conclusion .....	203
<b>XI</b>	<b>DETECTING LARD ADULTERATION OF BODY FATS OF CHICKEN, LAMB AND COW BY FOURIER TRANSFORM INFRARED SPECTROSCOPY .....</b>	<b>204</b>
	Introduction .....	204
	Materials and Methods .....	206
	Sample Preparation .....	206
	Instrumentation/Spectral Acquisition .....	207



Results and Discussion .....	208
Lamb Body Fat (LBF) .....	208
Chicken Fat (CF) .....	222
Cow Body Fat (CBF) .....	232
Conclusion .....	240
<b>XII CONCLUSIONS AND RECOMMENDATIONS .....</b>	<b>241</b>
Summary .....	241
Conclusions and Recommendations .....	246
<b>REFERENCES .....</b>	<b>248</b>
<b>APPENDICES .....</b>	<b>268</b>
<b>BIOGRAPHICAL SKETCH .....</b>	<b>274</b>



## LIST OF TABLES

Table		Page
1	Characteristic mid-infrared absorption of some common functional groups in edible fats and oils .....	12
2	Calibration and cross validation statistics for moisture content values measured by AOCS and IUPAC methods .....	65
3	Result of the developed calibration models of AOCS and IUPAC methods at the wavelength region (3074-3700 $\text{cm}^{-1}$ ) for moisture content of CPO samples <sup>a</sup> .....	71
4	Calibration statistics for moisture content values obtained by classical references and FTIR methods <sup>a</sup> .....	75
5	A general comparison between the three methods used in the study .....	77
6	Calibration and cross-validation for soap content in palm and groundnut oils by FTIR methods in comparison with chemical methods .....	87
7	Effect of different wavelength regions in developing calibration model for the determination of soap in palm and groundnut oils .....	93
8	Results from PLS calibration models using the 1675 to 1500 $\text{cm}^{-1}$ region in the FTIR spectrum to determine the soap content in palm and groundnut oils .....	94
9	Calibration statistics for soap content from data obtained by chemical analysis and FTIR spectroscopy .....	96
10	Calibration and cross-validation for hexane content in palm and groundnut oils by the FTIR method .....	111
11	Effect of different wavelength regions in developing calibration model for determining hexane residues in palm and groundnut oils .....	113
12	Results from calibration models using PLS of wavelength regions 2935 to 2817, 1490 to 1334 and 1200 to 1000 $\text{cm}^{-1}$ to determine the hexane content in palm and groundnut oils .....	114



13	Calibration statistics for hexane content from data obtained by GC analysis and FTIR spectroscopy .....	116
14	Calibration and validation statistics for aflatoxins determined by AOCS TLC and FTIR/ATR spectroscopic methods .....	129
15	Results of calibration models of AOCS TLC reference method for aflatoxins in reference standards (groundnut and groundnut cake) and FTIR/ATR methods .....	133
16	Calibration statistics for aflatoxin content in groundnut and groundnut cake obtained by classical TLC and the FTIR/ATR methods .....	135
17	Calibration and cross-validation statistics for malondialdehyde (TBARS) content in palm oil obtained by the PLS and PCR methods with zero (none) baseline .....	150
18	Results accuracy of prediction by the PLS and PCR calibration models of the reference method assessed against the FTIR spectroscopic method at the frequency regions 2900 - 2800 $\text{cm}^{-1}$ and 1800 - 1600 $\text{cm}^{-1}$ with different baseline types for malondialdehyde in palm oil .....	155
19	Calibration statistics for malondialdehyde (TBARS) content in palm oil obtained by reference and FTIR methods using the PLS and PCR techniques .....	158
20	SEP ratios of the different FTIR methods and their F-values (critical) at 95% confidence level .....	160
21	Calibration and cross-validation statistics for sesamol content in sesame groundnut and RBDPOo oils measured by FTIR spectroscopic method ...	171
22	Result of the developed calibration models for sesamol content of sesame groundnut and RBDPOo oils at the wavelength ranges 3650 – 3000, 1600 – 1450 and 1200 – 900 $\text{cm}^{-1}$ .....	172
23	Distribution of trisaturated, monounsaturated, diunsaturated, and triunsaturated triacylglycerols (TAG) in sesame, palm, groundnut oils and their blends .....	181
24	Sets of samples and frequency of bands a, b, c and d in $\text{cm}^{-1}$ of the FTIR spectra of sesame oil, Groundnut oil, Palm oil and mixtures .....	184
25	Gossypol contents determined by the AOCS and FTIR methods .....	195



26	Calibration and cross-validation using PLS of wavenumber regions 3600 to 2520 and 1900 to 800 $\text{cm}^{-1}$ for gossypol content in cottonseed oil samples .....	200
27	Calibration statistics for gossypol content using data obtained by chemical analysis and FTIR spectroscopy .....	200
28	Band frequencies in the Fourier transform infrared (FTIR) spectra of lamb body fat (LBF), lard and their blends in region (3009-3000 $\text{cm}^{-1}$ ) ...	214
29	Absorbance values in the FTIR spectra of LBF, lard, and their blends in region b (1418-1417 $\text{cm}^{-1}$ ) .....	216
30	Absorbance values in the FTIR spectra of CF, lard and their blends at 1417.85 $\text{cm}^{-1}$ and 1377.58 $\text{cm}^{-1}$ .....	229
31	Absorbance values in the FTIR spectra of cow body fat, and their blends at 966.22 $\text{cm}^{-1}$ .....	239





## LIST OF FIGURES

Figure		Page
1	Energy levels .....	6
2	Part of the electromagnetic radiation .....	8
3	Stretching and Bending Vibration Modes .....	10
4	Optical Diagram of Michelson Interferometer .....	14
5	Schematic representation of infrared transmission cells .....	18
6	Internal reflection element (IRE) .....	20
7	Schematic representation of an attenuated total reflectance (ATR) accessory .....	20
8	Polyethylene Infrared (IR) Card .....	22
9	Chemical structure of glycerol and triacylglycerol (TAG) .....	37
10	Mean spectrum of calibration set .....	67
11	Correlation spectrum, showing better correlation of water absorption at $3700 - 3074 \text{ cm}^{-1}$ than $1700 - 1500 \text{ cm}^{-1}$ .....	68
12	Variance spectrum obtained from the calibration set, determined by calculating the average absorbance at each wavenumber position over the entire calibration data set and then calculating the square root of variance about that mean for the entire data set generated the variance spectrum .....	69
13	Absorption changes at $3700 - 3074 \text{ cm}^{-1}$ with the changes in moisture content .....	70
14	American Oil Chemists' Society (AOCS) vacuum oven method compared with Fourier transform infrared (FTIR) spectroscopic predicted values, calculated with partial least square (PLS) calibration ...	73
15	International Union of Pure and Applied Chemistry (IUPAC) distillation method compared with FTIR spectroscopic predicted values, calculated	

