



UNIVERSITI PUTRA MALAYSIA

**THERMOSTABILITY AND PROTEIN STUDIES OF
NEWCASTLE DISEASE VIRUS**

ZURIDAH HASSAN

FSAS 1995 4

**THERMOSTABILITY AND PROTEIN STUDIES OF
NEWCASTLE DISEASE VIRUS**

By

ZURIDAH HASSAN

Thesis Submitted in Fulfilment of the Requirements for
the Degree of Master of Science in the Faculty of
Science and Environmental Studies,
Universiti Pertanian Malaysia.

October 1995



ACKNOWLEDGEMENTS

I am indebted to Assoc. Prof. Dr. Khatijah Mohd Yusoff for her guidance, advice, never-ending patience and many valuable discussions throughout the course of this work. I would also like to thank Assoc. Prof. Dr. Nor Aripin Shamaan for his technical support and comments and to Assoc. Prof. Dr. Norani Abdul Samad for constructive remarks and for access to use all the facilities in Lab 143 .

I would also like to thank various individuals and institutions who have helped me during the actual study:

Dr. Bill Jordan (Victoria University, New Zealand) who during his visit to Malaysia had spent some time in Lab 143 together with Dr. Khatijah and Dr. Nor Aripin to introduce to me the 2D-PAGE systems;

En. Kamudin and his team from INTAN, Bukit Kiara who helped me to get through the INTAN/JPA Quantitative test;



En. Abdul Ghani of the Faculty of Science and Environmental Studies for all the photography work;

The Deputy Director, Dato Dr. Ahmad Tajuddin, Hospital Kuala Lumpur and the Director, Dr. G. Duraisamy, National Blood Services Centre, Hospital Kuala Lumpur for granting the study leave;

All lab-mates, staff of Lab 143 (En. Ariffin, En. Husin, Omar, Rohana, Izan, Mazidah, Chin Hoon, Wen Siang, Goh, Najah, Kri, Mages, Sudani, Suzila, Fizah, Ban Kim, Muhajir), Nona, En. Karim, Syarifah, Zaharah and staff of the Graduate School, UPM.

To my husband, Syed Abdul Razak, my children and parents whose love and understanding have greatly helped me in my studies, I dedicate this thesis and Syukur Alhamdulillah, God has given me the will, patience and health to finish this work.

This study was sponsored by Jabatan Perkhidmatan Awam, Malaysia and partly by IRPA.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	ii
LIST OF TABLES	viii
LIST OF FIGURES	ix
LIST OF PLATES	x
LIST OF ABBREVIATIONS	xii
ABSTRACT	xiv
ABSTRAK	xvi
 CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	5
Newcastle Disease	5
Viral Structure	7
NDV Proteins	8
HN Protein	12
F Protein	13
M Protein	14
L Protein	15
P Protein	15
NP Protein	16



Thermostability of NDV Strains of Different Virulence	17
Separation of NDV Proteins by SDS-PAGE and 2D-PAGE	19
Peptide Mapping of NDV Polypeptides	20
Western Blot Analysis of NDV Proteins	22
III MATERIALS AND METHODS	24
Chemicals, Enzymes and Antibodies .	24
Viruses	24
Virus Cultivation	24
Inoculation Technique	25
Harvest Technique	27
Clarification of Allantoic Fluid	27
Purification	28
Protein Assays	28
Haemagglutination Assay	29
Preparation of Red Blood Cells for HA and Hemolysis Assays	30
Hemolysis	31
Neuraminidase Assay	31
Sodium Dodecyl Sulphate- Polyacrylamide Gel Electrophoresis (SDS-PAGE)	32
Casting of Discontinuous Polyacrylamide Gels	32
Sample Preparation	33



Peptide Mapping	34
Two-Dimensional Analysis of NDV Proteins	35
Sample Preparation for First- dimension (IEF) Run	36
Preparation of IEF Gels	36
Prefocussing of IEF Gels	37
Second-dimensional Analysis by SDS-PAGE	38
Western Blotting	40
Preparation for Blotting of SDS-PAGE and 2D-PAGE	40
Assembly of the Trans-Blot Semi-Dry Transfer Cell	40
Immunological Detection of Protein on Nitrocellulose Membrane	41
IV RESULTS AND DISCUSSION	44
Thermostability of NDV Strains ...	44
Stability of Haemagglutinating Activity	48
Inactivation Constant for HA Activity	50
Stability of Neuraminidase Activity	52
Stability of Hemolytic Activity	56
Analysis of NDV Proteins by SDS-PAGE and 2D-PAGE	59
Optimization of 2D-PAGE	64
Computer Analysis	67
Peptide Mapping	76



V	GENERAL DISCUSSION	82
VI	CONCLUSION	93
	REFERENCES	96
	APPENDICES	109
	A List of Enzymes and Chemicals ..	110
	B Protein Estimation by Lowry's Method	112
	C Additional Plates	113
	D Computer Printouts	116
	VITA	132



LIST OF TABLES

Table		Page
1	Functions of NDV Coded Proteins ...	11
2	Examples of Some Applications of Protein Analysis by 2D-PAGE	21
3	Stability of Hemagglutinating Activities of NDV Strain in 30 min of Heat Stress	46
4	Stability of Hemagglutinin of NDV Strains at 50°C	49
5	Comparison of Inactivation Constant (k per min) for the Haemagglutinating Activity in Various NDV Strains	53
6	Neuraminidase Activities of NDV Strains at 50°C	54
7	Hemolytic Activities of Various NDV Strains at 50°C	57
8	Summary of Protein Spots' x-coordinate Position on Nitrocellulose Membrane	74



LIST OF FIGURES

Figure		Page
1	Newcastle Disease Virion and Genome Structure	10
2	Inoculation of Chick Embryo via the Allantoic Cavity	26
3	Hemagglutinating Activities in NDV Strains at Different Temperature	47
4	Hemagglutinating Activities of Different NDV Strains at 50°C ...	51
5	Neuraminidase Activities of NDV Strains at 50°C	55
6	Hemolytic Activities of NDV Strains	58



LIST OF PLATES

Plate		Page
1	Electrophoretic Pattern of NDV Proteins Stained With Coomassie Brilliant Blue	60
2	Polypeptides of NDV Separated by SDS-PAGE	61
3	Polypeptides of NDV Separated by SDS-PAGE, Blotted and Detected Using Anti-F ₁ Monoclonal Antibodies	62
4	Polypeptides of NDV Separated by SDS-PAGE, Blotted and Detected Using Anti-P Monoclonal Antibodies	63
5	Two-Dimensional Gel Electrophoresis of NDV Strain V4 Stained by Silver Stain ..	66
6	Nitrocellulose Sheets Following Western Blotting of 2D-PAGE Gel Developed With Anti-HN Monoclonal Antibodies	69
7	Nitrocellulose Sheets Following Western Blotting of 2D-PAGE Gel Developed With Anti-HN Monoclonal Antibodies (Duplicate Set)	70
8	Nitrocellulose Sheets Following Western Blotting of 2D-PAGE Gel Developed With Anti-NP Monoclonal Antibodies	71
9	Nitrocellulose Sheets Following Western Blotting of 2D-PAGE Gel Developed With Anti-F ₁ Monoclonal Antibodies	72



10	Peptide Map of HN Proteins Digested With <i>Staphylococcus aureus</i> V8-Protease ..	77
11	Peptide Map of NP/P/F ₁ and M Proteins Digested with <i>Staphylococcus aureus</i> V8-Protease	79
12	Peptide Map of HN Protein Digested With <i>Pseudomonas fragi</i> Endoproteinase Asp-N in Various NDV Strains	80
13	Peptide Map of HN Proteins Digested With <i>Lysobacter enzymogenes</i> Endoproteinase Lys-C	81
14	Mini Protean II Dual Slab Cell	113
15	Mini Protean II 2-D Cell	114
16	View of the Trans-Blot SD Cell	115



LIST OF ABBREVIATIONS

ND	Newcastle disease
NDV	Newcastle disease virus
2D-PAGE	two-dimensional polyacrylamide gel electrophoresis
ts	temperature-sensitive
RNA	ribonucleic acid
mRNA	messenger RNA
MW	molecular weight
Kbp	kilobase pair
pI	isoelectric point
SDS	sodium dodecyl sulphate
SDS-PAGE	sodium dodecyl sulphate-polyacrylamide gel electrophoresis
Mab	monoclonal antibody
UPM	Universiti Pertanian Malaysia
JPA	Jabatan Perkhidmatan Awam
BSA	bovine serum albumin
HA	hemagglutinating activity
HL	hemolytic activity
NA	neuraminidase activity
NANA	N-acetylneuraminic acid



RBC	red blood cells
DTT	dithiothreitol
CHAPS	cholamidopropyldimethyl- hydroxypropanesulfonate
IEF	isoelectric focusing
TEMED	tetramethylethylenediamine
ncm	nitrocellulose membrane
NBT	nitrobluetetrazolium
BCIP	5-bromo-4-chloro-3-iodolyl-phosphate
EDTA	Ethylenediaminetetraacetic acid disodium salt
R_f	relative mobility
V	volt
Vh	volt-hour
v/v	volume/volume



Abstract of thesis submitted to the Senate of Universiti Pertanian Malaysia in fulfilment of the requirements for the degree of Master of Science.

**THERMOSTABILITY AND PROTEIN STUDIES OF
NEWCASTLE DISEASE VIRUS PROTEINS**

By

ZURIDAH HASSAN

October 1995

Chairman: Assoc. Prof. Dr. Khatijah Mohd Yusoff

Faculty: Science and Environmental Studies

The heat stability of four strains of Newcastle disease virus (NDV) belonging to three different pathotypes were studied. The strains were the lentogenic V4 and its heat stable variant V4-UPM, the mesogenic S strain and the velogenic strain AF2240. Analyses of their haemagglutination and neuraminidase activities (which are the functions of the HN protein) and the hemolytic activities at various temperatures showed that strains AF2240, V4 and V4-UPM were heat stable compared to strain S.

There were no differences observed in the mobilities of the various NDV proteins on sodium



dodecyl sulphate-polyacrylamide gel electrophoretic (SDS-PAGE) studies. However, analysis of the various peptides with *Staphylococcus aureus* protease showed that the digested HN proteins of strain V4-UPM was different from the strains V4, AF2240 and S. The peptide analysis was repeated using *Pseudomonas fragi* *Endoproteinase Asp-N* and *Lysobacter enzymogenes Lys-C* and found to be similar except in strain V4-UPM.

These proteins were further analysed by the two-dimensional polyacrylamide gel electrophoresis (2D-PAGE). The gels were then Western blotted and protein spots were identified using HN, NP and F₁ Mabs and then analysed by the UVP GDS Gel Documentation System (United Kingdom). It was observed that in the heat stable V4 strain the isoelectric point (pI) of the HN protein was in the acidic region, strains V4-UPM and AF2240 in the neutral/weak basic regions and in the thermolabile strain S, the HN protein was shifted to the basic end of the isoelectric focussing run. The pI changes in the NP protein was seen in strain S only. The F protein was at the basic region for all strain except strain V4-UPM. In strain S it was seen that the HN, NP and F proteins were in the basic region and this basic pI could be responsible for the different biological characteristics seen in the thermostable strains compared to the thermosensitive strain.



Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia untuk memenuhi keperluan Ijazah Master Sains.

**KESTABILAN SUHU DAN KAJIAN PROTIN
VIRUS NEWCASTLE DISEASE**

Oleh

ZURIDAH HASSAN

Oktober 1995

Pengerusi: Prof. Madya Dr. Khatijah Bt Mohd Yusoff

Fakulti: Sains dan Pengajian Alam Sekitar

Kajian keatas empat strain virus Newcastle disease (NDV) terdiri daripada tiga jenis patotip telah dijalankan. Strain tersebut adalah strain lentogenik V4 dan varian rintang suhu V4-UPM, strain mesogenik S dan strain velogenik AF2240. Analisis aktiviti haemagglutinasii dan neuraminidase (iaitu fungsi protein HN) dan aktiviti hemolitik pada berlainan suhu menunjukkan strain AF2240, V4 dan V4-UPM rintang kepada suhu berbanding dengan strain S.

Tidak ada perbezaan yang dapat diperhatikan di dalam pergerakan protein NDV didalam kajian elektroforetik sodium dodesil sulfat-gel poliakrilamida (SDS-PAGE).



Walau bagaimanapun, analisis peptida menggunakan protease *Staphylococcus aureus* menunjukkan protein HN yang terhadam bagi strain V4-UPM berbeza daripada strain V4, AF2240 dan S. Analisis peptida ini telah diulang menggunakan *Pseudomonas fragi* Endoproteinase Lys-C dan *Lysobacter enzymogenes* Lys-C tetapi tidak menunjukkan sebarang perbezaan melainkan strain V4-UPM.

Protein-protein ini telah dianalisis selanjutnya menggunakan kaedah elektroforesis dua dimensi (2D-PAGE). Gel diproses seterusnya dengan kaedah Western blotting dan titik protein dikenal pasti menggunakan Mabs HN, NP dan F dan dianalisis dengan UVP GDS Gel Documentation System (UK). Didapati bahawa bagi strain V4 titik isoelektrik (pI) bagi protein HN berada pada bahagian asid, strain V4-UPM dan AF2240 dibahagian neutral/basik rendah dan pada strain labil suhu S, protein HN telah berubah ke bahagian basik pemfokusan isoelektrik. Perubahan pada titik pI bagi protein NP hanya dilihat pada strain S sahaja. Protein F adalah di dalam kawasan basik kecuali strain V4-UPM. Bagi strain S, didapati bahawa protein HN, NP dan F berada di kawasan basik dan kemungkinan ini menunjukkan perubahan ciri-ciri biologi yang dilihat bagi strain stabil-suhu dan labil-suhu.



CHAPTER 1

INTRODUCTION

Newcastle disease virus (NDV) is an economically important avian paramyxovirus which causes a highly contagious and fatal disease in poultry known as Newcastle disease (ND). Its genome is a negative-stranded RNA which encodes six major proteins : the nucleocapsid (NP) protein, the phosphoprotein (P), matrix (M) protein, fusion (F) protein, haemagglutinin-neuraminidase (HN) protein and the large (L) protein (Samson, 1988). Several non-structural proteins (36,000 dalton and 33,000 dalton) have also been detected in NDV-infected cells (Iinuma and Simpson, 1974; Moore and Burke, 1974).

The severity of the disease depends on several factors, namely, (i) biological properties of the viral strain, (ii) species and age of the host, (iii) presence of other organisms and (iv) environmental factors. NDV strains can be classified into three major pathotypes : (a) velogenic, with sudden death and very high mortality; (b) mesogenic, with respiratory signs but low mortality and (c) lentogenic, with mild



infections. In addition, some strains are avirulent and cause no disease at all.

Poultry has a unique role in the livestock sector of the countries in the Asia-Pacific region. It is the only livestock species that is widely accepted by people from a variety of cultural background. Although ND in commercial poultry is effectively controlled by vaccination, the use of conventional ND vaccination in village poultry has been an unsatisfactory process. The chickens which are of multi ages, are scattered over the villages and are difficult to catch for formal vaccination. Moreover, many live ND vaccines are heat-labile (Lomniczi, 1975) and are thus not suitable in the Tropics. A new approach is therefore required to control ND in the village poultry.

Through a cloning technique, an immunogenic and heat tolerant vaccine of NDV lentogenic strain V4 designated as V4-UPM was isolated (Ideris et al., 1989). The thermostabilities of infectivity and haemagglutinin of V4-UPM were found to be greater than those of the parent V4. It took at least 5 h for the HA (haemagglutinating activity) titre of strain V4-UPM to decrease by 2 logarithms (base 2) compared to only 2 h for strain V4. In the case of



thermostability of infectivity, the time required for a decrease in titre 2 logarithmic orders (base 10) was within 1 h for strain V4 and 3 h for V4-UPM. This new vaccine is used to coat feed pellets which are readily eaten by the chickens. It is currently being commercialised by a joint venture between Universiti Pertanian Malaysia and a private vaccine company Remeo Holdings Sdn. Bhd.

Very little work has been done on the thermostabilities of the NDV proteins. Studies on temperature sensitive (ts) mutants at the molecular level of the mesogenic Beaudette C strain have shown an altered isoelectric point in the P protein of mutant ts172 (Samson et al., 1981) and the HN and M proteins of mutant ts53 (Harper et al., 1983). Sequence analysis of the latter has identified sites in the HN protein which may be important in folding and function (Hughes et al., 1991). These studies suggests that the pathotype of the NDV strain may not be directly related to thermostability since the two temperature sensitive mutants (ts172 and ts53) and the heat stable mutant V4-UPM were from different pathotypes. However, these studies were done in separate experiments. Therefore, the thermostabilities of NDV strains representatives of the three major pathotypes will be determined to



confirm the hypothesis that there is no direct relationship between pathogenicity and thermostability.

Examination of the viral proteins in gel electrophoresis may reveal minute difference in the proteins of the various NDV strains. It is possible to correlate these differences to the thermostability.

This study was undertaken

- (1) to determine whether the NDV strains of varying pathotypes have different response to heat based on their biological activities;
- (2) to examine these differences at the molecular level using several types of gel electrophoretic techniques; and
- (3) to determine the proteins which may be directly involved in thermostability.

CHAPTER II

LITERATURE REVIEW

Newcastle Disease

Newcastle disease (ND) was first reported in 1926 in Jakarta, Indonesia, by Doyle and Kraneveld (Allan, 1971) but the origin and epidemiology of ND may be obscure (Hanson, 1978). In the following year it was recognised in other parts of Asia and later in Europe and America (Allan et al., 1973). In England the disease was centred in Newcastle-upon-Tyne hence the name of the disease (Spradbrow, 1987).

ND is one of the most important viral disease in the poultry industry. It causes severe economic losses due to death of chickens and lowered production of eggs, increased cost of purchasing vaccines and running of eradication and quarantine programmes during an outbreak (Lancaster, 1981; Spradbrow, 1987). In some countries vaccination programmes were adopted as the control measure with



varying successes (Biggs, 1982). The requirement of immunization also varies with each programme depending on the levels of protection needed, the immune status of the birds, the type of field virus occurring in each locality and the relationship between poultry diseases and the administration of the ND vaccines (Lancaster, 1964).

Newcastle disease virus (NDV) which is the causative agent of ND, is a member of the genus paramyxovirus within the family *Paramyxoviridae* which includes the mumps, Sendai and parainfluenza viruses (Compans and Choppin, 1967). Three distinct pathotypes of NDV have been described viz. lentogenic, mesogenic and velogenic (Hanson and Brandly, 1955) and on the basis of tissue tropism, they can be grouped as viscerotropic, pneumotropic and neurotropic (Allan et al., 1973).

The severity of infection is highly strain dependent (Alexander, 1988a). Velogenic strains cause severe disease with high mortality even in adult birds. Mesogenic ones are of moderate virulence, causing mortalities of up to 50% and seriously reducing egg production. The lentogenic strains are of low virulence causing little mortality except in young



chicks but affected egg production (Hanson, 1978; Spradbrow, 1987, Waterson et al., 1967).

The disease can spread either naturally or due to factors associated with the transportation of eggs, birds, carcasses, poultry offal, vaccinating crew and their movements or even by frozen poultry meat or contaminated vaccines (Hanson, 1978; Alexander, 1988a, 1988b).

Viral Structure

NDV is an enveloped, negative stranded RNA virus (Compans and Choppin, 1967). The RNA is complementary to the viral mRNAs and are generated during transcription in infected cells (Davies et al., 1976). The virion contains approximately 67% by weight of protein, 1% RNA, 24% lipid and 7% carbohydrate (Blough and Lawson, 1968; Haslam et al., 1969). Under negative electron-microscopy, the viral particles are pleomorphic in shape with a diameter of 100-500 nm (Hosaka et al., 1966; Kingsbury, 1974, 1990).

The envelope is covered with spikes of glycoprotein, the hemagglutinin-neuraminidase (HN) protein and the fusion (F) protein, of 8 to 12 nm long. The nonglycosylated matrix protein (M) forms a shell on

