COVID-19 and Pregnancy Outcomes: An Increased Risk of Intrauterine Inflammation/Infection

PHUONG NGUYEN ^(D) DINA EL-KADY ^(D) JONATHAN ROSNER ^(D) SALMA RAHIMI CHERYL DINGLAS ^(D)

*Author affiliations can be found in the back matter of this article

Scientific Innovation

COLLECTION: GME RESEARCH DAY 2021

PUBLISHED ABSTRACT

Levy Library Press

ABSTRACT

Objective: The objective of this study is to determine the association between COVID-19 infection and pregnancy outcomes at our institution when universal testing was implemented for all patients admitted to Labor & Delivery.

Methods: This was an IRB-exempt, retrospective chart review of all obstetrical patients admitted and evaluated in L&D from March 30th to April 30th. COVID-19 testing was performed on all patients who were admitted and their support person, irrespective of the presence of symptoms. Data analysis was performed with baseline demographics compared. Continuous variables were compared via T-test and categorical values using Chi-square and Fisher exact. Significant values are those considered with p < .05.

Results: There were no differences in delivery outcomes between the two groups with regards to mode of delivery, preterm labor, premature rupture of membranes, preeclampsia, placental abruption, or fetal demise. However, there was an increase in intrauterine infection/inflammation among COVID positive patients (8.8% compare to 1.4%, p < .05) (*Tables 1* and *2*).

Conclusions: COVID positive patients were noted to have an increase in intrauterine infection/inflammation. Current published data demonstrates that SARS-Cov-2 infection during 3rd trimester of pregnancy is not associated with vertical transmission. "However, the possibility of viral load influencing the transmission risk should be of concern. Published studies have demonstrated a positive relationship between the viral load of some viruses and their ability to spread from mother to child" [1]. Given our findings, the possibility of SARS-CoV-2 infection resulting in intrauterine infection/inflammation should be further evaluated as the pandemic continues.

CORRESPONDING AUTHOR: Phuong Nguyen Mt Sinai South Nassau, US LONOI92@GMAIL.COM

KEYWORDS: COVID-19; pregnancy; intrauterine infection and inflammation

TO CITE THIS ARTICLE: Nguyen P, El-Kady D, Rosner J, Pabimi S, Dinglas C, COVID-19

Rahimi S, Dinglas C. COVID-19 and Pregnancy Outcomes: An Increased Risk of Intrauterine Inflammation/Infection. *Journal of Scientific Innovation in Medicine*. 2021; 4(2): 18, pp. 1–3. DOI: https://doi. org/10.29024/jsim.113

brought to you by

Innovation in Medicine

			D \/A1115
CHARACTERISTICS	COVID NEGATIVE N = 141	COVID POSITIVE N = 34	P-VALUE
Age (years)	32.0 ± 7.6	29.6 ± 7.1	.05
BMI (kg/m²)	32.2 + 6.5	31.7 + 5.5	.34
Nulliparous n (%)	52 (36.9)	14 (41.1)	.64
Race/Ethnicity n (%)			.001
White	48 (34)	4 (11.8)	
Hispanic	40 (28.4)	22 (64.7)	
Black	20 (14.2)	3 (8.9)	
Asian	2 (1.4)	0	
Other	31 (21.9)	5 (15)	
Smoking	4(3)	0	.99
COVID positive partner	10 (7.1)	17 (50)	<.001
Medical comorbidities*			.32
None	116 (82.3)	32 (94.1)	
Pregestational DM	2 (1.4)	1 (2.9)	
Chronic hypertension	5 (3.5)	0	
Respiratory disease	6 (4.3)	2 (5.9)	
Autoimmune disease	7 (4.9)	0	
Other	7 (4.9)	0	
Medication use			.21
None (aside from vitamins)	113 (80.1)	30 (88.2)	
Steroids	0	0	
Aspirin	1 (0.7)	1 (2.9)	
Heparin/Lovenox	2 (1.4)	0	
Other	25 (17.7)	3 (8.8)	

Nguyen et al. Journal of Scientific Innovation in Medicine DOI: 10.29024/jsim.113

Table 1 Baseline demographics:Patients who tested negativevs. positive for SARS-CoV2.

Continuous variables are reflected as mean (SD). Categorical values are reflected as n (%).

^{*} n greater than 141, and greater than 34, respectively as 2 patients had 2 or more medical comorbidities.

OUTCOMES	COVID NEGATIVE	COVID POSITIVE	P-VALUE
Gestational age at delivery (weeks)	38.7 <u>+</u> 2.5	38.4 <u>+</u> 1.5	.24
Vaginal delivery	87 (61.7)	21 (61.8)	.99
Preterm labor	4 (2.8)	0	1.0
Preterm Premature rupture of membranes	7 (4.9)	1 (2.9)	.61
Placental abruption	0	0	1.0
Fetal demise	2 (1.4)	0	1.0
Preeclampsia	9 (6.4)	3 (8.8)	.62
Intrauterine Infection	2 (1.4)	3 (8.8)	.01
Placental Pathology			
Normal	35 (24.8)	17 (50)	0.49
Abnormal	5 (3.5)	4 (11.8)	

Table 2 Comparison ofoutcomes between COVIDnegative and COVID positivepatients.

Continuous variables are reflected as mean (SD). Categorical values are reflected as n (%).

COMPETING INTERESTS

The authors have no competing interests to declare.

AUTHOR AFFILIATIONS

Phuong Nguyen (b) orcid.org/0000-0002-0569-7340 Mt Sinai South Nassau, US

Dina El-Kady 🕩 orcid.org/0000-0002-5809-5888 Mt Sinai South Nassau, US

Jonathan Rosner () orcid.org/0000-0003-4998-2431 Mt Sinai South Nassau, US

Salma Rahimi Mt Sinai South Nassau, US

Cheryl Dinglas 🕩 orcid.org/0000-0002-3128-9586 Mt Sinai South Nassau, US

REFERENCE

1. Pan Y, Zhang D, Yang P, Poon LL, Wang Q. Viral load of SARS-CoV-2 in clinical samples. *Lancet Infect Dis.* 2020. DOI: https://doi.org/10.1016/S1473-3099(20)30113-4

Nguyen et al. Journal of Scientific Innovation in Medicine DOI: 10.29024/jsim.113

TO CITE THIS ARTICLE: Nguyen P, El-Kady D, Rosner J, Rahimi S, Dinglas C. COVID-19 and Pregnancy Outcomes: An Increased Risk of Intrauterine Inflammation/Infection. *Journal of Scientific Innovation in Medicine*. 2021; 4(2): 18, pp. 1–3. DOI: https://doi. org/10.29024/jsim.113

Submitted: 04 May 2021 Accepted: 04 May 2021 Published: 28 May 2021

COPYRIGHT:

© 2021 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See http://creativecommons.org/ licenses/by/4.0/.

Journal of Scientific Innovation in Medicine is a peer-reviewed open access journal published by Levy Library Press.

