



Analysis of Prognostic Risk Factors Determining Poor Functional Recovery After Comprehensive Rehabilitation Including Motor-Imagery Brain-Computer Interface Training in Stroke Patients: A Prospective Study

Qiong Wu^{1,2}, Yunxiang Ge³, Di Ma², Xue Pang², Yingyu Cao⁴, Xiaofei Zhang⁵, Yu Pan^{2*}, Tong Zhang^{1*} and Weibei Dou^{3*}

OPEN ACCESS

Edited by:

Giovanni Morone,
Santa Lucia Foundation (IRCCS), Italy

Reviewed by:

Marco Iosa,
Sapienza University of Rome, Italy
Luca Sebastianelli,
Hospital of Vipiteno, Italy

*Correspondence:

Yu Pan
pya00561@btch.edu.cn
Tong Zhang
ZT61611@sohu.com
Weibei Dou
douwb@tsinghua.edu.cn

Specialty section:

This article was submitted to
Neurorehabilitation,
a section of the journal
Frontiers in Neurology

Received: 31 January 2021

Accepted: 20 April 2021

Published: 10 June 2021

Citation:

Wu Q, Ge Y, Ma D, Pang X, Cao Y, Zhang X, Pan Y, Zhang T and Dou W (2021) Analysis of Prognostic Risk Factors Determining Poor Functional Recovery After Comprehensive Rehabilitation Including Motor-Imagery Brain-Computer Interface Training in Stroke Patients: A Prospective Study. *Front. Neurol.* 12:661816. doi: 10.3389/fneur.2021.661816

¹ School of Rehabilitation Medicine, China Rehabilitation Research Center, Capital Medical University, Beijing, China, ² Department of Rehabilitation Medicine, Beijing Tsinghua Changgung Hospital, School of Clinical Medicine, Tsinghua University, Beijing, China, ³ Department of Electronic Engineering, Beijing National Research Center for Information Science and Technology (BNRist), Tsinghua University, Beijing, China, ⁴ School of Mechanical Engineering, Beijing Institute of Petrochemical Technology, Beijing, China, ⁵ Department of Clinical Epidemiology and Biostatistics, Beijing Tsinghua Changgung Hospital, Tsinghua University, Beijing, China

Objective: Upper limb (UL) motor function recovery, especially distal function, is one of the main goals of stroke rehabilitation as this function is important to perform activities of daily living (ADL). The efficacy of the motor-imagery brain-computer interface (MI-BCI) has been demonstrated in patients with stroke. Most patients with stroke receive comprehensive rehabilitation, including MI-BCI and routine training. However, most aspects of MI-BCI training for patients with subacute stroke are based on routine training. Risk factors for inadequate distal UL functional recovery in these patients remain unclear; therefore, it is more realistic to explore the prognostic factors of this comprehensive treatment based on clinical practice. The present study aims to investigate the independent risk factors that might lead to inadequate distal UL functional recovery in patients with stroke after comprehensive rehabilitation including MI-BCI (CRIMI-BCI).

Methods: This prospective study recruited 82 patients with stroke who underwent CRIMI-BCI. Motor-imagery brain-computer interface training was performed for 60 min per day, 5 days per week for 4 weeks. The primary outcome was improvement of the wrist and hand dimensionality of Fugl-Meyer Assessment (δ FMA-WH). According to the improvement score, the patients were classified into the efficient group (EG, δ FMA-WH > 2) and the inefficient group (IG, δ FMA-WH \leq 2). Binary logistic regression was used to analyze clinical and demographic data, including aphasia, spasticity of the affected hand [assessed by Modified Ashworth Scale (MAS-H)], initial UL function, age, gender, time since stroke (TSS), lesion hemisphere, and lesion location.

Results: Seventy-three patients completed the study. After training, all patients showed significant improvement in FMA-UL ($Z = 7.381$, $p = 0.000^{**}$), FMA-SE ($Z = 7.336$, $p = 0.000^{**}$), and FMA-WH ($Z = 6.568$, $p = 0.000^{**}$). There were 35 patients (47.9%) in the IG group and 38 patients (52.1%) in the EG group. Multivariate analysis revealed that presence of aphasia [odds ratio (OR) 4.617, 95% confidence interval (CI) 1.435–14.860; $p < 0.05$], initial FMA-UL score ≤ 30 (OR 5.158, 95% CI 1.150–23.132; $p < 0.05$), and MAS-H \geq level I+ (OR 3.810, 95% CI 1.231–11.790; $p < 0.05$) were the risk factors for inadequate distal UL functional recovery in patients with stroke after CRIMI-BCI.

Conclusion: We concluded that CRIMI-BCI improved UL function in stroke patients with varying effectiveness. Inferior initial UL function, significant hand spasticity, and presence of aphasia were identified as independent risk factors for inadequate distal UL functional recovery in stroke patients after CRIMI-BCI.

Keywords: motor-imagery brain-computer interface, regression analysis, stroke, upper limb, rehabilitation

INTRODUCTION

Previous studies have reported that 85% of stroke survivors suffer from upper limb (UL) dysfunction, which has significant long-term effects on activities of daily living (ADL), leisure activities, and work (1). Upper limb motor function recovery, especially distal function, is one of the main goals of stroke rehabilitation. Only 12% of patients with stroke regain full UL function, while other patients require long-term care (2). Brain-computer interface (BCI) is an interactive system for internal and external environment that directly reflects brain activity (3). Motor-imagery brain-computer interface (MI-BCI) can quantify and reinforce feedback from motor imagery tasks and affect changes in neural network plasticity. For patients with stroke, MI-BCI training significantly improves UL motor function (4), electromyography signals (EEG) (5), joint mobility, daily living skills (6), mood (7), and brain network connectivity (8).

The effect of BCI is multidimensional, including the improvement of the clinical score and some subclinical indicators. Previous studies have shown significant improvement in UL function after MI-BCI training in patients with stroke (9, 10). Subclinical effects with improvement in the neuroelectrophysiological index have also been observed (11, 12). The mechanism of motor function recovery in the distal UL differs from that in the proximal UL. Most patients with stroke with inadequate overall UL functional recovery show predominantly improved proximal UL function, for instance, shoulder and elbow function, which results from non-decussated corticospinal fibers stemming from the unaffected hemisphere (13, 14). However, the distal UL function is markedly affected by the integrity of the corticospinal tract (CST) (12), which lacks a compensatory mechanism and therefore recovers inadequately. Motor function in the distal UL has a substantive effect on ADL (10, 15).

Most stroke patients receive comprehensive treatment that not only includes MI-BCI but also routine training. Multiple factors affect the prognosis of UL functional recovery. First, there is a large variation in user performance between healthy

individuals, and the possible factors include cognitive function, sensory, motor imagery ability (16), psychological factors (17), age (18), gender (19), dominant hand (20), sensorimotor rhythm bias (21), cortical gray matter volume, and medical treatment. Second, studies on the prognosis prediction of UL motor recovery in patients with stroke have incorporated age, gender, dominant hand, lesion of hemisphere and location, initial UL function, and presence of comorbidities into models that can predict UL recovery type up to 6 months after onset (22).

In studies on MI-BCI fitness, gender, age, event-related potential, classification accuracy, neuropsychological score (23), EEG laterality index, and cortical activation intensity (24) were found to be useful to predict the performance of stroke patients who could manipulate MI-BCI devices, and the symmetry of the EEG signal could be used as a predictor of the degree of UL improvement (25, 26). Subacute phase is an essential rehabilitation period for patients with stroke, and an increasing number of studies have included subacute stroke patients in BCI studies. However, the effect of these factors on the effectiveness of comprehensive treatment including MI-BCI in patients with stroke is unclear. The elucidation of this aspect will enable us to better define the indications for MI-BCI, guide the development of individualized rehabilitation programs, and improve treatment effectiveness in patients with stroke.

Therefore, the present study aimed to investigate the independent risk factors that might lead to inadequate distal UL recovery of patients with stroke after comprehensive rehabilitation including MI-BCI (CRIMI-BCI).

SUBJECTS AND METHODS

General Information

Eighty-two patients with ischemic stroke hospitalized at Tsinghua Changgung Hospital, Beijing, between January 2018 and December 2019 were recruited for the study. The diagnostic criteria were based on the “Consensus on Clinical Research Specifications for Acute Stroke in China 2018,” along with

differential diagnosis by magnetic resonance imaging (MRI) or computerized tomography (CT) (27). The inclusion and exclusion criteria were chosen according to a previous study (8).

Inclusion Criteria

The following inclusion criteria were used: (1) age 18–75 years; (2) sufficient cognition [Montreal Cognitive Scale (MOCA) score >20]; (3) a history of first-ever unilateral brain lesion confirmed by MRI; (4) stroke occurrence [time since stroke (TSS)] 1–6 months prior to inclusion; (5) moderate to severe UL paralysis (Brunnstrom stages $\leq IV$); and (6) right-handedness (Edinburgh Handedness Inventory score ≥ 40).

Exclusion Criteria

The following exclusion criteria were used: (1) severe spasticity of the affected hand [Modified Ashworth Scale (MAS) score ≥ 3]; (2) open wound or deformity of the affected UL; (3) visual field deficit or unilateral spatial neglect; (4) severe aphasia [Boston Diagnostic Aphasia Examination (BDAE) score <3]; (5) currently undergoing antipsychotic treatment; (6) severe

dystonia and/or involuntary movements; (7) other severe neurological disorders such as epilepsy; and (8) currently undergoing neuromodulation treatment.

Nine patients were excluded at follow-ups conducted during the study due to changes in their conditions. Complete data were collected from 73 patients. The experimental flow chart is shown in **Figure 1**.

The study was approved by the Ethics Committee of Beijing Tsinghua Changgung Hospital (No. 18172-0-02) and registered at <http://www.chictr.org.cn> (No. ChiCTR1900022128). All patients signed an informed consent form prior to the trial. The following data were obtained from patients with stroke: gender, age, TSS, lesion hemisphere and location, and presence of concomitant aphasia. The lesion location was staged according to the Oxfordshire Community Stroke Study (OCPS) (28, 29). Aphasia diagnostic criteria were determined by the Aphasia Battery of Chinese (ABC), aphasia diagnostic criteria set as total score <25 (30).

General information of the 73 included patients is shown in **Table 1**.




TABLE 1 | General information of 73 patients.

	Mean/Median	Classification	Number	n (%)
Age (year)	61.00 (46.00, 67.00)	<65	50	68.49
		≥65	23	31.51
TSS (month)	3.00 (2.00, 5.00)	≤6	60	82.19
		>6	13	17.81
Lesion		TACI	29	39.73
		PACI	44	60.27
Gender		Female	25	34.25
		Male	48	65.75
Aphasia		With	49	67.12
		Without	24	32.88
Affected hemisphere		Right	31	42.47
		Left	42	57.53

PACI, Partial anterior circulation infarct; TACI, Total anterior circulation infarct.

EEG Acquisition

Scalp EEG potentials were collected from eight dry electrodes (according to 10–20 International System), band-passed by 2–60 Hz filter and a notch filter (48–52 Hz) to remove artifacts and power line interference, respectively; digitized at 256 Hz, and amplified by a commercial EEG system (g. LADYbird, g. Tec Medical Engineering GmbH, Schiedlberg, Austria) and then processed by a computer.

EEG signals were grounded to a unilateral earlobe and referenced at the other one. Electrodes were positioned over FC3, FC4, C3, C4, CP3, CP4, C1, and C2. Signals from the C3 and C4 were used for device control. Motor function related electrodes were used for offline analyses (left hemisphere: FC3, C3, and CP3; right hemisphere: FC4, C4, and CP4). Mu suppression, which reflects Event-Related Desynchronization (ERD), was due to increased neural activity. The mu suppression score provides information about motor innervation, and was posted on the screen, encouraged patients to get higher scores. For mu suppression score computation, EEG data from C3 to C4 were transformed into the frequency domain by a Fourier transform algorithm with a Hanning window covering the EEG data during the video period of the paradigm. The equation of mu suppression score are as follows (8):

$$\text{Musupp} = -\frac{\text{Mup}_{\text{task}} - \text{Mup}_{\text{rest}}}{\text{Mup}_{\text{rest}}} \times 100$$

Musupp: Mu suppression score, Mup_{test} : EEG power during MI; Mup_{rest} : EEG power during the resting state.

Comprehensive Rehabilitation Training Program

All patients received standard treatment for stroke in terms of medical care and rehabilitation, which consisted of conventional treatment, including an intensive occupational therapy focused on activities of daily live, such as grasping a toothpaste tube, eating, and reaching. A conventional treatment session lasted for 1 h per day, 5 days per week, for 4 weeks. MI-BCI training was

applied based on conventional treatment. The detailed protocol refers to previous studies (8).

MI-BCI Training Paradigms

To facilitate MI performance, patients were given the opportunity to execute tasks with the affected and unaffected hand several times before MI-BCI training. Meanwhile they were instructed to perform only MI tasks and to avoid movement attempts of the affected UL.

MI-BCI training consisted of 20 sessions, five sessions per week, for 4 weeks. During each session, the patient was comfortably seated in a soundproofed room, with their affected hand resting in an exoskeleton hand. A video of the unaffected hand grasping/opening was presented on a screen in front of the patient to guide the MI task. The exoskeleton hand provided mechanical support and assistance to the affected hand based on the mu suppression algorithm calculated during the follow up trial.

- (1) Movement observation: a dark screen was first displayed for 2 s, followed by a white cross for 2 s. A vocal cue of “hand grasp” or “hand open” was displayed for 2 s. Then a video clip was displayed for a duration of 6 s. Patients were requested to observe the video and avoid blinking, coughing, chewing, and performing head movements.
- (2) Exoskeleton hand assistance: If the mu suppression score was above 20, the exoskeleton hand would assist the hand grasping/opening task during the following 3 s.
- (3) End of process: the mu suppression score was then shown for 2 s. The trial ended with the display of a dark screen for 2 s. During each session, the trial was repeated 100 times for one session, and video clips of the grasping and opening hand were shown randomly. Patients were permitted to rest for 1 min after 10 trials.

Functional Evaluation

Before and after 20 training sessions, the motor function of the affected UL and muscle tone of the affected finger flexor were

evaluated. A simplified version of the Fugl-Meyer Assessment UL (FMA-UL), which was without sensory, passive joint mobility, and pain and reflex sections, was used to assess the motor function of the affected UL (31). The total FMA-UL score was 60, which was obtained by summing the five items on the FMA-UL scale: shoulder, elbow, wrist, hand, and coordination/speed. The sum of the wrist and hand score (FMA-WH) was 24. The sum of the shoulder and elbow score (FMA-SE) was 30. A higher score for each dimension implied that the patient was more functional. The simplified FMA-UL score was categorized as severe (0–12), severe-moderate (13–30), moderate-mild (31–47), and mild (48–60) (2). Because the initial FMA-UL scores of most patients were between severe-moderate to moderate-mild, a score of 30 was selected as the watershed in the following regression analysis.

The MAS was used to measure the muscle tone of the affected finger flexors and recorded as MAS-H (32, 33). For statistical convenience, the scores were recorded as follows: level 0 was set as 1, level I–I+ as 2, level II as 3, and level III–IV as 4. Because the initial MAS-H scores of most patients were between 2 and 3, level I+ was selected as the watershed.

All assessments were performed by one trained therapist, and the assessor was unaware of the therapeutic condition of the patients. Functional evaluation of the 73 patients is shown in **Table 2**.

Comparison of Group Characteristics

The 73 patients who completed all training and testing procedures were divided into two groups according to whether the increase in the FMA-WH score (δ FMA-WH) was higher than two points before and after treatment. The differences between the two groups were compared in terms of gender, age, TSS, lesion hemisphere and location, presence of aphasia, MAS-H, and FMA-UL and its component score.

Statistical Analysis

Data were analyzed using SPSS 22.0 statistical software, and measured values were expressed as mean and standard deviation ($x \pm s$) or median and interquartile range (25th percentile, 75th percentile). Data were verified for normality of distributions with the Kolmogorov-Smirnov test ($n > 50$) or Shapiro-Wilk test ($n < 50$). Because of the ordinal nature of the data, the study data were compared between groups using independent sample *t*-test and Mann-Whitney U test. Data were compared within the group using an independent *t*-test and a paired-sample Wilcoxon test. Categorical variables

were expressed as percentages and analyzed with the chi-square test or Fisher's exact test, and significance was set at $p < 0.05$. For regression analysis, the δ FMA-WH score was used as the criterion for grouping. Binary logistic regression was performed on the basis of the results of univariate analysis for $p < 0.10$ and by incorporating the parameters into the model through the Forward LR method. The results of multivariate analysis were expressed as odds ratio (OR) and 95% confidence interval (CI). A two-sided $p < 0.05$ was considered to be statistically significant.

RESULTS

Overall Efficacy of Comprehensive Rehabilitation

After comprehensive rehabilitation, 73 patients showed significant improvement in FMA-UL, FMA-SE, and FMA-WH (Wilcoxon signed-rank test, $p < 0.000$), as shown in **Table 2** and **Figure 2**.

Differences in Functional Recovery

It was difficult to perceive changes in hand function when δ FMA-WH ≤ 2 points (33) and by referring to the definition of minimum clinically important difference (MCID). Hence, δ FMA-WH ≤ 2 points before and after training was selected as the criterion for judging the efficacy (34, 35). The 73 patients were divided into the efficacy group (EG, with δ FMA-WH > 2 points, $n = 38$, 52.1%) and the inefficacy group (IG, δ FMA-WH ≤ 2 points, $n = 35$, 47.9%).

The difference in UL motor function status was compared between the two groups. The results showed no significant differences in the FMA-UL pre (Mann-Whitney U test, $p = 0.584$), FMA-SE pre (Mann-Whitney U test, $p = 0.615$), and FMA-WH pre (Mann-Whitney U test, $p = 0.950$) between the groups before training.

After training, the total score and the WH score in both EG and IG groups were significantly higher than those before the training [FMA-UL post (Mann-Whitney U test, $p = 0.026$) and FMA-WH post (Mann-Whitney U test, $p = 0.001$)]. The improvement in the total, SE, and WH scores of FMA-UL in the EG group was also significantly higher than that in the IG group [δ FMA-UL (Mann-Whitney U test, $p = 0.000$), δ FMA-WH (Mann-Whitney U test, $p = 0.000$), and δ FMA-SE (independent samples *t*-test, $p = 0.000$)], as listed in **Table 3** and **Figure 3**.

Risk Factor of Unfavorable Recovery

The univariate analysis revealed that TSS (chi-square test, $p = 0.040$), MAS-H (chi-square test, $p = 0.007$), presence of aphasia (chi-square test, $p = 0.025$), and FMA-UL pre (chi-square test, $p = 0.05$) might be associated with unfavorable hand function recovery after BCI training in patients with stroke. Age (chi-square test, $p = 0.989$), lesion (chi-square test, $p = 0.205$), gender (chi-square test, $p = 0.626$), and affected hemiplegia (chi-square test, $p = 0.377$) were not found to be relevant for the functional

TABLE 2 | UL function in 73 patients before and after training.

	Pre	Post	Z	p
FMA-UL	18 (8, 27.5)	30 (16, 45.5)	7.381	0.000**
FMA-SE	13 (8–19)	20 (12, 28.5)	7.336	0.000**
FMA-WH	3 (0, 8)	9 (4–16)	6.568	0.000**

** $p < 0.01$.

change of the distal UL after training in patients, as listed in **Table 4** and **Figure 4**.

Binary logistic regression analysis was performed based on the results of univariate analysis for $p < 0.10$. Presence of aphasia [odds ratio (OR) 4.617, 95% confidence interval (CI) 1.435–14.860; $p = 0.010$], FMA-UL pre ≤ 30 (OR 5.158, 95% CI 1.150–23.132; $p < 0.032$), and MAS-H \geq level I+ (OR 3.810, 95% CI 1.231–11.790; $p = 0.020$) were found to be risk factors for inadequate distal UL recovery in patients with stroke after CRIMI-BCI training, as shown in **Table 5**.

DISCUSSION

The results of the present study showed that for the target population, CRIMI-BCI significantly improved the overall and

distal motor function of the UL in stroke patients, but the degree of improvement varied. Classification according to distal functional improvement showed that although no significant differences were observed in UL motor function before training, the proportion of patients with aphasia, TSS > 6 months, and finger flexor tone >1+ grade on the affected hand was significantly higher in the IG group. Regression analysis showed that significant hand spasticity, poor initial function, and concomitant aphasia were the important factors that influenced the inadequate distal UL motor function after CRIMI-BCI in this study.

The results of the present study showed that CRIMI-BCI improved UL function, including distal motor function, in patients with stroke. This result is consistent with many studies (6, 25, 35). Bundy DT found significant improvements in grip




TABLE 3 | Comparison of changes before and after training between groups.

	EG (n = 38)	IG (n = 35)	t/Z	p
FMA-UL pre	18,000 (12.0, 24.0)	12,000 (6.0, 40.0)	-0.548	0.584
FMA-SE pre	13,000 (10.0, 18.0)	10,000 (6.0, 24.0)	-0.503	0.615
FMA-WH pre	4,000 (0.0, 7.0)	2,000 (0.0, 13.0)	-0.063	0.950
FMA-UL post	32,000 (23.0, 47.3)	18,000 (8.0, 44.0)	-2.226	0.026*
FMA-SE post	21,000 (16.0, 28.3)	17,000 (8.0, 29.0)	-1.261	0.207
FMA-WH post	12,000 (7.8, 16.3)	4,000 (0.0, 13.0)	-3.406	0.001**
ΔFMA-UL	14,000 (9.8, 18.3)	4,000 (3.0, 5.0)	-6.264	0.000**
ΔFMA-SE	17.42 ± 7.06	13.09 ± 6.24	2.770#	0.007**
ΔFMA-WH	7,000 (4.0, 11.0)	1,000 (0.0, 2.0)	-7.388	0.000**

* $p < 0.05$, ** $p < 0.01$, #Mann-Whitney U test.





FIGURE 3 | (A–C) UL function improvement in groups. Although the overall FMA-UL score improved, the FMA-WH scores in two patients regressed after training in IG. * $p < 0.05$, ** $p < 0.01$.

TABLE 4 | Comparison of the characteristics between groups.

	Classification	EG (%)	IG (n %)	Subtotal (n %)	χ^2	<i>p</i>
Age (year)	≤65	26 (68.42)	24 (68.57)	50 (68.49)	0	0.989
	>65	12 (31.58)	11 (31.43)	23 (31.51)		
TSS (month)	≤6	35 (92.11)	26 (74.29)	61 (83.56)	4.212	0.040*
	>6	3 (7.89)	9 (25.71)	12 (16.44)		
MAS-H	≤1+	28 (73.68)	15 (42.86)	43 (58.90)	7.152	0.007**
	>1+	10 (26.32)	20 (57.14)	30 (41.10)		
Lesion	TACI	15 (39.47)	19 (54.29)	34 (46.58)	1.606	0.205
	PACI	23 (60.53)	16 (45.71)	39 (53.42)		
Gender	Female	14 (36.84)	11 (31.43)	25 (34.25)	0.237	0.626
	Male	24 (63.16)	24 (68.57)	48 (65.75)		
Aphasia	Without	30 (78.95)	19 (54.29)	49 (67.12)	5.021	0.025*
	With	8 (21.05)	16 (45.71)	24 (32.88)		
Affected hemisphere	Left	18 (47.37)	13 (37.14)	31 (42.47)	0.780	0.377
	Right	20 (52.63)	22 (62.86)	42 (57.53)		
FMA-UL pre	≤30	34 (89.47)	25 (71.43)	59 (80.82)	3.827	0.050
	>30	4 (10.53)	10 (28.57)	14 (19.18)		

p* < 0.05, *p* < 0.01.



strength, grasp, and pinch function in stroke patients with chronic moderate to severe UL paresis, after using BCI combined with exoskeleton training in the unaffected hemisphere (35). Frolov et al. showed that in patients with stroke, BCI combined with exoskeleton training showed a significant advantage in the

grip and pinch function of the affected finger as compared to exoskeleton training alone (25).

Previous studies on factors influencing the clinical outcome of MI-BCI have shown no correlation between spasticity and training outcome in patients with stroke. However, the effect

TABLE 5 | Binary logistic regression of risk factors.

	B	SE	Z-value	Wald χ^2	P	OR	OR (95%CI)
MAS-H	1.338	0.576	2.321	5.386	0.020*	3.810	1.231–11.790
FMA-UL	1.641	0.766	2.143	4.591	0.032*	5.158	1.150–23.132
TSS	0.515	0.812	0.635	0.403	0.526	1.674	0.341–8.222
Aphasia	1.530	0.596	2.565	6.579	0.010*	4.617	1.435–14.860

B, estimate coefficient; SE, standard error; OR, odds ratio; CI, confidence interval. * $p < 0.05$.

of spasticity on UL motor function was demonstrated by numerous studies on the prediction of motor function in stroke patients. The present study showed that the MAS-H score was an independent factor that influenced inadequate functional improvement of the distal UL in patients after CRIMI-BCI. Finger and wrist flexor spasticity is very common in patients with stroke, and the recovery of wrist and hand movements vary greatly among patients with different degrees of spasticity. Patients with severe spasticity have minimal improvement in the FMA of the hand (34, 36). In contrast, patients with moderate and mild spasticity exhibit greater hand motion recovery. Hand spasticity affects the recovery of random finger movements as well as grasp and release functions. Spasticity can also impede motor learning ability after stroke (37). The severity of distal spasticity directly reflects the severity of the CST injury, as there is no compensation of the ipsilateral motor conduction. Spasticity of the hand is an extremely critical influencing factor in the recovery of motor function after stroke (36). The understanding of the relationship between hand spasticity and the effect of MI-BCI training in patients with stroke can be helpful in designing individualized rehabilitation programs.

Previous studies have suggested that complications such as depression, cognitive impairment, aphasia, hemianopia, and unilateral neglect after stroke also affect the recovery of motor function (38). The presence of spontaneous speech is a favorable factor for the recovery of motor function (39, 40). The present study showed that concomitant aphasia was an independent influencing factor of modest functional improvement of the distal UL in the patients. A possible reason for this was that although patients with concomitant aphasia in this study had mild speech impairment, the training process involved more complex instructions such as hearing vocal signals—watching videos—closing eyes—completing motor imagery—opening eyes to prepare for the next cycle; hence, speech dysfunction not only affected the reception and expression of information by the patients, but it may also have involved factors such as attention and working memory, resulting in inefficiency training and inadequate rehabilitation. Further research is necessary to understand the effect of aphasia on the effectiveness of MI-BCI in terms of the type, severity, and mechanism of the disorder.

Many studies have confirmed that the performance of MI-BCI manipulation does not correlate with motor function. However, in most stroke prediction models, the initial UL function score significantly affects prognosis. The patients in the present study had no significant differences in FMA-UL pre as observed in

Mann-Whitney U test; all of these patients had moderate or severe impairment. However, the difference in data dispersion between the groups can be observed from the box plot. There was an inter group difference in the trend of the initial FMA-UL degree (classified by 30 points), and a further increase in sample size might have cleared the difference. We incorporated the FMA-UL pre-score into the regression analysis based on clinical experience and prognostic studies. However, the initial motor function still showed a significant effect on treatment outcome in the regression analysis; this finding is consistent with the general pattern of UL functional recovery in patients with stroke. The result may be related to the initial FMA-UL score distribution or the interaction between the variables.

It is worth noting that, unlike previous studies, the present study did not show any influence of TSS, age, gender, and lesion on the effect of CRIMI-BCI. This may be explained by the distinctive population scope of the study. For example, in the intragroup comparison, the number of cases with TSS > 6 months was significantly greater in the IG group than in the EG group. However, in the multivariate analysis, TSS did not constitute an independent risk factor affecting the unfavorable functional recovery of the distal UL in stroke patients. Age has been reported to influence the performance of MI-BCI in healthy individuals, with older patients showing reduced EEG power, laterality, and significantly lower discrimination accuracy than younger patients (40, 41). Advanced age is also an influential factor in the inadequate prognosis of UL function after stroke (22, 42, 43). Regarding the effect of gender on outcome, many studies on the prognosis of stroke have shown that male patients have a better prognosis than female patients. However, Randolph reported that male patients had poorer ability to regulate μ -rhythms in the EEG and poorer control of the MI-BCI system than female patients (19). In terms of the location of lesions, the anterior putamen, internal capsule, thalamus, periventricular white matter, and premotor cortex were associated with inadequate UL recovery in patients with hemorrhagic stroke (44). Studies on chronic ischemic stroke have shown poorer recovery in patients with internal capsule injury (45). The prognosis of the TACI was worse than that of the other types (46, 47). In the present study, most patients with complete anterior circulation and severe posterior circulation could not complete MI-BCI training because of other comorbidities. In contrast, there is a POCI type in OCPS, but no patient with this type was not recruited in this study. This might be because such types of patients generally face less

obstacles and therefore hardly meet the test conditions. This may be related to the generally mild motor dysfunction of these patients. Therefore, the results of the present study were limited to the current population for whom MI-BCI is indicated, and the duration of the disease was restricted to the subacute phase. Additionally, cognitive impairment, severe aphasia, severe spasticity, depression, hemianopia, unilateral neglect, and other comorbidities were excluded, and the effect of the dominant hand was limited. The patient population in the present study was more homogeneous than that observed in actual clinical patients, and no remarkable differences were observed in TSS, age, gender, and lesion. Hence, the effect of the training might not be evident. This suggested that the study of factors influencing the effect of CRIMI-BCI on patients with stroke was very complicated, and the findings need to be tailored to its scope of application. Moreover, there is an urgent need to combine neuroimaging and neurophysiological investigations in the next step of this study.

The results of the present study answered the initial queries and showed the following findings: (1) CRIMI-BCI can improve UL motor function, especially the distal function, in patients with stroke in the subacute stage; (2) a heterogeneity was observed in the effect of CRIMI-BCI on distal UL function, with significantly better overall and distal UL function improvement in EG patients than in IG patients; and (3) in stroke patients with subacute phase UL paralysis who underwent CRIMI-BCI, poor initial UL function, severe hand spasticity, and concomitant aphasia were the influencing factors of inadequate distal function recovery.

LIMITATIONS

The generalization of these results is subject to certain limitations. First, although previous studies have compared the effects of conventional therapy with BCI training on patients with subacute stroke, the present study did not include a control group, and therefore, it cannot be determined whether the abovementioned findings differ from those of patients with conventional stroke. Second, the sample size was insufficient and limited by the strict screening requirement of the patients and the long-term training duration. Moreover, there was inadequate inclusion of factors influencing clinical outcomes, including NIHSS score and some early clinical predictors such as the presence of shoulder shrugging, active finger extension, and grip strength of the affected hand. Third, indicators reflecting the ability to manipulate MI-BCI, including classification rate and EEG activity, were lacking. Fourth, neuroimaging-related evidence was lacking. In addition, long-term and interaction effects were not analyzed. Therefore, controlled, long-term studies with larger sample sizes, more factors, and neuroimaging and neurophysiological evidence are needed to further validate the results of the present study.

CONCLUSION

The present study aimed to investigate the factors influencing the effect of CRIMI-BCI on the functional recovery of the distal UL

in patients with stroke. The main results suggest that CRIMI-BCI improved overall and distal UL function in patients with subacute stroke, but that the outcomes were varied. Poor initial UL function, severe hand spasticity, and concomitant aphasia were independent risk factors for inadequate MI-BCI training outcomes. This study was restricted to a specific population with subacute stroke who underwent CRIMI-BCI. It clarified the effect of BCI on distal UL function and explored further indications, prognosis, and protocols for rehabilitation. The study findings can be applied to design individualized rehabilitation programs for patients with stroke.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository and accession number can be found below: Chinese Clinical Trial Registry (ChiCTR), <http://www.chictr.org.cn>, No. ChiCTR1900022128.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by Ethics Committee of Beijing Tsinghua Changgung Hospital. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

QW, YP, WD, and TZ designed the study. DM, XP, and YC performed experiments. QW and YG wrote the paper. All authors contributed to the article and approved the submitted version.

FUNDING

This study was supported by the Beijing Hospitals Authority Youth Programme (QML20200902), the Opto-Mechatronic Equipment Technology Beijing Area Major Laboratory, the Beijing Institute of Petrochemical Technology (BTPT-OMET-OF-2020-1), the Beijing Municipal Natural Science Foundation (L182028), and the Beijing Municipal Science and Technology Commission (Z181100009218003). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

ACKNOWLEDGMENTS

We would like to like to acknowledge all the patients for participating in this experiment. We thank International Science Editing (<http://www.internationalscienceediting.com>) for editing this manuscript.

REFERENCES

- Aparicio HJ, Himali JJ, Satizabal CL, Pase MP, Romero JR, Kase CS, et al. Temporal trends in ischemic stroke incidence in younger adults in the Framingham study. *Stroke*. (2019) 50:1558–60. doi: 10.1161/STROKEAHA.119.025171
- Woytowicz EJ, Rietschel JC, Goodman RN, Conroy SS, Sorokin JD, Whittall J, et al. Determining levels of upper extremity movement impairment by applying a cluster analysis to the Fugl-Meyer assessment of the upper extremity in chronic stroke. *Arch Phys Med Rehabil*. (2017) 98:456–62. doi: 10.1016/j.apmr.2016.06.023
- Carino-Escobar RI, Carrillo-Mora P, Valdés-Cristerna R, Rodríguez-Barragan MA, Hernández-Arenas C, Quinzaños-Fresnedo J, et al. Longitudinal analysis of stroke patients' brain rhythms during an intervention with a brain-computer interface. *Neural Plast*. (2019) 2019:1–11. doi: 10.1155/2019/7084618
- Várkuti B, Guan C, Pan Y, Phua KS, Ang KK, Kuah CW, et al. Resting state changes in functional connectivity correlate with movement recovery for BCI and robot-assisted upper-extremity training after stroke. *Neurorehabil Neural Repair*. (2013) 27:53–62. doi: 10.1177/1545968312445910
- Ramos-Murguialday A, Broetz D, Rea M, Lärer L, Yilmaz O, Brasil FL, et al. Brain-machine interface in chronic stroke rehabilitation: a controlled study. *Ann Neurol*. (2013) 74:100–8. doi: 10.1002/ana.23879
- Young BM, Nigogosyan Z, Remsik A, Walton LM, Song J, Nair VA, et al. Changes in functional connectivity correlate with behavioral gains in stroke patients after therapy using a brain-computer interface device. *Front Neuroeng*. (2014) 7:25. doi: 10.3389/fneng.2014.00025
- Chowdhury A, Meena YK, Raza H, Bhushan B, Uttam AK, Pandey N, et al. Active physical practice followed by mental practice using BCI-driven hand exoskeleton: a pilot trial for clinical effectiveness and usability. *IEEE J Biomed Health Inform*. (2018) 22:1786–95. doi: 10.1109/JBHI.2018.2863212
- Wu Q, Yue Z, Ge Y, Ma D, Yin H, Zhao H, et al. Brain functional networks study of subacute stroke patients with upper limb dysfunction after comprehensive rehabilitation including BCI training. *Front Neurol*. (2019) 10:1419. doi: 10.3389/fneur.2019.01419
- Biasiucci A, Leeb R, Iturrate I, Perdakis S, Al-Khodairy A, Corbet T, et al. Brain-actuated functional electrical stimulation elicits lasting arm motor recovery after stroke. *Nat Commun*. (2018) 9:2421. doi: 10.1038/s41467-018-04673-z
- Daly JJ, Cheng R, Rogers J, Litinas K, Hrovat K, Dohring M. Feasibility of a new application of noninvasive Brain Computer Interface (BCI): a case study of training for recovery of volitional motor control after stroke. *J Neurol Phys Ther*. (2009) 33:203–11. doi: 10.1097/NPT.0b013e3181c1fc0b
- Cervera MA, Soekadar SR, Ushiba J, Millán J, Liu M, Birbaumer N, et al. Brain-computer interfaces for post-stroke motor rehabilitation: a meta-analysis. *Ann Clin Transl Neurol*. (2018) 5:651–63. doi: 10.1002/acn3.544
- Senesh MR, Reinkensmeyer DJ. Breaking proportional recovery after stroke. *Neurorehabil Neural Repair*. (2019) 33:888–901. doi: 10.1177/1545968319868718
- Curado MR, Cossio EG, Broetz D, Agostini M, Cho W, Brasil FL, et al. Residual upper arm motor function primes innervation of paretic forearm muscles in chronic stroke after brain-machine interface (BMI) training. *PLoS ONE*. (2015) 10:e0140161. doi: 10.1371/journal.pone.0140161
- Mrachacz-Kersting N, Jiang N, Stevenson AJ, Niazi IK, Kostic V, Pavlovic A, et al. Efficient neuroplasticity induction in chronic stroke patients by an associative brain-computer interface. *J Neurophysiol*. (2016) 115:1410–21. doi: 10.1152/jn.00918.2015
- Oujamaa L, Relave I, Froger J, Mottet D, Pelissier JY. Rehabilitation of arm function after stroke. Literature review. *Ann Phys Rehabil Med*. (2009) 52:269–93. doi: 10.1016/j.rehab.2008.10.003
- Pitt KM, Brumberg JS. Guidelines for feature matching assessment of brain-computer interfaces for augmentative and alternative communication. *Am J Speech Lang Pathol*. (2018) 27:950–64. doi: 10.1044/2018_AJSLP-17-0135
- Kadosh KC, Staunton G. A systematic review of the psychological factors that influence neurofeedback learning outcomes. *Neuroimage*. (2019) 185:545–55. doi: 10.1016/j.neuroimage.2018.10.021
- Zich C, Debener S, De Vos M, Frerichs S, Maurer S, Kranczioch C. Lateralization patterns of covert but not overt movements change with age: an EEG neurofeedback study. *Neuroimage*. (2015) 116:80–91. doi: 10.1016/j.neuroimage.2015.05.009
- Randolph AB. Not all created equal: individual-technology fit of brain-computer interfaces. In: *45th Hawaii International Conference on System Sciences*. Hawaii: IEEE (2012), p. 572–8. doi: 10.1109/HICSS.2012.451
- Jeunet C, Lotte F, Batail JM, Philip P, Micoulaud Franchi JA. Using recent BCI literature to deepen our understanding of clinical neurofeedback: a short review. *Neuroscience*. (2018) 378:225–33. doi: 10.1016/j.neuroscience.2018.03.013
- Zapala D, Zabielska-Mendyk E, Augustynowicz P, Cudo A, Jaśkiewicz M, Szewczyk M, et al. The effects of handedness on sensorimotor rhythm desynchronization and motor-imagery BCI control. *Sci Rep*. (2020) 10:2087. doi: 10.1038/s41598-020-59222-w
- van der Vliet R, Selles RW, Andrinopoulou ER, Nijland R, Ribbers GM, Frens MA, et al. Predicting upper limb motor impairment recovery after stroke: a mixture model. *Ann Neurol*. (2020) 87:383–93. doi: 10.1002/ana.25679
- Lyukmanov RK, Aziatskaya GA, Mokienco OA, Varako NA, Kovyazina MS, Suponeva NA, et al. Post-stroke rehabilitation training with a brain-computer interface: a clinical and neuropsychological study. *Zh Nevrol Psikhiatr Im S Korsakova*. (2018) 118:43–51. doi: 10.17116/jnevro201811808143
- Shu X, Chen S, Yao L, Sheng X, Zhang D, Jiang N, et al. Fast recognition of BCI-inefficient users using physiological features from EEG signals: a screening study of stroke patients. *Front Neurosci*. (2018) 12:93. doi: 10.3389/fnins.2018.00093
- Frolov AA, Mokienco O, Lyukmanov R, Biryukova E, Kotov S, Turbina L, et al. Post-stroke Rehabilitation training with a motor-imagery-based brain-computer interface (BCI)-controlled hand exoskeleton: a randomized controlled multicenter trial. *Front Neurosci*. (2017) 11:400. doi: 10.3389/fnins.2017.00400
- Mane R, Chew E, Phua KS, Ang KK, Robinson N, Vinod AP, et al. Prognostic and monitoring EEG-biomarkers for BCI upper-limb stroke rehabilitation. *IEEE Trans Neural Syst Rehabil Eng*. (2019) 27:1654–64. doi: 10.1109/TNSRE.2019.2924742
- Cerebrovascular Group of Neurology Branch of Chinese Medical Association. Chinese consensus for clinical study standards of acute stroke 2018. *China J Neurol*. (2018) 51:247–55. doi: 10.3760/cma.j.issn.1006-7876.2018.04.003
- Sung SF, Chen SC, Lin HJ, Chen CH, Tseng MC, Wu CS, et al. Oxfordshire Community Stroke Project classification improves prediction of post-thrombolysis symptomatic intracerebral hemorrhage. *BMC Neurol*. (2014) 14:39. doi: 10.1186/1471-2377-14-39
- Yang Y, Wang A, Zhao X, Wang C, Liu L, Zheng H, et al. The Oxfordshire Community Stroke Project classification system predicts clinical outcomes following intravenous thrombolysis: a prospective cohort study. *Ther Clin Risk Manag*. (2016) 12:1049–56. doi: 10.2147/TCRM.S107053
- Zheng Y, Zhong D, Huang Y, He M, Xiao Q, Jin R, et al. Effectiveness and safety of repetitive transcranial magnetic stimulation (rTMS) on aphasia in cerebrovascular accident patients: protocol of a systematic review and meta-analysis. *Medicine (Baltimore)*. (2019) 98:e18561. doi: 10.1097/MD.00000000000018561
- Liu L, Zhu L, Hou JX, Li XY, Yang F, Li SL, et al. Preliminary application analysis of Broetz hand test in patients with upper limb and hand dysfunction in acute stage of stroke. *Chin J Cerebrovasc Dis*. (2019) 16 (02):20–23; 29. doi: 10.3969/j.issn.1672-5921.2019.02.003
- Lu HT, Cui LH, Wang YZ. Comparison of effects of botulinum toxin type A injection under different guidance approaches on wrist and finger spasticity in stroke patients. *Chin J Rehabil Theory Pract*. (2019) 25:1352–5. doi: 10.3969/j.issn.1006-9771.2019.11.017
- Plantin J, Pennati GV, Roca P, Baron JC, Laurencikas E, Weber K, et al. Quantitative assessment of hand spasticity after stroke: imaging correlates and impact on motor recovery. *Front Neurol*. (2019) 10:836. doi: 10.3389/fneur.2019.00836
- Arya KN, Verma R, Garg RK. Estimating the minimal clinically important difference of an upper extremity recovery measure in subacute stroke patients. *Top Stroke Rehabil*. (2011) 18(Suppl 1):599–610. doi: 10.1310/tsr18s01-599
- Bundy DT, Souders L, Baranyai K, Leonard L, Schalk G, Coker R, et al. Contralateral brain-computer interface control of a powered exoskeleton for motor recovery in chronic stroke survivors. *Stroke*. (2017) 48:1908–15. doi: 10.1161/STROKEAHA.116.016304

36. Subramanian SK, Feldman AG, Levin MF. Spasticity may obscure motor learning ability after stroke. *J Neurophysiol.* (2018) 119:5–20. doi: 10.1152/jn.00362.2017
37. Delavaran H, Aked J, Sjunnesson H, Lindvall O, Norrving B, Kokaia Z, et al. Spontaneous recovery of upper extremity motor impairment after ischemic stroke: implications for stem cell-based therapeutic approaches. *Transl Stroke Res.* (2017) 8:351–61. doi: 10.1007/s12975-017-0523-9
38. Haselbach D, Renggli A, Carda S, Croquelois A. Determinants of neurological functional recovery potential after stroke in young adults. *Cerebrovasc Dis Extra.* (2014) 4:77–83. doi: 10.1159/000360218
39. Wortman-Jutt S, Edwards D. Poststroke aphasia rehabilitation: why all talk and no action. *Neurorehabil Neural Repair.* (2019) 33:235–44. doi: 10.1177/1545968319834901
40. Chen ML, Fu D, Boger J, Jiang N. Age-related changes in vibrotactile EEG response and its implications in BCI applications: a comparison between older and younger populations. *IEEE Trans Neural Syst Rehabil Eng.* (2019) 27:603–10. doi: 10.1109/TNSRE.2019.2890968
41. Rosjat N, Liu L, Wang BA, Popovych S, Tóth T, Viswanathan S, et al. Aging-associated changes of movement-related functional connectivity in the human brain. *Neuropsychologia.* (2018) 117:520–9. doi: 10.1016/j.neuropsychologia.2018.07.006
42. Kwah LK, Harvey LA, Diong J, Herbert RD. Models containing age and NIHSS predict recovery of ambulation and upper limb function six months after stroke: an observational study. *J Physiother.* (2013) 59:189–97. doi: 10.1016/S1836-9553(13)70183-8
43. Karaahmet OZ, Umay E, Gurcay E, Serçe A, Gundogdu I, Cakci A. The effect of premorbid features on post-stroke rehabilitation outcome. *Iran J Neurol.* (2018) 17:38–46.
44. Lee KB, Kim JS, Hong BY, Kim YD, Hwang BY, Lim SH. The motor recovery related with brain lesion in patients with intracranial hemorrhage. *Behav Neurol.* (2015) 2015:258161. doi: 10.1155/2015/258161
45. Schiemanck SK, Kwakkel G, Post MW, Kappelle LJ, Prevo AJ. Impact of internal capsule lesions on outcome of motor hand function at one-year post-stroke. *J Rehabil Med.* (2008) 40:96–101. doi: 10.2340/16501977-0130
46. Paci M, Nannetti L, D'Ippolito P, Lombardi B. Outcomes from ischemic stroke subtypes classified by the Oxfordshire Community Stroke Project: a systematic review. *Eur J Phys Rehabil Med.* (2011) 47:19–23.
47. Tedesco Triccas L, Kennedy N, Smith T, Pomeroy V. Predictors of upper limb spasticity after stroke? A systematic review and meta-analysis. *Physiotherapy.* (2019) 105:163–73. doi: 10.1016/j.physio.2019.01.004

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Wu, Ge, Ma, Pang, Cao, Zhang, Pan, Zhang and Dou. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.