



Congestive Heart Failure Exhibited Higher BMI With Lower Energy Intake and Lower Physical Activity Level: Data From the National Health and Examination Nutrition Survey

Tianyu Xu^{1†}, Haobin Zhou^{2†}, Zhuang Ma², Hao Zhang², Qingchun Zeng², Dingli Xu², Yuhui Zhang^{1*} and Jian Zhang^{1*}

¹ State Key Laboratory of Cardiovascular Disease, Heart Failure Center, National Center for Cardiovascular Disease, Fuwai Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China, ² State Key Laboratory of Organ Failure Research, Department of Cardiology, Nanfang Hospital, Southern Medical University, Guangzhou, China

OPEN ACCESS

Edited by:

Yuli Huang,
Southern Medical University, China

Reviewed by:

Jie Yu,
University of New South
Wales, Australia
Jun Lyu,
First Affiliated Hospital of Jinan
University, China

*Correspondence:

Jian Zhang
fwzhangjian62@126.com
Yuhui Zhang
yuhuizhangjoy@163.com

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Cardiovascular Metabolism,
a section of the journal
Frontiers in Cardiovascular Medicine

Received: 14 March 2021

Accepted: 08 April 2021

Published: 09 June 2021

Citation:

Xu T, Zhou H, Ma Z, Zhang H, Zeng Q,
Xu D, Zhang Y and Zhang J (2021)
Congestive Heart Failure Exhibited
Higher BMI With Lower Energy Intake
and Lower Physical Activity Level:
Data From the National Health and
Examination Nutrition Survey.
Front. Cardiovasc. Med. 8:680371.
doi: 10.3389/fcvm.2021.680371

Background: Despite that nutritional deficiency existed in congestive heart failure (CHF), there is a large amount of CHF patients suffering from obesity. This study aimed to identify the differences for increased BMI or obesity in CHF patients.

Methods: This cross-sectional study included adults from the National Health and Nutrition Examination Survey 2007–2016. Differences were compared between CHF participants vs. non-CHF participants, and BMI ≥ 30 kg/m² vs. BMI < 30 kg/m² CHF participants.

Results: CHF participants were with higher BMI, lower energy and macronutrient intake, lower physical activity level and longer rest time, and lower hematocrit and hemoglobin level (all $P < 0.05$) than non-CHF participants. The prevalence of BMI ≥ 30 kg/m² in participants with CHF was 53.48%. There was no significant difference observed in energy and macronutrient intake between CHF participants with BMI ≥ 30 kg/m² or < 30 kg/m². The water intake ($P = 0.032$), sedentary time ($P = 0.002$), and hematocrit ($P = 0.028$) were significantly different between CHF with BMI ≥ 30 kg/m² and with < 30 kg/m².

Conclusion: Compared with non-CHF participants, CHF participants exhibited higher BMI with lower energy and macronutrient intake, lower physical activity level, longer rest time, and hemodilution with lower hematocrit and hemoglobin level. Among CHF participants with BMI ≥ 30 kg/m², higher sedentary time and hematocrit were observed.

Keywords: congestive heart failure, body mass index, total nutrient intakes, physical activity, hemodilution, obesity

HIGHLIGHTS

- CHF participants were with higher BMI, lower energy and macronutrient intake, lower physical activity level and longer rest time, and lower hematocrit and hemoglobin level.
- Only water intake, sedentary time, and hematocrit were significantly different between CHF with BMI ≥ 30 kg/m² and with < 30 kg/m².

- Longer resting time and an unbalanced diet may be associated with a higher prevalence of increased BMI or obesity in CHF participants.

INTRODUCTION

Congestive heart failure (CHF) is a complex clinical condition that represents the final evolution of all cardiac diseases and a global public health problem which affects an estimated 26 million worldwide (1), which has a tremendous economic impact on the public healthcare system (2). Studies revealed that CHF is associated with alterations in cardiac energy metabolism (3) and that nutrition intake may influence the evolution of the disease progress (4). A deficiency in energy and macronutrient intake among CHF patients was reported, which caused an undernourished or malnutritional condition (5–7). Obesity [body mass index (BMI) ≥ 30 kg/m²], with up to 40% prevalence among CHF patients (8), is recognized as a major independent risk factor for the development of CHF (9) and has a paradoxical impact on the prognosis of CHF (10, 11). Studies indicated a non-linear U-shaped association between BMI and the risk of HF mortality, with a greater risk from being at the lowest group (mean BMI = 19.43 kg/m²), rather than being at the top category (mean BMI = 30.16 kg/m²) (12). The beneficial effects of weight loss in HF patients are still controversial, given the view that despite nutritional deficiency existed in CHF, there is a large amount of CHF patients suffering from obesity, and we still lacked studies to explain the reasons for the increased BMI in CHF patients. In this research, we used the data from the National Health and Examination Nutrition Survey (NHANES) 2007–2016 to identify the differences for increased BMI or obesity in CHF participants.

METHOD

Dataset

The NHANES data, a nationally representative multistage cross-sectional survey of the non-military and non-institutionalized population of the United States, was used as the data source (13). The data is released by the National Center for Health Statistics once every 2 years. Each survey cycle collects self-reported and directly measured information from participants who undergo a series of questionnaires in a detailed in-home interview such as health conditions, behaviors, and dietary intake, and a physical examination during mobile examination. Also, each participant provides their blood for laboratory tests. This survey included NHANES data from 2007 to 2016.

Study Population

To identify the study population, we implemented the following exclusion criteria: participants <18 years old, participants who were pregnant or breastfeeding at the time of the survey, and participants who lack information on CHF diagnosis, dietary data, examination data, laboratory data, or questionnaire data. The final sample for this study consisted of 660 CHF participants and 20,923 non-CHF participants.

Definition of CHF and Obesity

CHF was self-reported and was obtained from the medical condition files that were administered in the home by an interviewer using the Computer-Assisted Personal Interviewing system as part of the survey participant household interview. The final sample provided yes/no responses to the following question: “Has a doctor or other health professional ever told {you/SP} that {you/s/he}... had congestive HF?”

Height and weight were measured at the Mobile Examination Center (MEC) examination. BMI was calculated as weight in kilograms divided by the square of height in meters. BMI was analyzed as both a continuous variable and as a categorical variable dichotomizing into BMI ≥ 30 kg/m² (obese) and BMI < 30 kg/m² (non-obese) (14).

Sample Demographics, Dietary Condition, and Other Covariates

We obtained information on age and sex from the NHANES self-reported demographics data. Blood pressures and heart rate (resting) were measured in the MEC examination by a physician. Averages of up to four values for systolic and diastolic blood pressures were recorded. Intakes of energy, water, and macronutrients, which consisted of protein, carbohydrate, total sugars, total fat, and cholesterol, were obtained from self-reported dietary data, and an average of 2 days was recorded. Physical activity, including work, and recreational activities, was classified as vigorous, moderate, bicycling or walking, and sedentary and was obtained by self-reporting the number of days they engaged in that activity in a typical week and the average duration they engaged in that activity. For each activity, the Metabolic Equivalent of Task (MET)-hr-week was calculated by multiplying the number of days, the mean duration, and the respective MET level (MET-hr-week = days * duration * MET level). The MET levels for each activity are provided as vigorous work/recreational-related activity = 8 MET, moderate work/recreational-related activity = 4 MET, and walking or bicycling for transportation = 4 MET. It was analyzed as a continuous variable for total activates MET and MET in each activity. Hematocrit (%) and hemoglobin (g/dL), which may imply hemodilution (15), were acquired from blood collected during the MEC examination using methods described by the US Department of Health and Human Services.

Statistical Analysis

Data are expressed as the mean \pm standard deviation or as the number (%). Differences between groups were tested by the chi-square test for categorical variables and independent Student's *t*-test, as well as ANOVA with Tukey's *post-hoc* analysis for continuous variables, as appropriate. Since the sample size is large, a simple application of the Lyapunov or Lindeberg's Central Limit Theorem guarantees large sample convergence of the weighted mean to a standard normal distribution, ensuring that the *t*-statistic would have a limiting *t*-distribution. We also used restricted cubic splines with three knots at the 5th, 25th, 50th, 75th, and 95th centiles to flexibly model the association of sedentary time with BMI ≥ 30 kg/m² in CHF participants and non-CHF participants, respectively. We tested for potential non-linearity by using a likelihood ratio test comparing the

model with only a linear term against the model with linear and cubic spline terms. A P value < 0.05 was considered statistically significant. All tests were two-sided. All analyses were performed using R: A Language and Environment for Statistical Computing, version 3.1.3 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

Participants' Characteristics Between CHF and Non-CHF


The flowchart for participant selection is outlined in **Figure 1**. **Table 1** presents the clinical characteristics of CHF and non-CHF participants in NHANES 2007–2016. As noted, CHF participants were older than non-CHF participants ($P < 0.05$) with higher systolic blood pressure, lower diastolic blood pressure, and lower heart rate (all $P < 0.05$). Intake of total nutrients, including energy, protein, total carbohydrate, total sugar, total fat, and cholesterol, as well as water was much lower compared with that in non-CHF participants (all $P < 0.05$). Also, CHF participants had lower physical activity level ($P < 0.05$) and longer sedentary time ($P < 0.05$). Moreover, hemoglobin and hematocrit were lower in CHF participants ($P < 0.05$). However, BMI was higher in CHF participants than counterparts ($P < 0.05$).

Participants' Characteristics by BMI Category

As shown in **Table 2**, the proportions of BMI ≥ 30 kg/m² were 53.48% ($n = 353$) in CHF participants and 37.01% ($n = 7,744$) in non-CHF participants. Compared with BMI < 30 kg/m² CHF participants, BMI ≥ 30 kg/m² CHF participants showed significantly younger age, larger amount of water intake, longer sedentary time, and higher hematocrit (all $P < 0.05$). Among non-CHF participants, those with BMI ≥ 30 kg/m² were older and had higher blood pressure and heart rate and lower energy, protein, total carbohydrate, and total sugar intake but higher cholesterol and water intake (all $P < 0.05$). Physical activity level was lower and sedentary time was longer (both $P < 0.05$) in BMI ≥ 30 kg/m² non-CHF participants. Meanwhile, hemoglobin and hematocrit were lower in BMI ≥ 30 kg/m² non-CHF participants than BMI < 30 kg/m² ones (both $P < 0.05$). Restricted cubic spline showed that the risk of BMI ≥ 30 kg/m² in CHF participants was increased with the sedentary time (odds ratio 1.08, 95% confidence interval 1.03–1.13) (**Figure 2**), and such result could be also seen in participants without CHF (**Supplementary Figure 1**).

DISCUSSION

There are two major findings in the present study. First, compared with non-CHF participants, CHF participants exhibited higher BMI with lower energy and macronutrient intake, lower physical activity level, and longer rest time and hemodilution (lower hematocrit and hemoglobin). Second, higher water intake, sedentary time, and hematocrit were observed in CHF participants with BMI ≥ 30 kg/m²s, while energy and macronutrient intake and physical level were



similar between CHF participants with BMI ≥ 30 kg/m² and < 30 kg/m².

Excess body weight and a sedentary lifestyle are major public health problems worldwide (16). Obesity (BMI ≥ 30 kg/m²) has been considered as a major independent risk factor for cardiovascular disease (17). A previous study showed that up to 40% of patients with CHF suffered from obesity (8). The interplay between obesity and CHF is complex. Despite that obesity increases the risk of CHF, studies found that CHF patients who developed obesity or overweight (25–29.9 kg/m²) are associated with better prognosis compared with those who did not (18–20), with reductions in cardiovascular disease mortality (19 and 40%, respectively) and all-cause mortality (16 and 33%, respectively) compared to heart failure patients with normal weight (BMI 20–24.9 kg/m²), whereas heart failure patients with BMI < 20 kg/m² increased total and cardiovascular mortality (27 and 20%, respectively) (21). This phenomenon is termed as “obesity paradox,” and several potential reasons may account for it, such as earlier presentation, different etiology, greater metabolic reserves, protective cytokines, more tolerance of cardiac medications, different cardiorespiratory fitness level, and less cachexia (22). Based on previous studies, our study termed “obesity” as “BMI ≥ 30 kg/m²” to investigate the differences for increased BMI in CHF participants.

In our analysis, we recognized that CHF participants presented with higher BMI than non-CHF participants and a higher proportion of BMI ≥ 30 kg/m² among CHF participants. Meanwhile, we also noticed an inadequate energy and macronutrient intake in CHF participants, seen as significantly lower energy, protein, carbohydrate, total sugar, total fat, cholesterol, and water intake. This implied that nutritional deficiency existed in CHF participants. Malnutrition is commonly prevalent in CHF, with a reported incidence of 7.5 and 57% in severe or moderate malnutrition, respectively (23). These patients are older and frail, with negative calorie and nitrogen balance and energy availability (24) for physical

TABLE 1 | Characteristics of congestive heart failure group and non-congestive heart failure participants.

	Congestive heart failure (n = 660)	Non-congestive heart failure (n = 20,923)	P-value
Demographic and weight characteristics, mean (SD)			
Age (y)	66.46 (12.72)	48.65 (17.51)	<0.001
Sex			
Male (no., %)	360 (54.55)	10,153 (48.53)	<0.001
BMI (kg/m ²)	31.96 (8.54)	28.99 (6.80)	<0.001
SBP (mmHg)	129.82 (21.94)	123.26 (18.17)	<0.001
DBP (mmHg)	65.57 (15.03)	70.03 (12.76)	<0.001
Heart rate (beats/min)	71.02 (12.69)	72.81 (12.15)	<0.001
Total nutrient intake, mean (SD)			
Energy (kcal)	1734.71 (717.33)	2051.32 (872.86)	<0.001
Protein (gm)	69.85 (31.39)	80.65 (36.92)	<0.001
Carbohydrate (gm)	213.08 (94.19)	251.13 (111.62)	<0.001
Total sugars (gm)	93.56 (57.98)	110.98 (68.26)	<0.001
Total fat (gm)	66.36 (32.80)	76.58 (39.85)	<0.001
Cholesterol (mg)	266.10 (175.92)	286.80 (197.44)	0.0102
Water (gm)	531.16 (584.86)	646.00 (647.87)	<0.001
Physical activity, mean (SD)			
Total activity (MET-h/week)	24.54 (66.46)	57.68 (98.97)	<0.001
Vigorous activity (MET-h/week)	9.62 (42.83)	27.83 (70.54)	<0.001
Moderate activity (MET-h/week)	11.89 (28.80)	23.64 (42.33)	<0.001
Sedimentary activity (h/week)	6.81 (3.52)	5.79(3.39)	<0.001
Walk or bicycle (MET-h/week)	3.03 (14.95)	6.21 (20.27)	<0.001
Complete blood count—whole blood, mean (SD)			
Hemoglobin (g/dL)	13.48 (1.73)	14.08 (1.53)	<0.001
Hematocrit (%)	39.81 (4.82)	41.25 (4.24)	<0.001

BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; MET, metabolic equivalent of task; SD, standard deviation.

activity, and are triggered by multiple factors such as anorexia, malabsorption secondary to intestinal edema, high energy demand, and cytokine-induced hypercatabolism (7, 25) and worse outcomes (6, 26, 27). However, when compared with CHF participants with BMI < 30 kg/m², participants with BMI ≥ 30 kg/m² did not show a distinct increase in energy and macronutrient intake. A constellation of researches revealed that failing myocardium utilized glucose, switching from fatty acids, as the primary energetic substrate to produce ATP (28), which may exert a protective role in preventing cardiomyocytes from oxidative radical excess and cell damage (29, 30) but a lower energy production (31). With more total fat intake and less glucose intake, cells may have the less energetic substrate to produce enough energy. This would affect the metabolism in the body and cause adipose tissue accumulation and subsequently increased body weight. This may imply that inadequate or unbalanced energy and macronutrient intake accounts for the increase in BMI or obesity among CHF participants. A daily caloric intake of about 29 kcal/kg and a daily protein intake of 1.2–1.4 g/kg were recommended for CHF patients at normal weights and a less energy intake was required (20–24 kcal/kg/day), and a reduction in dietary fat intake to about 25% of total caloric intake (0.6–0.8 g/kg/day) was required for overweight and obese CHF patients (32, 33).

Our study found lower hematocrit and hemoglobin levels in CHF participants, while CHF with BMI ≥ 30 kg/m² participants had higher hematocrit and hemoglobin concentration than those with BMI < 30 kg/m². These results demonstrated that hemodilution and fluid retention are common in CHF, especially in those with lower BMI level (BMI < 30 kg/m²). This observation may be another potential reason to explain the “obesity paradox.” Considerable evidence demonstrated that total blood volume and cardiac output are positively correlated with the degree of excess body weight (18, 22, 34) and further preserved or even increased skeletal muscle mass (i.e., lean mass) (35, 36), which may cause BMI to be higher, and vice versa. This may explain the phenomenon of elevated hematocrit and hemoglobin concentration in CHF with BMI ≥ 30 kg/m² participants, and fluid retention may not be the reason for the increased BMI or obesity in CHF participants. Hemodilution was common in chronic heart failure (37) and has a deleterious effect as it may impair peripheral oxygen delivery (38) and is often neglected, as compensatory mechanisms may mask signs of volume (39). Our study found lower hematocrit and hemoglobin level in CHF participants, while CHF with BMI ≥ 30 kg/m² participants had higher hematocrit and hemoglobin concentration than BMI < 30 kg/m² ones. These results demonstrated that hemodilution and fluid retention are common in CHF, especially in those with

TABLE 2 | Characteristics of congestive heart failure group and non-congestive heart failure group (stratified by obese and non-obese).

BMI (kg/m ²)	Congestive heart failure			Non-congestive heart failure			*P-value
	BMI < 30 kg/m ² (n = 307)	BMI ≥ 30 kg/m ² (n = 353)	P value	BMI < 30 kg/m ² (n = 13,179)	BMI ≥ 30 kg/m ² (n = 7,744)	P value	
Demographic and weight characteristics, mean (SD)							
Age (y)	68.67 (12.86)	64.54 (12.30)	0.012	48.28 (18.17)	49.29 (16.30)	<0.001	<0.001
Sex							
Male (no., %)	178 (57.98)	181 (51.27)	<0.001	6,786 (51.49)	3,367 (43.48)	<0.001	<0.001
BMI (kg/m ²)	25.34 (3.03)	37.72 (7.56)	<0.001	24.95 (3.13)	35.86 (5.78)	<0.001	<0.001
SBP pressure (mmHg)	130.38 (23.38)	129.34 (20.63)	0.89	122.01 (18.53)	125.40 (17.33)	<0.001	<0.001
DBP (mmHg)	64.13 (14.56)	66.81 (15.33)	0.04	69.15 (12.45)	71.51 (13.16)	<0.001	<0.001
Heart rate (beats/min)	70.05 (12.37)	71.84 (12.92)	0.24	71.73 (11.9)	74.64 (12.23)	<0.001	<0.001
Total nutrient intakes, mean (SD)							
Energy (kcal)	1759.97 (716.89)	1712.74 (718.01)	0.90	2079.46 (894.56)	2003.43 (832.49)	<0.001	<0.001
Protein (gm)	70.93 (32.74)	68.91 (30.19)	0.90	81.30 (37.76)	79.55 (35.44)	0.005	<0.001
Carbohydrate (gm)	217.22 (95.26)	209.48 (93.24)	0.81	256.08 (114.39)	242.69 (106.21)	<0.001	<0.001
Total sugars (gm)	96.15 (59.90)	91.31 (56.23)	0.80	112.65 (70.08)	108.13 (64.96)	<0.001	<0.001
Total fat (gm)	65.65 (30.90)	66.98 (34.41)	0.97	76.48 (40.51)	76.73 (38.72)	0.97	<0.001
Cholesterol (mg)	261.71 (181.87)	269.92 (170.75)	0.95	282.87 (200.12)	293.50 (192.62)	<0.001	<0.001
Water (gm)	457.69 (484.71)	595.06 (653.72)	0.032	615.04 (621.65)	698.70 (687.04)	<0.001	<0.001
Physical activity, mean (SD)							
Total activity (MET-h/week)	28.17 (67.00)	21.38 (65.91)	0.81	60.85 (101.41)	52.28 (94.43)	<0.001	<0.001
Vigorous activity (MET-h/week)	11.36 (41.45)	8.11 (43.99)	0.93	29.53 (71.50)	24.93 (68.79)	<0.001	<0.001
Moderate activity (MET-h/week)	12.85 (29.27)	11.06 (28.41)	0.94	24.39 (42.80)	22.36 (41.49)	<0.001	<0.001
Sedimentary activity (h/week)	6.32 (3.26)	7.24 (3.69)	0.002	5.62 (3.32)	6.08 (3.48)	<0.001	<0.001
Walk or bicycle (MET-h/week)	3.97 (16.42)	2.21 (13.51)	0.68	6.93 (21.51)	4.99 (17.92)	<0.001	<0.001
Complete blood count—whole blood, mean (SD)							
Hemoglobin (g/dL)	13.34 (1.80)	13.61 (1.65)	0.121	14.12 (1.50)	14.00 (1.57)	<0.001	<0.001
Hematocrit (%)	39.31 (5.02)	40.25 (4.61)	0.028	41.32 (4.21)	41.13 (4.30)	0.009	<0.001

*P-value was analysis between four groups.


BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; MET, metabolic equivalent of task; SD, standard deviation.

lower BMI level (BMI<30 kg/m²). Previous studies indicated a higher mortality rate in patients with hemodilution than in those with hemoconcentration in acute heart failure patients (15), while fluid restriction could only improve signs and symptoms of chronic heart failure in patients in moderate to severe chronic heart failure (40) and aggressive fluid removal positively affected survival (41). In the present study, CHF with BMI ≥ 30 kg/m² participants had higher hematocrit and hemoglobin concentration than BMI < 30 kg/m² ones, indicating that obese CHF participants were less likely to develop hemodilution. This could be one explanation for the “obesity paradox.” Furthermore, hemodynamic changes including increased stroke volume and increased arterial pressure may compensate the impaired peripheral oxygen delivery, leading CHF patients with increased BMI or obesity to have better prognoses. Still, we could not ignore that there was fluid retention in our CHF group, which implied an inadequate usage of diuretics and fluid management should be enhanced.

Another finding in our analysis is that we demonstrated that CHF participants had lower physical activity levels and spent a longer time in rest, especially participants with BMI ≥ 30 kg/m². It was revealed that sedentary time caused

metabolic alterations at the muscle level and next influenced gross metabolic disturbances in the whole body (42). It has been reported that prolonged sedentary time would impair mitochondrial function by elevating oxidative stress levels (43), which decreased the mitochondrial respiration level (44) and caused insufficient ATP production for daily activity and metabolism. Emerging evidences have demonstrated a significant dose–response association between sitting time and cardiovascular disease mortality (45–48), and the relative risks associated with sedentary time were higher among participants without regular physical activity (49, 50). Meanwhile, there have been proven cardioprotective effects of regular physical activity on cardiovascular health, improving cardiac compliance, reducing arterial stiffness and ventricular afterload, and finally reducing the risk of future cardiac dysfunction and improving cardiovascular outcomes (42). Moreover, researches showed that an increasing lean mass with resistance exercise training could effectively improve muscular fitness in CHF (51, 52), which could prevent sarcopenia (53) or even cachexia in CHF patients and have better prognosis (54).

Our investigation has several limitations. We only included macronutrient in our analysis and lacked the data



of micronutrient intake. Therefore, the present study may have underestimated the nutritional deficit and its effects on obesity among participants. Also, details regarding the etiology, subtype (HF with reduced ejection fraction vs. HF with preserved ejection fraction), severity of HF, and complications were not available. Moreover, CHF participants included in our analysis were self-reported in the NHANES survey. This may lead to possible selection, reporting, and recall bias. Additionally, we did not include socioeconomic data, and thus we may neglect the socioeconomic impacts on nutrition intake and lifestyle self-management of CHF participants. Finally, we lacked follow-up data of these participants, including BMI changes and relative outcomes, so we were unable to recognize the effects of nutrients intake and physical activity on BMI changes and prognoses in participants with CHF.

CONCLUSION

Higher water intake, sedentary time, and hematocrit were observed in BMI ≥ 30 kg/m² CHF participants. It seems that longer resting time and an unbalanced diet may be associated

REFERENCES

- Ambrosy AP, Fonarow GC, Butler J, Chioncel O, Greene SJ, Vaduganathan M, et al. The global health and economic burden of hospitalizations for heart failure: lessons learned from hospitalized heart failure registries. *J Am Coll Cardiol.* (2014) 63:1123–33. doi: 10.1016/j.jacc.2013.11.053
- Ziaian B, Fonarow GC. Epidemiology and aetiology of heart failure. *Nat Rev Cardiol.* (2016) 13:368–78. doi: 10.1038/nrcardio.2016.25
- Lourenço BH, Vieira LP, Macedo A, Nakasato M, Marucci MF, Bocchi EA. Nutritional status and adequacy of energy and nutrient intakes among heart failure patients. *Arq Bras Cardiol.* (2009) 93:541–8. doi: 10.1590/S0066-782X2009001100016
- Sciatti E, Lombardi C, Ravera A, Vizzardi E, Bonadei I, Carubelli V, et al. Nutritional deficiency in patients with heart failure. *Nutrients.* (2016) 8:442. doi: 10.3390/nu8070442
- Makarewicz-Wujec M, Kozłowska-Wojciechowska M, Sygnowska E, Waśkiewicz A. Does heart failure determine the nutrition of patients? *Kardiol Pol.* (2014) 72:56–63. doi: 10.5603/KP.a2013.0184
- Chien SC, Lo CI, Lin CF, Sung KT, Tsai JP, Huang WH, et al. Malnutrition in acute heart failure with preserved ejection fraction: clinical correlates and prognostic implications. *ESC Heart Fail.* (2019) 6:953–64. doi: 10.1002/ehf2.12501
- Kinugawa S, Fukushima A. Malnutrition in heart failure: important but undervalued issue. *JACC Heart Fail.* (2018) 6:487–88. doi: 10.1016/j.jchf.2018.03.014
- Kapoor JR, Heidenreich PA. Obesity and survival in patients with heart failure and preserved systolic function: a U-shaped relationship. *Am Heart J.* (2010) 159:75–80. doi: 10.1016/j.ahj.2009.10.026
- Hu G, Jousilahti P, Antikainen R, Katzmarzyk PT, Tuomilehto J. Joint effects of physical activity, body mass index, waist circumference, and

with a higher prevalence of increased BMI or obesity in CHF participants. Future research is warranted to explore the mechanisms underlying this finding and whether intentional weight loss with combination of diet, exercise, and others could be contributing to better health outcomes in BMI ≥ 30 kg/m² CHF participants.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author/s.

ETHICS STATEMENT

This study was reviewed and approved by the National Center for Health Statistics research ethics review board, and written informed consent was obtained from all NHANES participants.

AUTHOR CONTRIBUTIONS

TX: study concept and design, acquisition, analysis, or interpretation of data, and drafting of the manuscript. HZho: study concept and design, acquisition, analysis, or interpretation of data, and statistical analysis. ZM and HZha: acquisition, analysis, or interpretation of data, and statistical analysis. QZ and DX: critical revision of the manuscript for important intellectual content, administrative, technical, or material support, and study supervision. YZ and JZ: study concept and design, critical revision of the manuscript for important intellectual content, administrative, technical, or material support, and study supervision. All authors read and approved the final manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcvm.2021.680371/full#supplementary-material>

- waist-to-hip ratio on the risk of heart failure. *Circulation*. (2010) 121:237–44. doi: 10.1161/CIRCULATIONAHA.109.887893
10. Oreopoulos A, Padwal R, Kalantar-Zadeh K, Fonarow GC, Norris CM, McAlister FA. Body mass index and mortality in heart failure: a meta-analysis. *Am Heart J*. (2008) 156:13–22. doi: 10.1016/j.ahj.2008.02.014
 11. Loprinzi PD. Physical activity, weight status, and mortality among congestive heart failure patients. *Int J Cardiol*. (2016) 214:92–4. doi: 10.1016/j.ijcard.2016.03.180
 12. Milajerdi A, Djafarian K, Shah-Bidar S, Speakman JR. Pre- and post-diagnosis body mass index and heart failure mortality: a dose-response meta-analysis of observational studies reveals greater risk of being underweight than being overweight. *Obes Rev*. (2018) 20:252–61. doi: 10.1111/obr.12777
 13. Zipf G, Chiappa M, Porter KS, Ostchega Y, Lewis BG, Dostal J. National health and nutrition examination survey: plan and operations, 1999–2010. *Vital Health Stat 1*. (2013) 1–37.
 14. Pi-Sunyer FX, Becker DM, Bouchard C, Carleton RA, Colditz GA, Dietz WH, et al. Executive summary of the clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. *Arch Intern Med*. (1998) 158:1855–67. doi: 10.1001/archinte.158.17.1855
 15. Zhou H, Xu T, Huang Y, Zhan Q, Huang X, Zeng Q, et al. The top tertile of hematocrit change during hospitalization is associated with lower risk of mortality in acute heart failure patients. *BMC Cardiovasc Disord*. (2017) 17:235. doi: 10.1186/s12872-017-0669-0
 16. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. (2014) 384:766–81. doi: 10.1016/S0140-6736(14)60460-8
 17. Lavie CJ, De Schutter A, Parto P, Jahangir E, Kokkinos P, Ortega FB, et al. Obesity and prevalence of cardiovascular diseases and prognosis—the obesity paradox updated. *Prog Cardiovasc Dis*. (2016) 58:537–47. doi: 10.1016/j.pcad.2016.01.008
 18. Lavie CJ, Alpert MA, Arena R, Mehra MR, Milani RV, Ventura HO. Impact of obesity and the obesity paradox on prevalence and prognosis in heart failure. *JACC Heart Fail*. (2013) 1:93–102. doi: 10.1016/j.jchf.2013.01.006
 19. Futter JE, Cleland JG, Clark AL. Body mass indices and outcome in patients with chronic heart failure. *Eur J Heart Fail*. (2011) 13:207–13. doi: 10.1093/eurjhf/hfq218
 20. Bozkurt B, Deswal A. Obesity as a prognostic factor in chronic symptomatic heart failure. *Am Heart J*. (2005) 150:1233–9. doi: 10.1016/j.ahj.2005.02.004
 21. Sharma A, Lavie CJ, Borer JS, Vallakati A, Goel S, Lopez-Jimenez F, et al. Meta-analysis of the relation of body mass index to all-cause and cardiovascular mortality and hospitalization in patients with chronic heart failure. *Am J Cardiol*. (2015) 115:1428–34. doi: 10.1016/j.amjcard.2015.02.024
 22. Parto P, Lavie CJ, Arena R, Bond S, Popovic D, Ventura HO. Body habitus in heart failure: understanding the mechanisms and clinical significance of the obesity paradox. *Future Cardiol*. (2016) 12:639–53. doi: 10.2217/fca-2016-0029
 23. Sze S, Pellicori P, Zhang J, Clark AL. Malnutrition, congestion and mortality in ambulatory patients with heart failure. *Heart*. (2019) 105:297–306. doi: 10.1136/heartjnl-2018-313312
 24. Sanches MDK, Dalira SPI, Clausell N, Corrêa SG. Adequacy of energy and nutrient intake in patients with heart failure. *Nutr Hosp*. (2015) 31:500–7. doi: 10.1590/s0066-782x2009001100016
 25. Aquilani R, Opasich C, Verri M, Boschi F, Febo O, Pasini E, et al. Is nutritional intake adequate in chronic heart failure patients? *J Am Coll Cardiol*. (2003) 42:1218–23. doi: 10.1016/S0735-1097(03)00946-X
 26. Maeda D, Kanzaki Y, Sakane K, Ito T, Sohmiya K, Hoshiga M. Prognostic impact of a novel index of nutrition and inflammation for patients with acute decompensated heart failure. *Heart Vessels*. (2020) 35:1201–8. doi: 10.1007/s00380-020-01590-4
 27. Nakamura T, Matsumoto M, Haraguchi Y, Ishida T, Momomura SI. Prognostic impact of malnutrition assessed using geriatric nutritional risk index in patients aged ≥ 80 years with heart failure. *Eur J Cardiovasc Nurs*. (2020) 19:172–7. doi: 10.1177/1474515119864970
 28. Taegtmeyer H, Sen S, Vela D. Return to the fetal gene program: a suggested metabolic link to gene expression in the heart. *Ann N Y Acad Sci*. (2010) 1188:191–8. doi: 10.1111/j.1749-6632.2009.05100.x
 29. Park TS, Goldberg IJ. Sphingolipids, lipotoxic cardiomyopathy, and cardiac failure. *Heart Fail Clin*. (2012) 8:633–41. doi: 10.1016/j.hfc.2012.06.003
 30. Libby P, Maroko PR, Braunwald E. The effect of hypoglycemia on myocardial ischemic injury during acute experimental coronary artery occlusion. *Circulation*. (1975) 51:621–6. doi: 10.1161/01.CIR.51.4.621
 31. Neubauer S. The failing heart—an engine out of fuel. *N Engl J Med*. (2007) 356:1140–51. doi: 10.1056/NEJMra063052
 32. Kopple JD. National kidney foundation K/DOQI clinical practice guidelines for nutrition in chronic renal failure. *Am J Kidney Dis*. (2001) 37:S66–70. doi: 10.1053/ajkd.2001.20748
 33. Arkinwall MJ, Bruce CR, Clark SA, Rickards CA, Burke LM, Hawley JA. Regulation of fuel metabolism by preexercise muscle glycogen content and exercise intensity. *J Appl Physiol* (1985). (2004) 97:2275–83. doi: 10.1152/jappphysiol.00421.2004
 34. Alpert MA. Obesity cardiomyopathy: pathophysiology and evolution of the clinical syndrome. *Am J Med Sci*. (2001) 321:225–36. doi: 10.1097/00000441-200104000-00003
 35. Collis T, Devereux RB, Roman MJ, de Simone G, Yeh J, Howard BV, et al. Relations of stroke volume and cardiac output to body composition: the strong heart study. *Circulation*. (2001) 103:820–5. doi: 10.1161/01.CIR.103.6.820
 36. Carbone S, Lavie CJ, Arena R. Obesity and heart failure: focus on the obesity paradox. *Mayo Clin Proc*. (2017) 92:266–79. doi: 10.1016/j.mayocp.2016.11.001
 37. Androne AS, Katz SD, Lund L, LaManca J, Hudaihed A, Hryniewicz K, Mancini DM. Hemodilution is common in patients with advanced heart failure. *Circulation*. (2003) 107:226–9. doi: 10.1161/01.CIR.0000052623.16194.80
 38. Chapler CK, Cain SM. The physiologic reserve in oxygen carrying capacity: studies in experimental hemodilution. *Can J Physiol Pharmacol*. (1986) 64:7–12. doi: 10.1139/y86-002
 39. Wilson JR, Lanoco V, Frey MJ, Ferraro N. Effect on peripheral arterioles of chronic fluid and sodium retention in heart failure. *J Am Coll Cardiol*. (1988) 12:202–8. doi: 10.1016/0735-1097(88)90375-0
 40. Philipson H, Ekman I, Forslund HB, Swedberg K, Schaufelberger M. Salt and fluid restriction is effective in patients with chronic heart failure. *Eur J Heart Fail*. (2013) 15:1304–10. doi: 10.1093/eurjhf/hft097
 41. Testani JM, Chen J, McCauley BD, Kimmel SE, Shannon RP. Potential effects of aggressive decongestion during the treatment of decompensated heart failure on renal function and survival. *Circulation*. (2010) 122:265–72. doi: 10.1161/CIRCULATIONAHA.109.933275
 42. Lavie CJ, Ozemek C, Carbone S, Katzmarzyk PT, Blair SN. Sedentary behavior, exercise, and cardiovascular health. *Circ Res*. (2019) 124:799–815. doi: 10.1161/CIRCRESAHA.118.312669
 43. Joseph AM, Adhithetty PJ, Buford TW, Wohlgenuth SE, Lees HA, Nguyen LM, et al. The impact of aging on mitochondrial function and biogenesis pathways in skeletal muscle of sedentary high- and low-functioning elderly individuals. *Aging Cell*. (2012) 11:801–9. doi: 10.1111/j.1474-9726.2012.00844.x
 44. Conley KE, Amara CE, Bajpeyi S, Costford SR, Murray K, Jubrias SA, et al. Higher mitochondrial respiration and uncoupling with reduced electron transport chain content in vivo in muscle of sedentary vs. active subjects. *J Clin Endocrinol Metab*. (2013) 98:129–36. doi: 10.1210/jc.2012-2967
 45. Young DR, Hivert MF, Alhassan S, Camhi SM, Ferguson JF, Katzmarzyk PT, et al. Sedentary behavior and cardiovascular morbidity and mortality: a science advisory from the American Heart Association. *Circulation*. (2016) 134:e262–79. doi: 10.1161/CIR.0000000000000440
 46. Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc*. (2009) 41:998–1005. doi: 10.1249/MSS.0b013e3181930355
 47. Chau JY, Grunseit AC, Chey T, Stamatakis E, Brown WJ, Matthews CE, et al. Daily sitting time and all-cause mortality: a meta-analysis. *PLOS ONE*. (2013) 8:e80000. doi: 10.1371/journal.pone.0080000

48. Pandey A, Salahuddin U, Garg S, Ayers C, Kulinski J, Anand V, et al. Continuous dose-response association between sedentary time and risk for cardiovascular disease: a meta-analysis. *JAMA Cardiol.* (2016) 1:575–83. doi: 10.1001/jamacardio.2016.1567
49. Biswas A, Oh PI, Faulkner GE, Bajaj RR, Silver MA, Mitchell MS, et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med.* (2015) 162:123–32. doi: 10.7326/M14-1651
50. Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N, Powell KE, et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet.* (2016) 388:1302–10. doi: 10.1016/S0140-6736(16)30370-1
51. Lavie CJ, Forman DE, Arena R. Bulking Up skeletal muscle to improve heart failure prognosis. *JACC Heart Fail.* (2016) 4:274–6. doi: 10.1016/j.jchf.2015.12.005
52. Ventura HO, Carbone S, Lavie CJ. Muscling up to improve heart failure prognosis. *Eur J Heart Fail.* (2018) 20:1588–90. doi: 10.1002/ejhf.1314
53. Prado CM, Heymsfield SB. Lean tissue imaging: a new era for nutritional assessment and intervention. *JPEN J Parenter Enteral Nutr.* (2014) 38:940–53. doi: 10.1177/0148607114550189
54. Bekfani T, Pellicori P, Morris DA, Ebner N, Valentova M, Steinbeck L, et al. Sarcopenia in patients with heart failure with preserved ejection fraction: Impact on muscle strength, exercise capacity and quality of life. *Int J Cardiol.* (2016) 222:41–6. doi: 10.1016/j.ijcard.2016.07.135

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Xu, Zhou, Ma, Zhang, Zeng, Xu, Zhang and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.