



UNIVERSITI PUTRA MALAYSIA

**BIOACTIVITY OF AZADIRACHTA EXCELSA JACK.
EXTRACTIVES ON SELECTED INSECTS**

MARIA LOURDES T. LARDIZABAL

FH 1998 1

**BIOACTIVITY OF *AZADIRACHTA EXCELSA* Jack.
EXTRACTIVES ON SELECTED INSECTS**

By

MARIA LOURDES T. LARDIZABAL

**Thesis Submitted in Fulfilment of the Requirements for the
Degree of Master of Science in the Faculty of Forestry,
Universiti Putra Malaysia**

April 1998



*Dedicated especially to,
Mom and Dad,
My brother Dominic,
and
My beloved Christopher,
who constantly inspired and prayed for me
throughout the completion of this thesis.*



ACKNOWLEDGEMENTS

First and foremost, praise be to God the Almighty for giving me the strength, courage, guidance and unending love without which this study could not have been completed.

I would like to express my profound gratitude to Assoc. Prof Dr. Ahmad Said Sajap, the chairman of my supervisory committee for his constant guidance, suggestions and encouragement throughout the course of my study and also in preparation of this thesis. My sincere appreciation is also extended to Assoc. Prof. Dr. Mohd. Hamami Sahri, Dr. Faizah Abood Haris and Assoc. Prof. Dr. Faujan Ahmad, members of my advisory committee, for their invaluable suggestions, comments and constructive criticisms which brought to the completion of this thesis.

I am indebted to the Ministry of Science, Technology and the Environment, Malaysia, for providing a study grant (PASCA Siswazah) which financially supported me throughout the course of my studies.

Acknowledgements are also extended to Mrs. Aida Marsidi, Mrs. Halimah Husin and all the staff of the Faculty of Forestry, University Putra Malaysia, who have



helped and provided me the facilities needed throughout the course of my studies. I would also like to record my appreciation to the field staff of the Plantation Unit and Wood Industry workshop at the Forest Research Institute Malaysia (FRIM) and to Puan Fatimah of the Insectory Unit at MARDI Station, Serdang, My grateful thanks also to the Field Unit staff of Universiti Putra Malaysia for their assistance in felling a few pine trees at the pine plantation UPM. I would also like to thank the Laboratory assistant of the Infrared Spectroscopy Laboratory at the Chemistry Department, UPM, for helping me to analyse my plant extractive samples.

Thank you very much, mom and dad for your constant support and undividing love and also to my dearest brother Dominic, for your dedicated assistance during my experiment, moral and financial support, encouragements and for your guidance and help in statistical analysis of my data. To all my dearest friends, Phillis, Shirley, Boyd, Audrey, Florence, Rose, John Keen, Evelyn, Patricia C., Ronnie, Jenny, Jean, and all my friends in CSSUPM, which could not be mentioned here, and especially to Dr. Mary Huang and Fr. Lawrence, thank you so much for your encouragement, moral support and prayers.

Finally, especially to Christopher Yong, thank you so much for your sincere love, care and concern and also for being the source of inspiration and light to my life at all times.



TABLE OF CONTENTS

		Page
ACKNOWLEDGEMENTS		iii
TABLE OF CONTENTS		v
LIST OF TABLES		ix
LIST OF FIGURES		xi
LIST OF PLATES		xiii
LIST OF ABBREVIATIONS		xiv
ABSTRACT		xvi
ABSTRAK		xix
 CHAPTER:		
I	GENERAL INTRODUCTION	1
II	LITERATURE REVIEW	5
	<i>Azadirachta excelsa</i> (Jack.) Jacobs.....	5
	Taxonomy.....	5
	Geographic Distribution.....	6
	Ecology.....	7
	Morphology.....	8
	Uses of <i>Azadirachta excelsa</i>	12
	Insecticidal Properties.....	13
	Properties and Potential of Natural Pesticides from the <i>Azadirachta</i> spp.....	15
	Azadirachtin.....	15
	Effects of Azadirachtin on Insects.....	17
	<i>Spodoptera litura</i> (Fabricus).....	19
	Distribution.....	19
	Host Range.....	20
	Biology.....	20
	Effect of Neem Extracts on <i>Spodoptera litura</i>	24
	Subterranean Termites.....	25
	Distribution.....	25
	Biology.....	25
	Termites as Pests.....	27
	<i>Coptotermes curvignathus</i> (Holmgren).....	27
	Termite Control.....	28
	<i>Callosobruchus</i> spp.....	30



	Distribution.....	30
	Biology.....	30
	<i>Callosobruchus chinensis</i> Linn.....	31
	Infestation.....	32
	Previous Studies on Weevil Control.....	34
III	EXTRACTION OF <i>AZADIRACHTA EXCELSA</i> PLANT EXTRACTIVES.....	35
	Introduction.....	35
	Materials and Methods.....	36
	Plant Samples.....	36
	Preparation of Plant Samples.....	37
	Extraction of Plant Extractives.....	37
	Preliminary Determination of Chemical Components Using TLC and Infrared Spectrometer.....	39
	Results and Discussion.....	40
	Summary.....	52
IV	BIOASSAY TEST ON <i>SPODOPTERA LITURA</i> FABRICUS.....	53
	Introduction.....	53
	Materials and Methods.....	55
	Test Insects.....	55
	Food Source.....	56
	Bioassay.....	56
	Assessment.....	59
	Data Analysis.....	59
	Results.....	61
	Leaf Area Consumed by <i>Spodoptera litura</i>	61
	<i>Spodoptera litura</i> Mortality.....	68
	Moulting and Larval Period.....	76
	Head Capsule Width.....	80
	Pupal Weight.....	84
	Discussion.....	88
V	BIOASSAY TEST ON <i>CALLOSOBRUCHUS</i> <i>CHINENSIS</i>.....	92
	Introduction.....	92
	Materials and Methods.....	95



	Mass Rearing of Insects.....	95
	Bioassay.....	96
	Data Analysis.....	97
	Results.....	98
	Effect on Oviposition Rate.....	98
	Effect on Adult Emergence.....	101
	Effect on Percentage of Adult Emergence.....	103
	Discussion.....	105
VI	BIOASSAY TEST ON <i>COPTOTERMES</i>	
	<i>CURVIGNATHUS</i>	108
	Introduction.....	108
	Materials and Methods.....	110
	Termite Species.....	110
	Termite Collection.....	110
	Filter Paper Treatment.....	115
	Exposure of Treated Filter Papers on Termites.....	115
	Data Analysis.....	116
	Termite Mortality.....	116
	Filter Paper Weight Loss.....	117
	Results.....	118
	Termite Mortality.....	118
	Filter Paper Weight Loss.....	128
	Discussion.....	137
VII	SENTANG WOOD CONSUMPTION TEST	
	BY TERMITES	140
	Introduction.....	140
	Materials and Methods.....	142
	A) Laboratory Choice Feeding Test.....	142
	B) Field Accelerated Stake Test.....	146
	Data Analysis.....	150
	Results.....	150
	A) Laboratory Test on Choice Feeding.....	150
	B) Accelerated Field Testing.....	153
	Discussion.....	155



VIII	SUMMARY AND CONCLUSION.....	158
IX	RECOMMENDATIONS.....	163
	REFERENCES.....	165
	APPENDICES.....	183
	Appendix A.....	183
	Appendix B.....	184
	VITA.....	190



LIST OF TABLES

Table	Page
2.1 Effect of Azadirachtin on Insects.....	18
3.1 R _f Values of TLC Results from Bark Samples.....	44
3.2 R _f Values of TLC Results from Leaves Samples.....	45
3.3 R _f Values of TLC Results from Trunk Samples.....	47
3.4 R _f Values of TLC Results from Branch Samples.....	48
3.5 R _f Values of TLC Results from Twigs Samples.....	49
3.6 Infrared Spectroscopy Results of Plant Extracts (ν_{max} values in cm^{-1}).....	51
4.1 Mean Moulting and Larval Period (days) of <i>S.litura</i> Larva Exposed to <i>A.excelsa</i> Leaf Extracts.....	77
4.2 Mean Moulting and Larval Period (days) of <i>S.litura</i> Larva Exposed to <i>A.excelsa</i> Bark Extracts.....	79
4.3 Mean (\pm SE) Head Capsule Width (mm) of Last Instar <i>S.litura</i> Larva Exposed to <i>A.excelsa</i> Extractives.....	81
4.4 Mean (\pm SE) Pupal Weight (g) of <i>S.litura</i> Larva Exposed to <i>A.excelsa</i> Plant Extractives.....	85
5.1 Efficacy of <i>A.excelsa</i> Leaf and Bark Extracts on Oviposition Rate, Emergence and Percentage Adult Emergence of <i>C.chinensis</i>	100
6.1 Mean (\pm SE) Mortality of <i>C.curvignathus</i> Exposed to <i>A.excelsa</i> Plant Extractives.....	118
6.2 Analysis of Variance Table for Termite Mortality.....	127



6.3 Mean (\pm SE) Weight Loss (mg) of Filter Paper Treated with <i>A.excelsa</i> Plant Extractives.....	129
6.4 Analysis of Variance Table for Filter Paper Weight Loss	136
7.1 Wood Density Range.....	143
7.2 Mean (\pm SE) Wood Consumed by Termites on Different Wood Species (Choice Feeding-Laboratory Test).....	150
7.3 Mean (\pm SE) Wood Consumed by Termites on Different Wood Species (Field Accelerated Test).....	153



LIST OF FIGURES

Figure	Page
2.1 Geographical Distribution of <i>Azadirachta excelsa</i> (Jack) Jacobs.....	6
2.2 Chemical Structure of Marrangin.....	15
2.3 Chemical Structure of Azadirachtin.....	16
4.1 Mean Leaf Area Consumed by <i>S. litura</i> Larva Exposed to <i>A. excelsa</i> Leaf Extracts.....	62
4.2 Mean Leaf Area Consumed by <i>S. litura</i> Larva Exposed to <i>A. excelsa</i> Bark Extracts.....	64
4.3 Comparison of Mean Leaf Area Consumed by <i>S. litura</i> larva Exposed to <i>A. excelsa</i> Hexane Extracts of the Leaf and Bark.....	66
4.4 Comparison of Mean Leaf Area Consumed by <i>S. litura</i> Larva Exposed to <i>A. excelsa</i> Methanol Extracts of Leaf and Bark.....	67
4.5 Mean (%) Mortality of <i>S. litura</i> Larva Exposed to <i>A. excelsa</i> Leaf Extracts.....	70
4.6 Mean (%) Mortality of <i>S. litura</i> Larva Exposed to <i>A. excelsa</i> Bark Extracts.....	71
4.7 Comparison of Mean Mortality (%) of <i>S. litura</i> Exposed to <i>A. excelsa</i> Hexane Extracts of Leaf and Bark.....	73
4.8 Comparison of Mean Mortality (%) of <i>S. litura</i> Exposed to <i>A. excelsa</i> Methanol Extracts of Leaf and Bark.....	75
4.9 Head Capsule Width of Last Instar <i>S. litura</i> Larvae Exposed to <i>A. excelsa</i> Plant Extracts.....	83
4.10 Pupal Weight of <i>S. litura</i> Larvae Exposed to <i>A. excelsa</i> Extractives.....	87



6.1 Mean Termite Mortality (%) of <i>C. curvignathus</i> Exposed to <i>A. excelsa</i> Leaf Extracts.....	121
6.2 Mean Termite Mortality (%) of <i>C. curvignathus</i> Exposed to <i>A. excelsa</i> Bark Extracts.....	123
6.3 Mean Termite Mortality (%) of <i>C. curvignathus</i> Exposed to <i>A. excelsa</i> Hexane extracts of Leaf and Bark.....	125
6.4 Mean Termite Mortality (%) of <i>C. curvignathus</i> Exposed to <i>A. excelsa</i> Methanol Extracts of Leaf and Bark.....	126
6.5 Mean Weight Loss of Filter Paper Treated with <i>A. excelsa</i> Leaf Extracts.....	131
6.6 Mean Weight Loss of Filter Paper Treated with <i>A. excelsa</i> Bark Extracts.....	133
6.7 Mean Weight Loss of Filter Paper Treated with <i>A. excelsa</i> Hexane Extracts of Leaf and Bark.....	134
6.8 Mean Weight Loss of Filter Paper Treated with <i>A. excelsa</i> Methanol Extracts of Leaf and Bark.....	135
7.1 Mean Wood Weight Loss of Timber Species exposed to Termites (Choice-Feeding Lab. Test).....	152
7.2 Mean Wood Weigh Loss of Different Timber Species Exposed to Termites (Field Test).....	154



LIST OF PLATES

Plate		Page
1	<i>Azadirachta excelsa</i> Leaf Clusters.....	9
2	<i>Azadirachta excelsa</i> Leaf.....	11
3	Adult Male and Female <i>Spodoptera litura</i>	22
4	Mature <i>Spodoptera litura</i> Caterpillar.....	22
5	<i>Spodoptera litura</i> Pupa.....	23
6	Infested Cowpea Seeds.....	33
7	Adult <i>Callosobruchus chinensis</i> Emergence Hole.....	33
8	Rotary Evaporator.....	38
9	25 Compartment Plastic Plate.....	58
10	Culture Jars Capped with Muslin Cloths.....	95
11	Wooden Stakes Sandwich Bundle.....	111
12	Infested Wood Bundle Carefully Taken Out from the Ground.....	113
13	Infested Wood Bundles Kept in Plastic Containers.....	114
14	Termite Collection Method.....	114
15	Remnants of Treated Filter Paper After Exposure to Termites.....	128
16	Test Blocks Placed Radially in Kilner Jars.....	145
17	Wood Bundle of Accelerated Test Stakes.....	148



LIST OF ABBREVIATIONS

ANOVA	=	Analysis of Variance
AWPA	=	American Wood Preservers' Association
AZA	=	Azadirachtin
BRCs	=	Brain-ring Gland Complexes
CA	=	Corpora Allata
DMRT	=	Duncan's Multiple Range Test
IGR	=	Insect Growth Regulator
IR	=	Infrared
JPSM	=	Jabatan Perhutanan Semenanjung Malaysia (Forestry Department Peninsular Malaysia)
LHW	=	Light Hardwood
MARDI	=	Malaysian Agriculture Research and Development Institute
MAPPS	=	Malaysian Plant Protection Society
M.C.	=	Moisture Content
MHW	=	Medium Hardwood
MSTAT-C	=	Microcomputer Statistical Program
MTIB	=	Malaysian Timber Industry Board
PROC GLM	=	Procedure Generalised Linear Model
PTTH	=	Prthoracicotropic Hormone
R_f	=	Retention Factor
RH	=	Relative Humidity
RRIM	=	Rubber Research Institute Malaysia
SAS	=	Statistical Analysis System
S.E.	=	Standard Error

(continued).....



LIST OF ABBREVIATIONS (continued)

TLC	= Thin Layer Chromatography
UPM	= Universiti Putra Malaysia (formerly known as Universiti Pertanian Malaysia)
UV	= Ultraviolet
λ_{\max}	= Lambda max
w/w	= Weight per weight



Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfilment of the requirements for the degree of Master of Science.

**BIOACTIVITY OF *AZADIRACHTA EXCELSA* Jack.
EXTRACTIVES ON SELECTED INSECTS.**

By

MARIA LOURDES T. LARDIZABAL

April 1998

Chairman : Associate Professor Ahmad Said Sajap, Ph. D.

Faculty : Forestry

Azadirachta excelsa Jack. is a fast growing timber species that has been newly chosen as a potential forest plantation species to overcome timber deficit for domestic consumption in future. This species like *Azadirachta indica* is suspected to contain biologically active compounds that is detrimental on insects. Many studies have been done on the *A. indica*, a relative species to *A. excelsa*, however, studies on *A. excelsa* are still in preliminary stages. Therefore, the aim of this study was to evaluate the effect of *A. excelsa* crude extractives towards three selected insect pests, (i.e. *Spodoptera litura*, *Callosobruchus chinensis* and *Coptotermes curvignathus*). A preliminary test to evaluate the susceptibility of *A. excelsa* wood towards termites, compared to *Koompasia malaccensis*, *Hevea brasiliensis* and *Pinus caribaea* was also conducted.



In the first part of the study, the *A. excelsa* plant extracts were extracted from different plant parts (i.e. leaf, bark, trunk, twigs and branch) using a range of polar to non-polar solvents, namely hexane, petroleum ether, toluene, ethyl acetate and methanol. The plant extractives were sampled and analysed through Thin Layer Chromatography (TLC) and Infrared Spectroscopy (IR). In the second part of the study, bioassay tests were conducted on the selected insects. The insects were exposed to the plant extractives prepared at three concentration levels (i.e. 250 ppm, 500 ppm and 1000 ppm) and the solvents (0 ppm) as the control, by feeding the insects with treated food source. The *S. litura* larvae were fed with castor leaf discs soaked in the respective extractives. The *C. chinensis* adults were exposed to mungbean seeds that had been soaked in the plant extractives while the *C. curvignathus* were fed with treated filter paper. In the last part of this study, the *A. excelsa* wood together with the other three wood species mentioned earlier, were exposed to the termites both in the laboratory and field.

The TLC and IR results on the *A. excelsa* extractives showed that extractives from the different plant parts contained more non-polar chemical compounds. Among the five plant parts, the leaf contained the highest number of chemical compounds. The functional groups that were identified in the methanol and hexane extracts of leaf, bark and twigs, were hydroxyl, carboxyl, aromatic and unsaturated aldehydes.



The bioassay on *S. litura* showed that the larvae consumed less leaf discs that were treated with higher concentration of extractive. Mortality mostly occurred at the early larval stages. Although less mortality occurred at the lower concentration, but the moulting and larval periods were extended to twice the normal period and thus the larvae consumed more leaf discs. However, the growth parameters (i.e. head capsule measurement and pupal weight) at the last instar stage was not effected by the treatments. Generally, bark extract caused a higher mortality and less food consumption compared to the leaf extract.

The *A. excelsa* crude extract was not a good protectant on mungbean seeds. Results obtained from the treatments show that the oviposition rate and percentage adult emergence were not significant.

The termites bioassay test showed that *A. excelsa* extractives caused high termite mortality coupled with less consumption on filter paper treated with bark extracts. Choice feeding test on wood blocks from four different species showed that *A. excelsa* was the least preferred wood species compared to the other species, both in the field and laboratory trials.



Abstrak tesis yang dikemukakan kepada senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains.

**BIOAKTIVITI EKSTRAK TUMBUHAN *AZADIRACHTA EXCELSA* JACK.
TERHADAP SERANGGA-SERANGGA TERPILIH**

Oleh

MARIA LOURDES T. LARDIZABAL

April 1998

Pengerusi : Profesor Madya Ahmad Said Sajap, Ph. D.

Fakulti : Perhutanan

Azadirachta excelsa Jack. merupakan spesies pokok kayu cepat tumbuh yang baru dipilih sebagai spesies tanaman ladang hutan yang berpotensi untuk mengatasi masalah kekurangan bekalan kayu bagi kegunaan tempatan pada masa akan datang. Spesies ini seperti *Azadirachta indica*, disyaki mengandungi komponen biokimia aktif di dalam ekstrak tumbuhannya yang boleh mengawal serangga perosak. Banyak kajian telah dijalankan terhadap spesies *A. indica*, yang merupakan spesies serupa dengan *A. excelsa*. Walau bagaimanapun, kajian - kajian terhadap *A. excelsa* masih lagi pada peringkat awal. Tujuan kajian ini dijalankan adalah untuk menilai kesan ekstraktif mentah *A. excelsa* terhadap spesies serangga-serangga perosak terpilih, iaitu *Spodoptera litura*, *Callosobruchus chinensis* dan *Coptotermes curvignathus*. Kajian awal untuk menguji kerentanan kayu *A. excelsa* terhadap anai-anai, dibandingkan dengan *Koompasia malaccensis*, *Hevea brasiliensis* dan *Pinus caribaea* telah juga dijalankan.



Pada awal kajian ini, ekstrakatif tumbuhan *A. excelsa* telah diekstrak daripada bahagian-bahagian tumbuhan yang berlainan (i.e. daun, kulit kayu, batang kayu, ranting kayu dan cabang), dengan menggunakan pelarut julat polar hingga tidak polar. Antara pelarut yang digunakan adalah heksana, petroleum ether, toluene, etil asetat, dan metanol. Ekstraktif tumbuhan tersebut telah disampel dan dianalisa melalui “Thin Layer Chromatography” (TLC) dan “Infrared Spectroscopy” (IR). Dalam bahagian kedua kajian ini, ujian “bioassay” telah dijalankan terhadap serangga-serangga terpilih. Serangga-serangga tersebut telah didedahkan kepada ekstrakatif tumbuhan yang telah dilarutkan kepada tiga kepekatan berbeza (i.e. 250 ppm, 500 ppm dan 1000 ppm) serta pelarut yang digunakan semasa ekstraksi sebagai kawalan (0 ppm), dengan memberikan makanan yang telah dirawat. Larva *S. litura* telah diberikan kepingan daun pokok jarah yang telah direndamkan ke dalam ekstrakatif berkenaan. Serangga dewasa *C. chinensis* telah didedahkan kepada biji kacang hijau yang telah dirawat dengan ekstrakatif tumbuhan *A. excelsa*, manakala *C. curvignathus* telah diberikan kertas turas yang telah dirawat sebagai sumber makanan. Pada bahagian terakhir kajian ini, kayu *A. excelsa* bersama dengan spesies kayu yang telah disebutkan lebih awal telah didedahkan kepada anai-anai di lapangan dan juga di makmal.

Keputusan analisa TLC dan IR terhadap ekstrakatif *A. excelsa* menunjukkan bahawa ekstrakatif daripada bahagian tumbuhan yang dikaji mengandungi lebih

banyak kompaun kimia yang tidak polar. Di antara kelima-lima bahagian tumbuhan, bahagian daun mengandungi bilangan komponen kimia yang paling banyak. Kumpulan berfungsi yang telah dikenalpasti di dalam ekstrak metanol dan heksana daripada daun, kulit kayu dan ranting adalah hidroksil, karboksil serta aldehid aromatik dan tidak tepu.

Ujian “bioassay” terhadap *S. litura* menunjukkan bahawa jumlah pemakanan daun bagi larva serangga ini kurang pada daun yang telah dirawat dengan ekstrak pada kepekatan yang paling tinggi. Sungguhpun kematian larva rendah pada kepekatan yang paling rendah, namun jangkamasa pengelongsongan dan jangkamasa larva telah diunjurkan dua kali ganda lebih panjang daripada tempoh biasa. Oleh yang demikian, larva tersebut terus memakan daun yang diberikan. Walau bagaimanapun, ekstrak tersebut tidak memberi kesan signifikan terhadap berat pupa dan juga saiz kulit kepala pada peringkat larva terakhir. Pada keseluruhannya, ekstrak kulit kayu menyebabkan kematian yang lebih tinggi dan pemakanan daun yang kurang berbanding dengan rawatan ekstrak daun.

Ekstraktif mentah *A. excelsa* bukanlah pelindung biji kacang hijau yang baik. Keputusan yang diperolehi menunjukkan bahawa kadar peneluran dan peratus dewasa yang keluar adalah tidak signifikan.

Ujian “bioassay” terhadap anai-anai menunjukkan bahawa ekstrakatif *A. excelsa* telah menyebabkan kematian anai-anai yang tinggi dan juga pemakanan yang rendah ke atas kertas turas yang dirawat dengan ekstrak kulit kayu . Kajian pemakanan pilihan (Choice-feeding Test) ke atas empat spesis kayu yang telah diberikan menunjukkan bahawa kayu *A. excelsa* adalah spesis yang kurang diminati berbanding dengan spesis kayu yang lain. Keputusan yang sama telah diperolehi dalam ujian lapangan dan juga makmal.



CHAPTER I

GENERAL INTRODUCTION

Many insect species are beneficial to human beings, especially as pollinators, source of food and also source of medicine. However, when insects compete with man for food and organic resources, while others spread diseases or cause discomfort they become pests. Particularly in agriculture and forestry, these insects constitute an important constraint to crop and timber production. Over the years, means have been developed to reduce drastically the harmful effects of these insect pests. To a large extent, this process has been achieved through trial and error, including most current control methods which are for the most part empirical.

Therefore, to alleviate these pest problems, extensive attempts were made using synthetic pesticides in order to control these insect pests. In Malaysia, for example, chemical insecticides have been widely used in combating insect pest outbreaks.



However, owing to the extensive use of chemical and synthetic insecticides, various drawbacks have been observed in man and the environment. These include toxicity to non - target organisms, development of pest resistance and environmental degradation (Schmutterer 1988, Anon.,1991). In developing countries, inadequate product knowledge, supply uncertainties and high prices cause inefficient pesticide use and also create additional socioeconomic problems between the rich and poor. Consequently, the search for new, preferably environmentally sound, effective and low-cost insecticides are thus needed. Due to this matter, interest in plant-based insecticides has grown rapidly during recent years.

Plant-based pesticides offer a safer, environmentally friendly and effective alternative to synthetic insecticides. Phytochemical extracts from the neem tree, *Azadirachta indica* A. Juss for instance, have been extensively studied in the last decade. They have shown several properties useful for the management of insect pests. These include repellency, feeding and oviposition deterrence, insect growth regulatory activity, low mammalian toxicity, and minimal persistence in the environment (reviewed in Schmutterer 1990, Koul and Isman 1991, Ascher 1993, Mordue and Blackwell 1993). Neem may also have less impact on natural enemies than most conventional insecticides (Hoelmer *et al.* 1990, Stark 1992, McCloskey *et al.* 1993, Lowery and Isman 1994). Larval feeding inhibition and insect growth regulation are due primarily to the tetranortriterpenoid compound azadirachtin (Isman *et al.* 1990), which is found in great concentrations in the seed kernels.