



UNIVERSITI PUTRA MALAYSIA

**FERTILIZER REQUIREMENTS AND NUTRIENT DYNAMICS
OF TEAK (TECTONA GRANDIS L.F.) PLANTATIONS
IN PENINSULAR MALAYSIA**

MUHAMMAD TAHIR SIDDIQUI

FH 1998 10

**FERTILIZER REQUIREMENTS AND NUTRIENT DYNAMICS
OF TEAK (*TECTONA GRANDIS* L.f.) PLANTATIONS
IN PENINSULAR MALAYSIA**

MUHAMMAD TAHIR SIDDIQUI

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA
1998**



**FERTILIZER REQUIREMENTS AND NUTRIENT DYNAMICS
OF TEAK (*TECTONA GRANDIS* L.f.) PLANTATIONS
IN PENINSULAR MALAYSIA**

BY

MUHAMMAD TAHIR SIDDIQUI

**Dissertation Submitted in Fulfilment of the Requirements for the
Degree of Doctor of Philosophy in the Faculty of Forestry
Universiti Putra Malaysia.**

February 1998



Dedicated to My Beloved Father

(Late) **HAJI MUHAMMAD ISHAQUE SIDDIQUI**

whose dream comes true.



ACKNOWLEDGEMENTS

All praise to Allah S.W.T whose countless blessings enabled me to accomplish this study. I wish to express my deep gratitude to my supervisor Assoc. Prof. Dr. Haji Sheikh Ali Abod for his unflinching interest, concern and inspiration throughout the period of this study. His discreet assistance of many kinds has helped me to steer uphill, especially during the write-up. I gratefully acknowledge my indebtedness to my co-supervisors Prof. Dr. Azizah Hashim and Dr. Amir Husni Mohd. Shariff who furnished invaluable comments and suggestions which significantly improved the thesis. Thanks are also due to Dr. Mohammad Idrees Ahmad for his help in statistical analyses and my friends for their help and encouragement.

I take this opportunity to express my thanks to Forest Research Institute Malaysia (FRIM) for the permission given to conduct the study in its sub-station at Mata Ayer, Perlis and also to the field staff for their help in the field work and data collection.

I gratefully acknowledge the financial support from the Public Services Department Malaysia and University of Agriculture, Faisalabad- Pakistan for granting study leave.



Special thanks are due to my brothers, sisters and relatives for their encouragement and support throughout the period of this study. I would like to express my profound appreciation to my wife Shagufta Shaheen and my lovely daughters Wajeeha Siddiqui and Zanira Siddiqui for their sacrifices, patience and moral support that enabled me to successfully carry out this research.



TABLE OF CONTENTS

		Page
ACKNOWLEDGEMENTS.....		iii
LIST OF TABLES		viii
LIST OF FIGURES.....		xi
ABSTRACT		xiv
ABSTAK		xviii
CHAPTER		
I	INTRODUCTION	1
	Objectives	5
	Teak (<i>Tectona grandis</i>) - A Profile	6
	Distribution.....	6
	Physiognomy	7
	Ecological Requirements.....	7
	Silvicultural Aspects	8
	Properties and Uses	9
II	LITERATURE REVIEW.....	11
	Plantation Forestry - A Vivid Strategy	11
	Sustainable Plantation Forestry	16
	Morphology and Physiology of Teak	18
	Nutrient Deficiencies Diagnosis	20
	Visual Symptoms	21
	Plant Tissue Analysis	24
	Soil Analysis	25
	Biological Assays	26
	Forest Fertilization	26
	General Principles and Practices	26
	Rates of Fertilizer Application	28
	Frequency of Fertilizer Application	30
	Time of Application of Fertilizers.....	31
	Fertilization and Stand Development	32
	Nutrient Dynamics within Tree	33
	Teak Nutrition Research - An Overview.....	39
	Forest Fertilization Research in Malaysia	41

III	POT TRIALS OF <i>T. GRANDIS</i> SEEDLINGS	43
	Introduction	
	Materials and Methods	44
	Potting Medium	44
	Seedlings	44
	Experimental Design and Treatments	45
	Measurements for Growth and Nutrient Element	
	Parameters and Data Analysis	46
	Results	48
	Statistical Analysis	48
	Height Increment	52
	Diameter Increment	52
	Root Length	55
	Total Leaf Area	55
	Dry Matter Production	57
	Discussion	60
IV	FIELD TRIALS OF <i>T. GRANDIS</i> PLANTATIONS	67
	Introduction	67
	Materials and Methods	68
	General Description of the Experimental Site	68
	Experimental Design and Treatments	71
	Growth Measurements and Data Analysis	73
	Results	74
	Height Increment	74
	Diameter Increment	74
	Dry Matter Production	78
	Discussion	79
V	FOLIAR NUTRIENT CONCENTRATIONS IN	
	<i>T. GRANDIS</i> PLANTATIONS	82
	Introduction	82
	Materials and Methods	83
	Results	86
	Relationship between Foliar Nutrient	
	Concentrations and Growth	86
	Fertilizer Effects	88
	Effect of Chronosequence and Crown Sampling Positions	
	on Foliar Nutrient Concentrations	89
	Chronosequence	89



	Crown Strata	91
	Nitrogen (N)	91
	Phosphorus(P)	92
	Potassium (K)	92
	Calcium (Ca)	94
	Magnesium (Mg)	94
	Zinc (Zn)	94
	Manganese (Mn)	96
	Copper (Cu)	96
	Effect of Crown Strata on Nutrient Variability	98
	Discussion	102
VI	NUTRIENT CONCENTRATIONS IN STEMS OF TEAK (<i>T. GRANDIS</i>) GROWN IN PLANTATIONS	110
	Introduction	110
	Materials and Methods	111
	Results	112
	Fertilizer Effects	112
	Chronosequence Effect on Nutrient Concentrations in Stem Components	114
	Relationship between Nutrient Concentrations in Stem Components and Growth	119
	Discussion	
VII	ROOT NUTRIENT CONCENTRATIONS IN TEAK (<i>T. GRANDIS</i>) PLANTATIONS	129
	Introduction	129
	Materials and Methods	130
	Results	132
	Discussion	138
VIII	GENERAL DISCUSSION	143
IX	SUMMARY, CONCLUSIONS AND RECOMMENDATIONS	153
	BIBLIOGRAPHY	157
	APPENDIX- Additional Figures	180
	VITA	185



LIST OF TABLES

Table	Page
1 Areas of Plantations Including Woodlots in the Tropics and Hotter Subtropics by Continents (000s ha)	13
2 Mean Annual Increments from Tropical Plantations	14
3 Comparison Between Productivity of Planted Forests and Indigenous Forests	15
4 Chemical Properties of the Soil Sand Mix	44
5 Levels of Nutrients Applied	46
6 Fertilizer Treatments	46
7 Results of Analyses of Variance on Teak Growth Plantations	50
8 Comparison Between Treatments for Different Growth Parameters of <i>T. grandis</i> Seedlings after 12 Months of Fertilizer Application	51
9 Levels of Nutrients Applied	72
10 Fertilizer Treatments	72
11 Comparison between Treatments and Growth Parameters of <i>T. grandis</i> Seedlings 12 Months after Fertilizer Application	77
12 Analysis of Variance for Height and Diameter of Teak Seedlings in the Field	77
13 Detail of Trees Sampled for Foliar Analysis	84
14 Coefficient of Correlation between Foliar Nutrient Concentrations and Tree Growth	87
15 Stepwise Multiple Regression between Growth Parameters and Foliar Nutrient Concentrations	88



16	Effect of Fertilizers on Foliar Nutrient Concentrations in 1 and 2-year-old Teak Seedlings	89
17	Foliar Nutrient Concentrations (% dry wt.) in an Age Series of Un-fertilized <i>T. grandis</i> Plantations	90
18	Nutrient Concentrations in Different Crown Positions in Different Ages (year)	91
19	Coefficient of Variation of Foliar Nutrient Concentrations in Un-fertilized Trees at Various Crown Positions and Ages	99
20	Analysis of Variance (F-values) on Crown Positions and Ages of Teak Trees	102
21	Reported Foliar Nutrient Concentrations (% oven dry weight) in Mature Leaves of <i>T. grandis</i> Taken from the Upper Part of the Crown of Dominant Trees	108
22	Effect of Fertilizer Application on Various Compartments of Teak Stem	113
23	Nutrient Concentrations in Sapwood and Heartwood of Teak Trees	119
24	Correlation Coefficient for Stemwood Nutrients and Stembark Nutrients with Diameter and Age Trees	120
25	Multiple Regression for Estimating the Nutrient Concentrations in Stemwood from Diameter and Tree Age	121
26	Multiple Regression for Estimating the Nutrient Concentrations in Stembark from Diameter and Tree Age	122
27	Comparison of Nutrient Concentrations in Stem Components of Different Aged Teak Trees	127
28	Anova of Main Effect of Age, Treatment and Diameter and Their Interactions on Nutrient Concentrations in Roots of 1 and 2-year-old Teak Trees	133



29	Main Effect of Age, Fertilizer and Diameter on the Nutrient Concentrations in Roots of Teak Trees	134
30	Effect of Fertilizer Application on Nutrient Concentration in Roots of 1 and 2-year-old Teak Trees	135
31	Anova of Main Effects of Age, Diameter and their Interactions on the Nutrient Concentrations in Roots of Unfertilized Teak Trees	135
32	Main Effects of Age on the Nutrient Concentrations in Roots of Unfertilized Teak Trees	135
33	Main Effect of Diameter on the Nutrient Concentrations in Roots of Unfertilized Teak Trees	136
34	Characteristics of Penambang Series Soils at Mata Ayer, Perlis	136
35	Soil Properties under Different Aged Teak Plantations	137



LIST OF FIGURE

Figure		Page
1	Main Effects of Different Levels of N, P, K on Height Increment after 12 Months	53
2	Main Effects of Different Levels of N on Height Growth Monitored at 2 Monthly Intervals for 12 Months	53
3	Height Growth at Different Levels of P Monitored at 2 Monthly Intervals for 12 Months	53
4	Height Growth at Different Levels of K Monitored at 2 Monthly Intervals for 12 Months	53
5	Effects of NP Interaction on Height Increment after 12 Months	53
6	Effects of NK Interaction on Height Increment after 12 Months	53
7	Effects of PK Interaction on Height Increment after 12 Months After Treatment	54
8	Main Effects of Different Levels of N, P, K on Diameter Increment 12 Months After Treatment	54
9	Main Effects of Different Levels of N on Diameter Monitored at 2 Monthly Intervals 12 Months after Treatment	54
10	Main Effect of Different Levels of P on Diameter Monitored at 2 Monthly Intervals 12 Months after Treatment	54
11	Main Effects of Different Levels of K Monitored at 2 Monthly Intervals 12 Months after Treatment	54
12	Effects of NP Interaction on Diameter Increment 12 Months after Treatment	54
13	Effects of NK Interaction on Diameter Increment after 12 Months	56



14	Effects of PK Interaction on Diameter Increment after 12 Months	56
15	Effects of Different Levels of N, P, K on Root Length	56
16	Effects of Different Levels of N, P, K on Leaf Area	56
17	Effects of Different Levels of N, P, K on the Total Plant Weight 12 Months after Treatment	59
18	Effect of N, P, K Fertilizer on Leaf Weight 12 Months after Treatment	59
19	Effects of Different Levels of N, P, K on Shoot Weight	59
20	Effect of N, P, K Fertilizer on Root Weight	59
21	Effects of NPK Fertilizer on Root-shoot Ratio	59
22	Lay-out Plan of the Experiment	73
23	Effects of Combined NP Fertilizer on Height Increment Monitored 12 Months after Treatment	75
24	Height Growth at Different Levels of N	75
25	Height Growth at Different Levels of P	75
26	Effect of Combined NP Fertilizer on Diameter Increment Monitored 12 Months after Treatment	76
27	Diameter Increment at Different Levels of N Fertilizer	76
28	Diameter Increment at Different Levels of P	76
29	Effects of Sample Position and Age on Foliar N Concentration	93
30	Effects of Sample Position and Age on Foliar P Concentration	93
31	Effects of Sample Position and Age on Foliar K Concentration	93



32	Effect of Sample Position and Age on the Foliar Ca Concentration	95
33	Effect of Sample Position and Age on Foliar Mg Concentration	95
34	Effect of Sample Position and Age on Foliar Zn Concentration	95
35	Effect of Sample Position and Age on the Foliar Mn	97
36	Effect of Sample Position and Age on Foliar Cu Concentration	97
37	Effect of Age on Nutrient Concentration of Bark of Teak Trees	115
38	Effect of Age on Nutrient Concentrations of Stembark in Teak Trees . . .	115
39	Effect of Age on the Nutrient Concentrations in Twigs of Teak Trees . . .	117
40	Effect of Age on the Nutrient Concentrations in Twigs of Teak Trees . . .	117
41	Effect of Age on Nutrient Concentrations in Branches of Teak Trees . . .	117
42	Effect of Age on Nutrient Concentrations in Branches of Teak Trees . . .	118
43	Effect of Age on the Nutrient Concentrations in Stemwood of Teak Trees	118
44	Effect of Age on Nutrient Concentrations in Stemwood of Teak Trees . .	118
45	Diagram Showing Sub-divisions of Sampling Zone for Roots	131
46	Map of Peninsular Malaysia Showing the Experimental Area	181
47	Rainfall Data Recorded During Study Period from July, 1995 to June, 1996 at Chuping Meterological Station near Experimental Site	183
48	Temperature Data Recorded During Study Period from July, 1995 to June, 1996 at Chuping Meterological Station near Experimental Site	184



LIST OF PLATES

Plate		Page
1	Foliar Sample Positions in the Crown	85
2	<i>Tectona grandis</i> Seedlings in the Greenhouse	182



ABSTRACT

Abstract of thesis submitted to the Senate of the Universiti Putra Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy.

FERTILIZER REQUIREMENTS AND NUTRIENT DYNAMICS OF TEAK (*TECTONA GRANDIS* L.f.) PLANTATIONS IN PENINSULAR MALAYSIA

By

MUHAMMAD TAHIR SIDDIQUI

February, 1998

Chairman: Associate Professor Dr. Haji Sheikh Ali Abod

Faculty: Forestry

Teak (*Tectona grandis* L.f) is one of the highly sought after timbers of the world. It is not indigenous to Malaysia and was introduced as plantation species in the northern states of Kedah and Perlis in early 1950's. To-date, 1100 ha of teak plantation has been established in Malaysia and more acreage are planned in the future. However, comprehensive fertilizer requirements of this species are not adequately known.

The objective of the present study was to determine the fertilizer requirements of *T. grandis* and their effects on the nutrient dynamics in the trees. Three



experiments were conducted for this purpose: pot trial, field trial and destructive sampling of various aged trees. Soil and plant analyses were carried out to determine the effects of fertilizer application and age on the nutrient concentrations in the trees. Results of the pot trial revealed that 564 kg/ha of ammonium sulphate and 300 kg/ha of P_2O_5 with 75 kg/ha of K_2O promoted the growth of *T. grandis* seedlings. The results of field experiment clearly indicated that N and P are equally important and act additively to boost the growth of teak seedlings. It was found that 200 kg/ha ammonium sulphate and 300 kg/ha triple superphosphate gave maximum height and diameter of the teak seedlings. The disparity in fertilizer dosages is due to large number of plants per ha in pot experiment.

Application of fertilizers increased the N, P, K, Ca, Mn and Zn concentrations in the leaves of teak trees. Magnesium and Cu concentrations were however, reduced as a result of fertilizer application. This is attributed as antagonism process. Nutrient concentrations i.e. N, P and K in the leaves reduced as the tree advanced in age whereas Ca, Mg, Mn, Zn and Cu concentrations increased with increase in tree age.

The foliar sampling experiment revealed that nutrient concentrations were influenced by the position in the crown. On the basis of low nutrient variability, the results showed that sampling should be carried out in the morning and from the upper or middle crown for N, P, K, and from the lower crown for Ca, Mg, Mn and Cu.

Stem analysis showed that there was lowest nutrient concentrations in the wood especially in the heartwood. Bark contained maximum concentrations of Ca. Fertilizer application increased N, P, K and Ca concentrations in the bark and stemwood. In contrast, Mg and Zn concentrations decreased as a result of fertilizer application. Nitrogen concentration decreased in twigs of fertilized plants. There was a decreasing trend in nutrient concentrations due to increase in age except for Ca which increases with age.

Nutrient concentrations in teak roots were also influenced by fertilizer additions. Macro nutrients were high in fine roots (<2mm) except Mg which was high in control plants. Manganese and Zn also increased as a result of fertilizer addition irrespective of root size. The results also revealed that nutrient concentrations decreased with increase in root diameter class and age.

The results of the present study revealed that the combination of pot and field experiments, analysis of tree components i.e. foliar, stem and root and soil analysis could be a useful technique in determining the fertilizer requirements of *T. grandis* in Peninsular Malaysia. The nutrient concentrations as affected by fertilizer additions and age should also serve as a guideline for evaluating the nutritional status of other quality timber species in the tropics.

ABSTRAK

Abstrak tesis dikemukakan kepada Senat Univesiti Putra Malaysia bagi memenuhi keperluan Ijazah Doktor Falsafah.

KEPERLUAN BAJA DAN DINAMIK NUTRIEN PADA JATI (*TECTONA GRANDIS* L.f.) DI SEMENANJUNG MALAYSIA

Oleh

MUHAMMAD TAHIR SIDDIQUI

Februari, 1998

Pengerusi: Madya Professor Dr.Hj. Sheikh Ali Abod

Fakulti: Perhutanan

Jati (*Tectona grandis* L.f.) merupakan salah satu kayu balak yang sangat penting di dunia. Ia bukan spesies yang asli di Malaysia tetapi telah diperkenalkan sebagai spesies perladangan di negeri-negeri di bahagian utara semenanjung iaitu Kedah dan Perlis pada awal tahun 1950 an. Sehingga kini, terdapat 1100 ha. ladang jati di Malaysia dan keluasan perladangan ini akan diperbesarkan lagi pada masa depan. Walau bagaimanapun, penggunaan baja yang paling sesuai untuk spesies jati belum diketahui lagi.

Objektif kajian ini adalah untuk menentukan baja yang diperlukan oleh *Tectona grandis* dan kesannya ke atas dinamik nutrien di dalam pokok. Tiga eksperimen telah dijalankan iaitu percubaan berpasu, perenbaan di ladang dan persampelan destruktif ke atas pokok pelbagai umur. Analisis tanah dan tanaman telah dijalankan untuk



menentukan kesan pemberian baja dan umur ke atas konsentrasi nutrien di dalam pokok. Keputusan dari percubaan berpasu menunjukkan bahawa 564 kg/ha ammonium sulfat dan 300 kg/ha P_2O_5 dengan 75 kg/ha K_2O menggalakkan pertumbuhan anak benih *T. grandis*. Keputusan dari percubaan ladang menunjukkan dengan jelas bahawa N dan P adalah sama penting dan bertindak secara sinergistik untuk menggalakkan lagi pertumbuhan anak benih pokok jati Ia juga menunjukkan bahawa 200 kg/ha ammonium sulfat dan 300 kg/ha triple superphosphate memberikan ketinggian dan perepang batang yang maksimum kepada anak benih pokok jati.

Pemberian baja meningkatkan kadar N, P, K, Ca, Mg, Mn dan Zn pada daun pokok jati. Walau bagaimanapun, konsentrasi Mg dan Cu berkurangan dengan pemberian baja. Ini adalah disebabkan berlakunya proses antagonisma. Konsentrasi nutrien, contohnya N, P dan K pada daun didapati berkurangan apabila umur pokok meningkat manakala Ca, Mg, Mn, Zn dan Cu pula bertambah.

Ujikaji persampelan daun telah menunjukkan bahawa konsentrasi nutrien dipengaruhi oleh kedudukan daun pada silara pokok. Berasaskan kepada perubahan nutrien yang rendah, keputusan kajian menunjukkan bahawa persampelan hendaklah dibuat pada awal pagi dan sampel diambil dari bahagian atas atau tengah silara pokok untuk analisis N, P, K, dan dari bahagian bawah silara pokok untuk analisis Ca, Mg, Mn dan Cu.



Analisis batang menunjukkan bahawa kayu jati terutamanya di bahagian kayu terasnya mempunyai konsentrasi nutrien yang rendah. Konsentrasi Ca yang maksimum didapati pada bahagian kulit. Pemberian baja telah meningkatkan konsentrasi N, P, K dan Ca pada kulit dan batang kayu. Sebaliknya, konsentrasi Mg dan Zn berkurangan dengan pemberian baja. Konsentrasi nitrogen adalah rendah pada ranting pokok yang telah dibaja. Keputusan turut menunjukkan pengurangan konsentrasi nutrien dengan pertambahan umur pokok kecuali unsur Ca yang didapati meningkat dengan bertambahnya umur pokok.

Konsentrasi nutrien pada akar jati juga dipengaruhi oleh pemberian baja. Nutrien makro adalah tinggi pada akar halus (< 2 mm) kecuali unsur Mg yang didapati rendah pada pokok kawalan. Pemberian baja dengan tidak mengambil kira saiz akar juga telah meningkatkan unsur Mn dan Zn dalam akar. Keputusan yang diperolehi juga menunjukkan bahawa konsentrasi nutrien berkurangan dengan bertambahnya kelas diameter akar dan umur pokok.

Ketiga-tiga kajian yang telah dijalankan boleh digunakan sebagai satu garis panduan yang baik untuk menentukan paras baja yang diperlukan oleh *T. grandis* di Semenanjung Malaysia. Konsentrasi nutrien yang dipengaruhi oleh pemberian baja dan umur pokok boleh digunakan sebagai panduan untuk menilai status nutrien pada spesis balak bermutu di kawasan tropika.

CHAPTER I

INTRODUCTION

Malaysia is blessed with rich natural resources which have brought wealth and prosperity to this nation. The country is recognized as one of the world's leading exporters of tropical hardwoods. The total forest area in Malaysia is 19.07 million hectares about 58.1 percent of the total land mass of the country. Out of this, 6.19 million hectares are in Peninsular Malaysia whereas 4.44 million hectares and 8.44 million hectares are in Sabah and Sarawak, respectively (Ministry of Primary Industries, 1996). However, the World Bank has estimated that the tropical forest is fast disappearing at a rate of 15-20 million ha per year. At the level of demand prevailing then, the remaining tropical forest would disappear in 60 to 80 years (Evans, 1992). The most extreme pressures lie in countries in the tropics. Malaysia is not an exception in this case.

In Peninsular Malaysia, approximately 360,121 ha of forested lands are being cleared annually in the recent years. Seventy-five percent of the total forest lands cleared were alienated for agricultural and industrial development (Anon, 1979). In addition, there is also a general increase in demand for wood, paper and paper products. The production pattern however, does not seem to keep pace with this timber demand. It is



anticipated that Peninsular Malaysia will experience an acute shortage of timber well before the year 2000 (Freezaillah, 1982)

The present precarious forest situation in Malaysia stems from an apparently nonchalant forest management practice in the past which was influenced by the misconstrued belief that Malaysia's forest resources were inexhaustible. Hence, from the early 1960s onwards, the green gold of Malaysia became the target of exploitation. Several strategies are therefore under way to ensure that this renewable resource is being perpetuated at sustainable levels, one of which is to raise large scale plantations of suitable exotic and indigenous species. It is therefore, safe to conclude that plantation forestry will play an important role in the future. The growing demand for wood, hampered by the low productivity of natural forests, has greatly enhanced the role of plantation forestry. Plantation forestry also enhances the creation of resources to meet the demand for wood and wood products, development of a flexible resource able to yield different kind and size of products for both internal and external markets. Intensive reforestation programme is currently being carried out in P. Malaysia under a programme of Compensatory Forest Plantation (Yong, 1984). However, this project purely stresses on the supply of medium quality timber grown on short rotation to meet the basic requirement of the lower income group. So far there has been no large scale programme to grow high quality timber species which is also expected to be in short supply in the near future (Hashimand Zainudin, 1983).

One of the current high quality timber species greatly appreciated by Malaysians is teak (*Tectona grandis* L.f). Teak occupies a dominant position in the highly sought after timbers in the world. This species originates from countries with tropical monsoon climates, chiefly India, Myanmar, Thailand and Cambodia. The annual pronounced dry season followed by wet weather in these countries concerned imparts a beautiful grain structure to teak (Borota, 1991). At the same time teak wood is characteristically strong, durable and easily workable. Teak rules the world market because of its sterling qualities. It is difficult to find a happy blend of beauty, strength, stability and durability in any other tropical timber. Rightly teak wood is considered to be the best general utility timber with worldwide reputation, being extensively used for ship building, bridges and wharves, railway carriages and wagons, ordnance, shingles, wheels, carving and general carpentry (Appanah and Weinland, 1993). As a result, it is much sought after throughout the world and fetches a high price as compared to other species of wood.

Currently, all teak utilized in Malaysia is imported from Myanmar, Thailand and Indonesia. Owing to a decline in supply from the countries of origin, the price has spiraled over the years. This situation is expected to continue, unless efforts are made to establish large scale plantations of teak in Malaysia.

Teak has been reported to require very specific soil and climatic conditions for optimum growth. It needs well- drained and well- aerated deep alluvial soils for good growth. Griffith and Gupta (1947) observed that a forest of fine teak growth changed to