



UNIVERSITI PUTRA MALAYSIA

**PHENOLOGY AND AVAILABILITY OF FRUIT TREES AND THEIR
INFLUENCE ON THE ABUNDANCE OF SELECTED ANIMALS
IN LOGGED AND PRIMARY FOREST OF SUNGAI LALANG
FOREST RESERVE, SELANGOR**

ROMEO M. LOMOLJO

FH 2003 18

**PHENOLOGY AND AVAILABILITY OF FRUIT TREES AND THEIR
INFLUENCE ON THE ABUNDANCE OF SELECTED ANIMALS
IN LOGGED AND PRIMARY FOREST OF SUNGAI LALANG
FOREST RESERVE, SELANGOR**

By

ROMEO M. LOMOLJO

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of Requirements for the
Degree of Master Science**

October 2003



DEDICATION

This Piece of Work is dedicated to; my late father Jorge C. Lomoljo

And Mother Anicita M. Lomoljo and to my Dearest

Wife Eva and Children Maruxa Linda

And Marcella MIA

**Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
Fulfilment of the requirements for the degree of Master of Science**

**PHENOLOGY AND AVAILABILITY OF FRUIT TREES AND THEIR
INFLUENCE ON THE ABUNDANCE OF SELECTED ANIMALS
IN LOGGED AND PRIMARY FOREST OF SUNGAI LALANG
FOREST RESERVE, SELANGOR**

By

ROMEO M. LOMOLJO

October 2003

Chairman: Associate Professor Mohamed Zakaria Hussin, Ph.D.

Faculty: Forestry

The abundance of fruit tree species as food source for wildlife in logged and primary forest was evaluated. The phenological data collection was carried out from September 1999 to October 2000 in Sungai Lalang Forest Reserve, Semenyih, Selangor, Malaysia. The general objective of this study is to determine the fruit tree species that serves as food source for wildlife in logged and primary forest. The specific objectives are: to compare the availability of fruits to animals in three different compartments. To compare fruit trees distribution in three compartments within block. And to correlates the abundance of animals in relation to food availability. Three different Compartments were selected namely: Compartment 24 (VJR), Compartment 33 (10-year-old logged forest) and Compartment 18 (5-year-old logged forest). Within each Compartment,

three blocks were established in different location (e.g. ridge top, mid-slope and valley bottom). All trees bigger than 10 cm dbh were tagged and identified and monitored every month for leafing, flowering and fruiting activities. The Duncan's test indicated that the leafing pattern in Compartment 24 (VJR) was significantly different ($P < 0.05$) from Compartment 33 (10-year-old logged forest) and Compartment 18 (5-year-old logged forest) ($P < 0.05$). The flowering pattern however, showed no significant difference among the three different Compartments. The percentage of fruiting was significantly lower in Compartment 18 (5-year-old logged forest) than in the primary forest (VJR) ($P < 0.05$).

In general, the availability of the food sources such as leaves, flowers and fruits was almost similar in primary and the two-logged forest, however only the primary forest (VJR) tends to provide more food to wildlife. This study showed that less trees or fruit trees meant less food; likewise more fruit trees meant more food. This indicated that the survival of the animals in the forest especially the logged forest fully depended on the fruit trees left after the logging activities. Logging activity influenced the distribution and availability of food sources and it is frequently correlated with the behaviour pattern of animal species.

**Abstrak thesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains**

**FENOLOGI DAN KEDAPATAN POKOK BERBUAH DAN PENGARUH
KEATAS LIMPAHAN HAWAN TERPILIH DALAM HUTAN SELEPAS
PEMBALAKAN DAN HUTAN ASLI HUTAN SIMPAN SUNGAI
LALANG, SELANGOR**

Oleh

ROMEO M. LOMOLJO

October 2003

Pengerusi: Professor Madya Mohamed Zakaria Hussin, Ph.D.

Fakulti: Perhutanan

Sejumlah spesis pokok berbuah sebagai sumber makanan untuk hidupan liar di hutan selepas dibalok dan hutan asli telah dinilai. Pengumpulan data fenologi telah dijalankan daripada September 1999 hingga Oktober 2000 di Hutan Simpan Sungai Lalang, Semenyih, Selangor, Malaysia. Oleh yang demikian objektif umum kajian ini adalah untuk mengenal spesis pokok berbuah yang merupakan sumber makanan untuk hidupan liar di hutan selepas dibalok dan di hutan asli. Objektif utama adalah untuk membuat perbandingan jumlah buah yang terdapat dengan kehadiran haiwan di tiga kompartmen yang berbeza, yang dinamakan; Kompartmen 24 (VJR), Kompartmen 33 (10 tahun selepas pembalakan) dan Kompartmen 18 (5 tahun selepas pembalakan). Pada setiap kompartmen, tiga blok telah dibuat di lokasi yang berbeza (cth: Cerun atas, Lereng tengah dan Lembah). Kesemua pokok lebih besar daripada 5 cm dbh

akan dikenalpasti, ditag, dicamkan dan dinilai setiap bulan samada dari segi jumlah daun, pembungaan dan pengeluaran buah. Keputusan menunjukkan bahawa corak pengeluaran daun di dalam Kompatmen 24 (VJR) perbezaan bererti ($P < 0,05$) berbanding dengan Kompatmen 33 (hutan 10-tahun selepas pembalakkan) dan Kompatmen 18 (hutan 5-tahun selepas penbalakkan) ($P < 0,05$). Walaubagaimanapun, corak pembungaan tidak menunjukkan perbezaan yang ketara di antara ketiga – tiga Kompatmen. Peratusan pengeluaran buah meneunjukkan perbezaan bererti di Kompatmen 18 (hutan 5- tahun selepas pembalakkan) berbanding dengan hutan primer (VJR) ($P < 0.05$). Secara amnya, kehadiran sumber – sumber makanan seperti daun, bunga dan buah adalah hampir sama dengan hutan primer dan kedua – dua hutan yang telah dibalak, walaubagaimanapun hanya hutan primer (VJR) yang menyumbangkan lebih makanan kepada hidupan liar.

Kajian ini menunjukkan bahawa kurangnya pokok – pokok atau pokok buah bermakna kurang makanan, dan sebaliknya lebih pokok buah, lebih makanan. Ini menunjukkan bahawa kemandirian hidupan di dalam hutan terutamanya hutan yang telah dibaik adalah bergantung sepenuhnya kepada pokok buah – buahan yang tinggal selepas aktiviti pembalakkan. Kegiatan pembalakkan mempengaruhi dan kalempahan sumber- sumber makanan dan ia biasanya dihubungkan dengan corak kelakuan sepsis hidupan.

ACKNOWLEDGEMENTS

My sincere appreciation is expressed to my supervisor Assoc. Prof. Dr. Mohamed Zakaria Hussin for the knowledge and insights he generously gave during the entire course of the study. My deepest gratitude is also to Assoc. Prof. Dr. Faridah Hanum Ibrahim for her invaluable encouragement and support. My special thanks also goes to Dr. Abdullah Mohd. and Pn. Khamzia Abdul Kudus for their useful comments, encouragements and suggestions.

Special gratitude is also due to Intensified Research Priority Areas (IRPA) through Associate Professor Mohamed Zakaria Hussin for providing research grant during my study. To SEAMEO SEARCA for giving me support to complete my thesis writing. Sincere appreciation is due to the botanist from FRIM Mr. Kamarudin Salleh for helping me in identifying the fruit trees in the three compartments, to our project coordinator En. Abdul Rahim who help me with my data collection, to En. Mohammad Yusoph Yaakob for his invaluable help as research assistant during my data collection, to Zamri Rosli for helping me preparing the draft of thesis, to Norfarikah Haneda for invaluable help in doing my analysis, to Evelyn Bigcas for editing my thesis and to all wilder members during our happy moments in the jungle, friends and kababayans whom in their own special ways had greatly helped me finished this research study.

Finally, to my loving wife Eva and my two daughters En-En and Barbie for the love and support they unselfishly give to me. And above all, praise and glory to the heavenly father “Jehovah”.



I certify that an Examination Committee met on October 28, 2003 to conduct the final examination of Romeo M. Lomoljo on his Master of Science thesis entitled "Phenology and Availability of Fruit Trees and Their Influence on the Abundance of Selected Animals in Logged and Primary Forest of Sungai Lalang Forest Reserve, Selangor" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

AHMAD SAID SAJAP, Ph.D.

Professor
Faculty of Forestry
Universiti Putra Malaysia
(Chairman)

MOHAMED ZAKARIA HUSSIN, Ph.D.

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Member)

FARIDAH HANUM BT. IBRAHIM, Ph.D.

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Member)

ABDULLAH MOHD, Ph.D.

Faculty of Forestry
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 05 MAR 2004

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

MOHAMED ZAKARIA HUSSIN, Ph.D.

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Chairman)

FARIDAH HANUM BT. IBRAHIM, Ph. D.

Associate Professor
Faculty of Forestry
Universiti Putra Malaysia
(Members)

ABDULLAH MOHD, Ph. D.

Faculty of Forestry
Universiti Putra Malaysia
(Members)




AINI IDERIS, Ph.D.

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 MAR 2004

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledge. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or other institution.



ROMEO M. LOMOLJO

Date: 03 MAR 2024

TABLE OF CONTENTS

		Page
DEDICATION		ii
ABSTRACT		iii
ABSTRAK		v
ACKNOWLEDGEMENTS		vii
APPROVAL SHEETS		viii
LIST OF TABLES		xv
LIST OF FIGURES		xvi
LIST OF PLATES		xvii
LIST OF ABBREVIATION		xviii
CHAPTER		
1	INTRODUCTION	
	1.1 Problem Statement	2
	1.2 Justification	4
	1.3 General Objective	5
	1.4 Specific Objective	
2	LITERATURE REVIEW	
	2.1 Tropical Rainforest	6
	2.2 Characteristics of the Tropical Rainforest	7
	2.3 Habitat	7
	2.4 Phenology Patterns of Plant in Tropical forest	8
	2.5 Flowering and Fruiting	9
	2.6 Plant as Food	12
	2.7 Dispersal, Fruit Utilization and Seed Predation	13
	2.8 Effects of Logging on Plants	15
	2.9 Effects of Logging on Animals	17
	2.10 Effects of Logging on Food Resources	19
	2.11 Effects Of Logging on Microclimate	20
3	MATERIALS AND METHODS	
	3.1 Site Description	22
	3.2 Sampling Procedure	25
	3.3 Phenological Study	25
	3.3.1 Data Collection	26
	3.4 Data Analysis	27
	3.5 Selectively Logged Forest	28
	3.5.1 Compartment 18 (5-year-old logged forest)	28
	3.5.2 Compartment 33 (10-year-old logged forest)	31
	3.6 Primary Forest	33
	3.7 Climate	34
	3.8 Topography	36

4	RESULTS	
4.1	Plant Composition in the Three Compartment	37
4.1.1	Comparison of Plant and Stand Density Among Three Compartments	38
4.2	Fruit Tree Composition in the Three Compartments	39
4.2.1	Compartment 24 (VJR)	39
4.2.2	Compartment 18 (5-year-old logged forest)	40
4.2.3	Compartment 33 (10-year-old logged forest)	42
4.3	Phenological Results in the Three Compartments	43
4.3.1	Leafing Pattern	43
4.3.2	Flowering Pattern	47
4.3.3	Fruiting Pattern	50
4.4	Species Distribution of Fruit Trees in the Three Compartments	53
4.4.1	Compartment 24 (VJR-ridgetop)	56
4.4.2	Compartment 24 (VJR-midslope)	54
4.4.3	Compartment 24 (VJR-valley bottom)	55
4.4.4	Compartment 33 (10-year-old logged forest -ridgetop)	62
4.4.5	Compartment 33 (10-year-old logged forest – midslope)	64
4.4.6	Compartment 33 (10-year-old logged forest – valley bottom)	66
4.4.7	Compartment 18 (5-year-old logged forest – ridgetop)	68
4.4.8	Compartment 18 (5-year-old logged forest – midslope)	70
4.4.9	Compartment 18 (5-year-old logged forest – valley bottom)	72
4.5	Diversity of Fruit Trees in the Three Different Study Sites	74
4.5.1	Species Diversity	74
4.5.2	Species Richness	74
4.5.3	Species Evenness	75
4.6	Frequency of Flushing	77
4.7	Types of Fruits Available to birds, Primates and Small Mammals Among the Three Compartments	87
4.8	Correlation Between Phenological Activities, Frugivorous and Insectivorous Birds and the Pheasants	89
4.10	Correlation between Phenological activities, primates and small mammals	92
5	DISCUSSION	
5.1	Plant Composition in the Three Compartment	106
5.1.1	Comparison of Plant and Stand Density Among the Three Compartments	107
5.2	Fruit Tree Composition in the Three Compartments	107
5.3	Phenological Results in the Three Compartments	108
5.3.1	Leafing	108

5.3.2 Flowering	109
5.3.3 Fruiting	110
5.4 Species Distribution of Fruit Trees in the Three Compartments	112
5.5 Diversity of Fruit Trees in the Three Compartments	113
5.5.1 Species Diversity	113
5.5.2 Species Richness	114
5.5.3 Species Evenness	114
5.6 Fruits Available to birds, Primates and Small Mammals in the Three Compartments	112
5.7 Correlation Between Phenological Activities, Frugivorous, Insectivorous Birds and Pheasants	116
5.8 Correlation Between Phenological Activities, Primates and Small Mammals	117
5.12 Distribution of Wildlife Based on the Fruit tree Species	119
6 CONCLUSION AND RECOMMENDATION	120
REFERENCES	123
APPENDICES	132
BIODATA OF THE AUTHOR	154

LIST OF TABLES

Tables	Page
4.1 Total Number of Trees, Families, Genera, and Fruit Trees in Three Study Sites	37
4.2 Forest Composition and Stand Density (tree/ha) Among Plot Elevation Within a Compartment	38
4.3 List of fruit trees and Number of Individual Fruit Trees in C24 (VJR)	40
4.4 List of Fruit Tree Species in C18 (5-year-old logged forest)	41
4.5 List of Fruit Trees and Number of Individual Fruit Trees in C33 (10-year-old logged forest)	43
4.6 Monthly Leafing Pattern of Fruit Trees in the Three Compartments	44
4.7 Monthly Flowering Pattern of Fruit Trees in All Study Sites	47
4.8 Monthly Comparison Fruiting Pattern of Fruit Trees in All Study Sites	50
4.9 Distribution of Species in the Three Plots in Compartment 24 (VJR)	53
4.10 Distribution of Species in the Three Plot in Compartment 33 (10-year-old logged forest)	54
4.11 Comparison of Fruit Tree Diversity in Three Study Sites	76
4.12 Comparison of Shannon's Index Values by Plot in Three Different Study Sites	77
4.13 Ranking of Species Based on the Total Frequency of Leafing in C24 (VJR)	67
4.14 Ranking of Species Based on the Total Frequency of Flowering in C24 (VJR)	68
4.15 Ranking of Species Based on the Total Frequency of Fruiting in C24 (VJR)	80
4.16 Ranking of Species Based on the Total Frequency of Leafing in C33 (10-year-old logged forest)	81
4.17 Ranking of Species Based on the Total Frequency of Flowering in C33 (10-year-old logged forest)	82
4.18 Ranking of Species Based on the Total Frequency of Fruiting in C33 (10-year-old logged forest)	83
4.19 Ranking of Species Based on the Total Frequency of Leafing in C18 (5-year-old logged forest)	84
4.20 Ranking of Species Based on the Total Frequency of Flowering in C18 (5-year-old logged forest)	85
4.21 Ranking of Species Based on the Total of Frequency of Fruiting in C18 (5-year-old logged forest)	86
4.22 Fruit Availability in C24 (VJR) in Sg. Lalang According to Fruit Type	87

4.23 Fruit Availability in C33 (10-year-old logged forest) in Sg. Lalang According to Fruit Type	87
4.28 Fruit Availability in C18 (5-year-old logged forest) in Sg. Lalang According to Fruit Type	88
4.29 Simple Correlation Between Phenological Activities, Frugivorous and Insectivorous Birds and Pheasants	90
4.30 Simple Correlation Between Phenological Activities, Primates and Small Mammals	91

LIST OF FIGURES

Figures	Page
3.1 Location of Study Sites (C18), (C33) and (C24) of Sungai Lalang forest Reserve Semenyih, Selangor, Malaysia.	23
3.2 Map Showing in Full Scale the Study Plot in Compartment 18 of Sungai Lalang Forest Reserve.	29
3.3 Map Showing in Full Scale the study Plot in Compartment 33 of Sungai Lalang Forest Reserve	32
3.4 Map Showing in Full Scale the Study Plot in Compartment 24 in Compartment 24 (VJR) of Sungai Lalang Forest Reserve.	35
3.5 Monthly Rainfall of Sungai Lalang Forest Reserve Semenyih, Selangor.	36
4.1 Number of Fruit Trees Leafing in the Three Different Study Sites of Sungai Lalang Forest Reserve.	46
4.2 Percentage of Fruit Trees Leafing in Three Different Study Sites of Sungai Lalang Forest Reserve.	46
4.3 Number of Fruit Trees Flowering in Three Different Study Sites of Sungai Lalang Forest Reserve.	49
4.4 Percentage of Fruit Trees Flowering in Three Different Study Sites of Sungai Lalang Forest Reserve.	49
4.5 Number of Fruit Trees Fruiting in Three Different Study Sites of Sungai Lalang Forest Reserve.	52
4.6 Percentage of Fruit Trees Fruiting in Three Different Study Sites of Sungai Lalang Forest Reserve	52
4.7 Species Distribution of Fruit Trees in C24 (VJR) Ridgetop of Sungai Lalang Forest Reserve.	57
4.8 Species Distribution of Fruit Trees in C24 (VJR) Midslope of Sungai Lalang Forest Reserve.	59
4.9 Species distribution of Fruit Trees in C24 (VJR) Valley Bottom of Sungai Lalang Forest Reserve.	61
4.10 Species Distribution of Fruit Trees in C33 (10-year-old logged forest) Ridgetop of Sungai Lalang Forest Reserve.	63
4.11 Species Distribution Fruit Trees in C33 (10-year-old logged forest) Midslope of Sungai Lalang Forest Reserve	65
4.12 Species Distribution of Fruit Trees in C33 (10-year-old logged forest) Valley Bottom of Sungai Lalang Forest Reserve.	67
4.13 Species Distribution of Fruit Trees in C18 (5-year-old logged forest) Ridge Top of Sungai Lalang Forest Reserve.	69
4.14 Species Distribution of Fruit Trees in C18 (5-year-old logged forest) Midslope of Sungai Lalang Forest Reserve.	71
4.15 Species Distribution of Fruit Trees in C18 (5-year-old	

logged forest) Valley Bottom of Sungai Lalang Forest Reserve.	73
4.16 Leafing Pattern in the Study Sites	93
4.17 Flowering Activities in the study Sites	93
4.18 Flowering Activities in the Study Sites	94
4.19 <i>Durio</i> sp. Fruiting in the Study Sites	94
4.20 <i>Artocarpus</i> sp. Fruiting in the Study Sites	95
4.21 <i>Artocarpus</i> sp. Fruiting the Study Sites	95
4.22 <i>Artocarpus</i> sp. Fruiting in the Study Site	96
4.23 <i>Macaranga</i> sp. Fruiting in the study Sites	96
4.24 <i>Aporosa</i> sp. Fruiting in the Study Sites	97
4.25 <i>Streblus elongates</i> Fruits Eaten by Squirrels in Study Sites	97
4.26 <i>Nephelium intermedium</i> Fruits Eaten by Small Mammals and Primates	98
4.27 <i>Nephelium intermedium</i> Fruits Eaten by Small Mammals and Primates	98
4.28 <i>Parkia speciosa</i> fruits eaten by Primates and Squirrels in the Study Site	99
4.29 <i>Calamus</i> sp. Eaten by Small Mammals in the Study Sites	99
4.30 <i>Adenanthera bicolor</i> Fruiting in the Study Sites	100
4.31 <i>Baccaurea minor</i> Eaten by Primates and Small Mammals in the study sites	100
4.32 <i>Thaipingensis</i> sp. Fruits Eaten by Obscura Monkey in the Study Sites	101
4.33 <i>Artocarpus</i> sp. Fruits Eaten by Primates and Squirrels in the Study Site	101
4.34 <i>Baccaurea</i> sp. Fruits in the study sites	102
4.35 <i>Baccaurea</i> sp. Fruits Eaten by Primates in the Study Sites	102
4.36 <i>Parkia speciosa</i> Fruiting in the Study Site	103
4.37 <i>Canarium</i> sp. Fruiting in the Study Site	103
4.38 <i>Baccaurea reticulata</i> Fruits Eaten by Small Mammals and Primates in the Study Sites	104
4.39 <i>Bouea oppositifolia</i> Fruits in the Study Sites	104
4.30 <i>Bouea oppositifolia</i> Fruits Scattering in the Forest Floor	105

LIST OF ABBREVIATIONS

Archbu	<i>Archidendron bubalinum</i>
Artoin	<i>Artocarpus integer silvestris</i>
Artolan	<i>Artocarpus lanceifolius</i>
Artonit	<i>Artocarpus nitidus graffethii</i>
Bacmin	<i>Baccaurea minor</i>
Bacret	<i>Baccaurea reticulata</i>
Bouop	<i>Bouea oppositifolia</i>
Canlit	<i>Canarium litorale purpurescens</i>
Canruf	<i>Canarium litorale rufom</i>
Canpil	<i>Canarium pilosum</i>
Caschef	<i>Castanopsis schefferiana</i>
Calatro	<i>Callerya atropurpurea</i>
Diamain	<i>Dialium maingayi</i>
Diaplat	<i>Dailium platysepalum</i>
Elatap	<i>Elateriospermum tapus</i>
Garmal	<i>Garcinia mallaccensis</i>
Garscor	<i>Garcinia scortechii</i>
Garpy	<i>Garcinia pyrifera</i>
Manggra	<i>Mangifera gracilipes</i>
Mangqua	<i>Mangifera quadrifida</i>
Nepcus	<i>Nephelium cuspidatum</i>
Nepin	<i>Nephelium intermedium</i>
Nepmain	<i>Nephelium maingayi</i>
Parspec	<i>Parkia speciosa</i>
Sapbac	<i>Sapium bacatum</i>
Sanlae	<i>Santeria laevigata</i>
Dacros	<i>Dacryodes rostrata</i>
Xeroin	<i>Xerospermum intermedium</i>
Xeronor	<i>Xerospermum noronhianum</i>
C24	Compartment 24
C33	Compartment 33
C18	Compartment 18
VJR	Virgin Jungle Reserve

CHAPTER ONE

INTRODUCTION

The tropical rainforest is one of the most complex ecosystems in the world. The forest plays an important role not only in the production of timber but most importantly in providing environmental services, which include the maintenance of biological diversity, soil and watershed protection, regulation of climate and nutrients cycle. However, for development purposes many forested areas in the tropical region are cleared and used for agricultural production, industrial estate and human settlements.

Many tropical countries especially in Southeast Asia are concerned with the effects of selective logging on fauna and flora. The issue is important because there is a decreased in forested areas and there is a need for greater public awareness. There are a few major problems facing wildlife conservation at a global level today, and among them is the increasing rate of habitat loss due to human pressure on the environment especially the core wildlife habitat such as tropical forest. The tropical hardwood for example, is an extremely valuable economic resource for the Southeast Asian countries. However, the increasing rate of forest exploitation will cause extensive tropical forest of the region to become secondary forest by the beginning of the twenty-first century (Myers 1980).

In Malaysia alone, it is extremely rich in both plants and animal species (Tho 1993). Its high diversity offers a wide variety of natural habitats for flora and fauna. It is estimated that Malaysia has ca. 8000 flowering plant of which ca. 2,650 are tree species. There are also over 800 species of non – flowering plants that have been recorded. The animal is equally diverse with ca. 1000 species of vertebrates and an estimated 20-80 thousand invertebrate's species. Many of these species are indigenous and can be found only in peninsular Malaysia. The destruction of wildlife habitats especially forest areas has reduced the number of wildlife in the forest (John 1986). Many birds and other animals left the logging area as soon as the logging activities begin. Logging and other human activities have destroyed most of their feeding and breeding habits considering that their existence in the forest is highly governed by their food supply. Thus, any form of destruction to their habitat has decreased the food supply and eventually affected their existence in the forest.

Problem Statement

It is unlikely that any single factor limits the density of an animal population (Leighton 1983). Although food resources availability may be ultimately limiting. Distribution and abundance of food frequently correlated with the behaviour pattern of animal species (Dawson 1979). Changes in distribution of food types occurring as a result of selective logging will cause the changes

of feeding on a certain food. Animals feed more on leaf materials in logged and primary forest like primate species, which may be correlated with a reduced availability of flowers and fruits.

This study focused on the phenology and availability of fruit trees and their influence on the abundance of selected animals in logged and primary forest in three different compartments such as Virgin Jungle Reserve (VJR), ten-year-old logged forest (C33) and five-year old logged forest (C18). The reason of choosing this three study sites:

The Compartment 24 (VJR) and the two logged forest study were selected for the following reasons: I) The compartment 24 (VJR) forest areas have been established and is presently used by some researchers working in botany. References to work in the area are readily available and the present work will form part of an integrated research program. Ii) The study area are adjacent to each other, increasing the likelihood that pre-logging conditions (diversity and composition of flora and fauna) in the compartment 33 (10 year old logged forest) and in the compartment 18 which is (5 year old logged forest) were similar to those in the unlogged site. This similarity is crucial if comparisons of phenology activities in relation to different animals of the areas are to be made. Iii) Both areas were far from any human settlements and hunting was almost unknown, increasing the possibility that changes to the plant and animal populations were due only to logging activities.

Justification

Most of the forest areas in Peninsular Malaysia especially the lowland forest had been logged and become degraded and formed into secondary forest. This effect is very much dependent on the intensity of logging destruction and volume of timber extracted and this might be severed to certain species of animals or birds (especially understorey birds) but not the other animals. The indirect effects of logging towards animal's population are the reduction of food and shelter (Johns 1986). For the primate's species, the logging activities seemed to cause less impact on their populations (Sundai 2000). This is probably because, the removal of higher number of dipterocarp trees during logging had cause less impact to their population because dipterocarp are less preferred as food resources by the primates (Chivers 1977a, 1977b; Johns 1986). In addition, the removal of emergent trees and the opening of the canopy cover would increase the amount of sunlight reaching the forest floor. This will encourage the growth of smaller trees especially for non-dipterocarp species, which could provide a potential food sources (e.g. leaves flower and fruits) for the primates. Thus, there is an urgent need to determine the extent of this disturbance towards the fruit tree species that serves as food for animals in the forest. It is important to note that the information gained will become the basis or guide line in formulating the effective management plan on forest utilization and forest operation to ensure the survival and to see these animals in the future.

Objectives of the Study

General Objective

The main objectives of the study were to identify the fruit tree species and determine the food resources for selected animals in logged and primary forest.

Specific Objectives

The specific objectives of this study were

1. To study the phenology of fruit trees, which serve as food resources for selected animals in 10-year old logged forest, 5-year old logged forest and VJR (primary forest).
2. To compare the food resources for selected animals in 10-year old logged forest, 5-year old logged forest and VJR (primary forest).
3. To compare the fruit trees distribution in three compartments within plots.
4. To correlate the abundance of selected animals in relation to fruits available in three different study sites namely; 10-year-old logged forest, 5-year-old logged forest and VJR (primary forest).