



UNIVERSITI PUTRA MALAYSIA

**GERMPLASM PRESERVATION:
STORAGE OF SEED, POLLEN AND TISSUE OF
SELECTED MALAYSIAN ORCHID SPECIES**


SALEH B. KADZIMIN

FP 1987 1

It is hereby certified that we have read this thesis entitled 'Germplasm Preservation: Storage of Seed, Pollen and Tissue of Selected Malaysian Orchids' by Saleh bin Kadzimin, and in our opinion it is satisfactory in terms of scope, quality, and presentation as partial fulfilment of the requirements for the degree of Doctor of Philosophy


.....
ALANG P. ZAINUDDIN, Ph.D.

Assoc. Professor/Dean of Graduate Studies
Universiti Pertanian Malaysia
(Chairman Board of Examiners)


.....
YONEO SAGAWA, Ph.D.

Professor/Director
Lyon Arboretum
University of Hawaii at Manoa
Honolulu, Hawaii
(External Examiner)


.....
ISMAIL HAMZAH, Ph.D.

Assistant Director
Harrisons Malaysian Plantation Bhd
(Research and Advisory Scheme)
(External Examiner)


.....
ZALIHA C. ALANG, Ph.D.

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Pertanian Malaysia
(Internal Examiner)

Chin Hoong Fong

.....
CHIN HOONG FONG, Ph.D.

Professor

Faculty of Agriculture

Universiti Pertanian Malaysia

(Internal Examiner/Supervisor)

This thesis was submitted to the Senate of Universiti
Pertanian Malaysia and was accepted as partial fulfilment of
the requirements for the degree of Doctor of Philosophy.



ALANG P. ZAINUDDIN, Ph. D.
Associate Professor/Dean of Graduate Studies

Date: 8th October 1987

GERMPLASM PRESERVATION:
STORAGE OF SEED, POLLEN AND TISSUE OF
SELECTED MALAYSIAN ORCHID SPECIES

by

Saleh B. Kadzimin

A thesis submitted in partial fulfilment of the
requirements for the degree of Doctor of Philosophy
in the Faculty of Agriculture
Universiti Pertanian Malaysia

April 1987



Lovingly dedicated to

S H A Z R I N

S H A Z W A N

K A M A R I A H



ACKNOWLEDGEMENTS

I wish to express my sincere appreciation and thanks to my supervisor, Professor Chin Hoong Fong, of the Department of Agronomy and Horticulture, Universiti Pertanian Malaysia, for his invaluable advice, guidance and encouragement throughout the study.

I am grateful to the staff of the Electron Microscope Unit of the University, Puan Aminah Jusoh and Encik Ho Oi Kuan for their assistance in electron microscopy work.

Special thanks are extended to Encik Mohd. Said Saad, Lecturer, Department of Agronomy and Horticulture, Universiti Pertanian Malaysia, for his kind assistance in the statistical analyses of the study. I also wish to extend many thanks to Puan Rahimah Abdullah of the same Department for her patience in typing the manuscript.

Lastly, but not the least, I wish to thank my wife, Kamariah, for her help and encouragement, understanding and tolerance throughout the duration of the study.



TABLE OF CONTENTS

| | PAGE |
|--|-------|
| ACKNOWLEDGEMENTS | iii |
| TABLE OF CONTENTS | iv |
| LIST OF TABLES | viii |
| LIST OF FIGURES | ix |
| LIST OF PLATES | xi |
| LIST OF APPENDICES | xiv |
| LIST OF ABBREVIATIONS | xvii |
| ABSTRACT | xviii |
| | |
| CHAPTER 1 - INTRODUCTION | 1 |
| | |
| CHAPTER 2 - LITERATURE REVIEW | 7 |
| STORAGE OF SEED | 7 |
| Introduction | 7 |
| The Orchid Seed | 8 |
| Orchid Seed Storage | 9 |
| STORAGE OF POLLEN | 14 |
| Introduction | 14 |
| The Orchid Pollen | 16 |
| Orchid Pollen Storage | 16 |
| STORAGE OF TISSUE | 19 |
| Introduction | 19 |
| General Features of Cryopreservation | 20 |



| | PAGE |
|--|-----------|
| Preconditioning of Cultured Tissue | 23 |
| Cooling Rate, Thawing and Post-thaw Recovery .. | 30 |
| CHAPTER 3 - MATERIALS AND METHODS | 37 |
| THE SPECIES UNDER STUDY | 37 |
| SEED | 38 |
| Seed Processing | 38 |
| Determination of Seed Moisture Content | 39 |
| Establishment of Seed Moisture Content | 40 |
| Determination of Seed Viability | 40 |
| Electron Microscopic Observation of Seed | 43 |
| Experiment I. Effect of Relative Humidity on Seed Moisture Content | 43 |
| Experiment II. Effect of Initial Seed Moisture Content on Germination | 44 |
| Experiment III. Effect of Seed Moisture Content on Germination Rate | 44 |
| Experiment IV. Effect of Seed Moisture Content on Viability in Conventional Storage | 45 |
| Experiment V. Effect of Seed Moisture Content, Storage Duration, and Thawing Rates on Viability in Cryogenic Storage | 45 |
| Experiment VI. Effect of Storage Temperature on Seed Viability | 46 |
| Experiment VII. Effect of Cryogenic Temperature on Rate of Germination | 49 |
| Experiment VIII. Effect of Cryogenic Storage on Seed Moisture Content | 47 |



| | PAGE |
|---|------|
| POLLEN | 49 |
| Pollen Processing | 49 |
| Determination of Pollen Viability | 49 |
| Experiment on the Effect of Storage Temperature on Pollen Viability | 50 |
| Electron Microscopic Observation of Pollen | 51 |
| TISSUE | 51 |
| General Procedure in the Cryopreservation of Orchid Protocorm | 51 |
| Experiment I. Effect of Incubation Time and Concentration of Cryoprotectant on Viability of Protocorm Section | 56 |
| Experiment II. Effect of Prefreezing Temperature on Viability of Protocorm Section | 56 |
| Experiment III. Effect of Storage in Liquid Nitrogen on Viability of Protocorm Section | 57 |
| Electron Microscopic Observations of Protocorm Tissue | 58 |
| CHAPTER 4 - RESULTS AND DISCUSSION | 60 |
| THE SPECIES UNDER STUDY | 60 |
| STORAGE OF SEED | 62 |
| Seed Morphology | 62 |
| Comparison on Seed Viability Tests | 65 |
| Experiment I. Effect of Relative Humidity on Seed Moisture Content | 72 |
| Experiment II. Effect of Initial Seed Moisture Content on Germination | 75 |



| | PAGE |
|--|------|
| Experiment III. Effect of Seed Moisture Content on Germination Rate | 79 |
| Experiment IV. Effect of Seed Moisture Content on Viability in Conventional Storage | 82 |
| Experiment V. Effect of Seed Moisture Content, Storage Duration, and Thawing Rates on Viability in Cryogenic Storage | 87 |
| Experiment VI. Effect of Storage Temperature on Seed Viability | 92 |
| Experiment VII. Effect of Cryogenic Temperature on Rate of Germination | 98 |
| Experiment VIII. Effect of Cryogenic Storage on Seed Moisture Content | 101 |
| STORAGE OF POLLEN | 107 |
| Pollen Structure | 107 |
| Experiment on the Effect of Storage Temperature on Pollen Viability | 113 |
| STORAGE OF TISSUE | 121 |
| Protocorm Development | 121 |
| Experiment I. Effect of Incubation Time and Concentration of Cryoprotectant on Viability of Protocorm Section | 125 |
| Experiments II and III. Effect of Prefreezing Temperature on Survival of Protocorm Section | 130 |
| Electron Microscopic Observations of Protocorm Tissue | 143 |
| CHAPTER 5 - SUMMARY AND CONCLUSION | 153 |
| BIBLIOGRAPHY | 166 |
| APPENDICES | 176 |



LIST OF TABLES

| TABLE | | PAGE |
|-------|---|------|
| 1 | Comparison of Germination Percentages as Determined by Germination and Differential Staining Techniques Using Fresh Seeds | 71 |
| 2 | Variations in Initial Seed Moisture Content and Corresponding Germination of Three Orchid Species | 76 |
| 3 | Summary of <u>V. hookeriana</u> Protocorm Development Following Reculturing in Cryoprotectant-free Medium | 138 |



LIST OF FIGURES

| FIGURE | | PAGE |
|--------|---|------|
| 1 | The Relative Humidity and Moisture Content Equilibrium Relationship of Seeds of <u>A. graminifolia</u> , <u>S. plicata</u> and <u>C. finlaysonianum</u> | 73 |
| 2 | Effect of Seed Moisture Content on Rate of Germination of <u>C. finlaysonianum</u> Seed Following Storage Under Laboratory Conditions .. | 80 |
| 3 | Effect of Storage Under Laboratory Conditions on Percentage Germination of <u>A. graminifolia</u> Seed of Various Levels of Seed Moisture Content | 83 |
| 4 | Effect of Storage Under Laboratory Conditions on Percentage Germination of <u>C. finlaysonianum</u> Seed of Various Levels of Seed Moisture Content | 84 |
| 5 | Effect of Storage Under Laboratory Conditions on Percentage Germination of <u>S. plicata</u> Seed of Various Levels of Seed Moisture Content | 85 |
| 6 | Effect of Storage Duration on Percentage Germination of <u>C. finlaysonianum</u> Seed of Various Levels of Moisture Content (Rapid Thaw) | 89 |
| 7 | Effect of Storage Duration on Percentage Germination of <u>C. finlaysonianum</u> Seed of Various Levels of Moisture Content (Slow Thaw) | 90 |
| 8 | Effect of Seed Moisture Content and Thawing Rate on Viability of <u>C. finlaysonianum</u> Seeds Stored in Liquid Nitrogen | 91 |
| 9 | Effect of Storage under Different Temperature Regimes on Percentage Germination of <u>A. graminifolia</u> Seed | 94 |
| 10 | Effect of Storage under Different Temperature Regimes on Percentage Germination of <u>C. finlaysonianum</u> Seed | 95 |



| FIGURE | | PAGE |
|--------|---|------|
| 11 | Effect of Storage under Different Temperature Regimes on Percentage Germination of <u>S. plicata</u> Seed | 97 |
| 12 | Effect of Storage Duration on Rate of Germination of <u>S. plicata</u> Seed | 99 |
| 13 | Effect of Storage Duration on Seed Moisture Content of <u>S. plicata</u> | 102 |
| 14 | Effect of Storage Duration on Viability of <u>A. graminifolia</u> Pollen | 114 |
| 15 | Effect of Storage Duration on Viability of <u>S. plicata</u> Pollen | 116 |
| 16 | Effect of Storage Duration on Viability of <u>V. hookeriana</u> Pollen | 117 |
| 17 | Effect of Incubation Time on Percentage Viability of <u>V. hookeriana</u> Protocorm | 126 |
| 18 | Effect of Sub-zero Temperature on Percentage Survival of <u>V. hookeriana</u> Protocorm | 132 |



LIST OF PLATES

| PLATE | | PAGE |
|-------|--|------|
| I | The species under study. (A) <u>A. graminifolia</u> (B) <u>C. finlaysonianum</u> , (C) <u>S. plicata</u> , and (D) <u>V. hookeriana</u> | 61 |
| II | Scanning Electron, Micrographs of Whole Seed (A) <u>A. graminifolia</u> x 130, (B) <u>C.</u> <u>finlaysonianum</u> x 100, (C) <u>S. plicata</u> x 130, (D) <u>V. hookeriana</u> x 200 | 64 |
| III | Scanning Electron Micrographs of <u>A. graminifolia</u> Seed after 20 Weeks in Storage. (A) Whole Seed x 130, (B) 5°C x 1000, (C) -15°C x 860, (D) -196°C x 720 | 66 |
| IV | Scanning Electron Micrographs of <u>C.</u> <u>finlaysonianum</u> Seed After 20 Weeks in Storage (A) Whole Seed x 100, (B) 5°C x 600, (C) -15°C x 660, (D) -196°C x 440 | 67 |
| V | Scanning Electron Micrographs of <u>S. plicata</u> Seed After 20 Weeks in Storage (A) Whole Seed x 130, (B) 25°C x 720, (C) -15°C x 660, (D) -196°C x 540 | 68 |
| VI | Scanning Electron Micrographs of <u>V.</u> <u>hookeriana</u> Seed After 20 Weeks in Storage (A) Whole Seed x 200, (B) 5°C x 1600, (C) -15°C x 132 (D) -196°C x 1000 | 69 |
| VII | Scanning Electron Micrographs of Fresh Pollinia (A) <u>A. graminifolia</u> x 72, (B) <u>C.</u> <u>finlaysonianum</u> x 32, (C) <u>S. plicata</u> x 54, (D) <u>V. hookeriana</u> x 44 | 108 |
| VIII | Scanning Electron Micrographs of Fresh Pollen Grains. (A) <u>A. graminifolia</u> x 1000, (B) <u>C. finlaysonianum</u> x 1000, (C) <u>S. plicata</u> x 1000, (D) <u>V. hookeriana</u> x 2000 | 109 |
| IX | Scanning Electron Micrographs of Pollen Grains of <u>A. graminifolia</u> After 8 weeks in Storage. (A) -15°C x 1000, (B) -15°C x 2000, (C) -196°C x 2000 | 110 |



| PLATE | | PAGE |
|-------|---|------|
| X | Scanning Electron Micrographs of Pollen Grains of <u>S. plicata</u> After 8 Weeks in Storage (A) 25°C x 2000, (B) -15°C x 1000, (C) -196°C x 2000 | 111 |
| XI | Scanning Electron Micrographs of Pollen Grains of <u>V. hookeriana</u> After 8 weeks in Storage (A) 5°C x 2000, (B) -15°C x 1000, (C) -196°C x 1000 (D) -196°C x 1500 | 112 |
| XII | <u>Vanda hookeriana</u> Protocorm (A) Incubation with Cryoprotectant, (B) Aggregates of Protocorm, (C) Stages of Differentiation | 122 |
| XIII | Protocorm Showing Regrowth Following Storage in Liquid Nitrogen (A) 12 Weeks, (B) 18 Weeks, After Reculture | 124 |
| XIV | Scanning Electron Micrographs of Fresh Protocorm Section (A) Side View x 40, (B) Top View x 54, (C) Wall Structure x 480 ... | 145 |
| XV | Scanning Electron Micrographs Showing Undamaged Cells Following Storage in Liquid Nitrogen with Cryoprotection from (A) 5% Glucose x 480, (B) 10% Glucose x 1000, (C) 5% DMSO + 5% Sucrose x 480, (D) 10% DMSO + 10% Sucrose | 146 |
| XVI | Scanning Electron Micrographs Showing Cell Collapse Following Storage in Liquid Nitrogen. Specimen Treated With: (A) 5% DMSO x 480, (B) 5% Glycerol x 480, (C) 10% DMSO + 10% Ethylene Glycol x 1800, (D) 10% DMSO + 5% Glycerol x 1000 | 147 |
| XVII | Transmission Electron Micrographs of Protocorm (A) Control x 7700, (B) Treated With VW Medium x 2150 | 149 |
| XVIII | Transmission Electron Micrographs of Protocorm After Storage in Liquid Nitrogen, Cryoprotected by (A) 5% DMSO x 3550, (B) 10% DMSO x 2750 | 150 |



| PLATE | | PAGE |
|-------|---|------|
| XIX | Transmission Electron Micrographs of Protocorm After Storage in Liquid Nitrogen, Cryoprotected by 10% DMSO + 5% Glucose (A) x 1650, (B) x 4600 | 151 |
| XX | Transmission Electron Micrographs of Protocorm After Storage in Liquid Nitrogen, Cryoprotected by (A) 5% DMSO + 5% Sucrose x 3550, (B) 5% DMSO + 10% Glycerol x 2750 | 152 |



LIST OF APPENDICES

| APPENDIX | PAGE |
|----------|---|
| 1 | Relationship Between Concentration and Vapor Pressure of Potassium hydroxide (KOH) and Equilibrium Relative Humidities at 20°C 176 |
| 2 | Medium as Used for the Germination of <u>A. graminifolia</u> Seed 177 |
| 3 | Medium As Used for the Germination of <u>C. finlaysonianum</u> and <u>V. hookeriana</u> Seeds 178 |
| 4 | Medium As Used for the Germination of <u>S. plicata</u> seeds 179 |
| 5 | Medium As Used for the Culture of Explant from Shoot Tip and Protocorm of <u>V. hookeriana</u> 181 |
| 6 | Preparation of Orchid Pollen and Protocorm Tissue for Scanning Electron Microscopy 182 |
| 7 | Preparation of Orchid Protocorm Tissue for Transmission Electron Microscopy 183 |
| 8 | Cryoprotectants Used in the Cryopreservation of Orchid Protocorm 184 |
| 9 | Analysis for the Comparison of Germination Percentage as Determined by Germination and Differential Staining Techniques Using Fresh Seeds 185 |
| 10 | Equilibrium Seed Moisture Content Following Exposure to Various Levels of Relative Humidity 186 |
| 11 | Analysis for Variations in Initial Seed Moisture Content and Corresponding Germination of Three Orchid Species 188 |
| 12 | Effect of Seed Moisture Content on Germination Rate of <u>C. finlaysonianum</u> under Laboratory (25°C) Conditions 191 |



APPENDIX

PAGE

| | | |
|----|---|-----|
| 13 | Effect of Seed Moisture Content on Germination of <u>A. graminifolia</u> Seed Stored under Laboratory (25°C) Conditions | 194 |
| 14 | Effect of Seed Moisture Content on Germination of <u>C. finlaysonianum</u> Seed Stored under Laboratory (25°C) Conditions | 196 |
| 15 | Effect of Seed Moisture Content on Germination of <u>S. plicata</u> Seed Stored under Laboratory (25°C) Conditions | 198 |
| 16 | Effect of Seed Moisture Content on Germination of <u>C. finlaysonianum</u> Seeds Stored in Liquid Nitrogen | 200 |
| 17 | Effect of Storage Temperature on Germination of <u>A. graminifolia</u> Seed | 203 |
| 18 | Effect of Storage Temperature on Germination of <u>C. finlaysonianum</u> Seed | 205 |
| 19 | Effect of Storage Temperature on Germination of <u>S. plicata</u> Seed | 207 |
| 20 | Analysis for the Effect of Storage Temperature and Duration on Germination of Three Orchid Species | 209 |
| 21 | Effect of Cryogenic Storage on Rate of Germination of <u>S. plicata</u> Seed | 211 |
| 22 | Effect of Cryogenic Storage on Seed Moisture Content of <u>S. plicata</u> | 213 |
| 23 | Effect of Storage Temperature on Germination of <u>A. graminifolia</u> Pollen | 215 |
| 24 | Effect of Storage Temperature on Germination of <u>S. plicata</u> Pollen | 217 |
| 25 | Effect of Storage Temperature on Germination of <u>V. hookeriana</u> Pollen | 219 |
| 26 | Survival of <u>V. hookeriana</u> Protocorm Sections Following Preculturing in Various Cryoprotectants | 221 |



APPENDIX

PAGE

- 27 Cryoprotective Effect of DMSO, Ethylene glycol, Glycerol, Glucose, and Sucrose on V. hookeriana Protocorm Sections Cooled to Various Sub-zero Temperatures 224
- 28 Survival of V. hookeriana Protocorm Sections Following Prefreezing to Different Temperatures and Storage in Liquid Nitrogen ... 227



LIST OF ABBREVIATIONS

| | | |
|-----|---|---|
| Å | : | Angstrom |
| °C | : | Degrees Celcius |
| cm | : | Centimeter |
| CV | : | Coefficient of variation |
| DF | : | Degree of freedom |
| F | : | Calculated F-value |
| g | : | Gram |
| Hg | : | Mercury |
| kg | : | Kilogram |
| kV | : | Kilo Volt |
| kPa | : | Kilo Pascal |
| m | : | Meter |
| mg | : | Milligram |
| min | : | Minute |
| ml | : | Milliliter |
| mm | : | Millimeter |
| M | : | Molar |
| MS | : | Mean square |
| psi | : | Pound per square inch |
| REP | : | Replication |
| SE | : | Standard error |
| SED | : | Standard error of difference between means |
| ug | : | Microgram |
| v/v | : | Volume-to-volume |
| w/v | : | Weight-to-volume |



An abstract of the thesis presented to the Senate of Universiti
Pertanian Malaysia in partial fulfilment of the requirements
for the degree of Doctor of Philosophy

GERMPLASM PRESERVATION: STORAGE OF SEED, POLLEN AND
TISSUE OF SELECTED MALAYSIAN ORCHID SPECIES

by

Saleh B. Kadzimin

April, 1987

Supervisor : Professor Chin Hoong Fong

Faculty : Agriculture

In an attempt to develop methods for cheap, long-term preservation of orchid germplasm, seed, pollen and tissue of four local species were subjected to various storage conditions and subsequently determined of their effects on viability. The resultant quality of stored materials varied over the range of conditions and species investigated.

Storage of seeds of Arundina graminifolia, Cymbidium finlaysonianum and Spathoglottis plicata at seed moisture content of less than 5 percent and at storage temperature of less than 5°C gave percentage viability that ranged between 60 and 90 percent for a storage duration of more than twelve weeks. Storage in liquid nitrogen (-196°C) gave the highest



mean among the temperatures tested, confirming the orthodox nature of the seed.

Over the range of storage temperatures tested, storing pollen of A. graminifolia, S. plicata and Vanda hookeriana in liquid nitrogen gave the highest mean of viability (15, 64 and 57 percent respectively) for up to fourteen weeks of storage.

Protocorm tissue of V. hookeriana treated with cryoprotectant and prefrozen to a sub-zero temperature briefly before effecting storage in liquid nitrogen, offers a potential system to long-term preservation of the orchid species. The highest mean of survival (5.6 percent) following liquid nitrogen storage was afforded by cryoprotecting with 5 percent dimethyl sulfoxide in combination with 5 percent sucrose, and subsequent prefreezing to -40°C prior to storage.

Cryopreservation of orchid seed, pollen and tissue presents a number of potential practical advantages. The present study explored this technology as a supplement or extension to existing conventional systems of germplasm preservation.



Abstrak tesis yang dikemukakan kepada Senat Universiti
Pertanian Malaysia sebagai memenuhi sebahagian daripada syarat
untuk ijazah Doktor Falsafah

PENGEKALAN JANAPLASMA: PENYIMPANAN BIJI BENIH, DEBUNGA
DAN TISU SPESIS PILIHAN ORKID MALAYSIA

oleh

Saleh B. Kadzimin

April, 1987

Penyelia : Profesor Chin Hoong Fong

Fakulti : Pertanian

Dalam usaha mencipta teknik untuk pengekalan janaplasma orkid dalam jangkamasa yang panjang, biji benih, debunga dan tisu dari empat spesis tempatan telah dirawat di dalam beberapa keadaan simpanan dan kemudiannya dikaji kesan-kesan rawatan ke atas daya percambahannya. Mutu bahan yang tersimpan berbeza-beza mengikut keadaan simpanan dan spesis yang dikaji.

Penyimpanan biji benih Arundina graminifolia, Cymbidium finlaysonianum dan Spathoglottis plicata pada paras kelembapan kurang dari 5 peratus dan pada suhu kurang dari 5°C memberi daya percambahan biji benih diantara 60 hingga 90 peratus dalam jangkamasa simpanan melebihi dua belas minggu. Penyimpanan bijibenih dalam cecair nitrogen (-196°C) memberi purata



tertinggi di antara suhu-suhu yang diuji, mengesahkan sifat 'orthodox' biji benih tersebut.

Di antara suhu-suhu yang diuji, penyimpanan debunga A. graminifolia, S. plicata dan Vanda hookeriana dalam cecair nitrogen memberi purata daya percambahan yang tertinggi (15, 64 dan 57 peratus tiap-tiap satu) dalam jangkamasa simpanan selama empat belas minggu.

Tisu protokom V. hookeriana yang telah dirawat dengan 'cryoprotectant' dan dibeku di bawah paras 0°C sebelum disimpan di dalam cecair nitrogen, merupakan satu sistem pengekalan jangkapanjang yang berpotensi bagi spesis orkid. Purata kemandirian tertinggi (5.6 peratus) selepas penyimpanan dalam cecair nitrogen telah diperolehi dari rawatan 5 peratus dwimetil sulfoksida bercampur 5 peratus sukros, dan pembekuan awal ke -40°C sebelum disimpan.

Penyimpanan krio biji benih, debunga dan tisu mempunyai beberapa kebaikan. Penyelidikan ini mengkaji teknologi tersebut sebagai satu tambahan kepada sistem-sistem pengekalan janaplasma yang sedia ada.

