

UNIVERSITI PUTRA MALAYSIA

EFFECTS OF GAMMA IRRADIATION ON IN VITRO CULTURES OF SELECTED ORCHID HYBRIDS

ADRAIN LING CHIENG KUANG

FP 1999 15



EFFECTS OF GAMMA IRRADIATION ON IN VITRO CULTURES OF SELECTED ORCHID HYBRIDS

ADRAIN LING CHIENG KUANG

MASTER OF AGRICULTURAL SCIENCE UNIVERSITI PUTRA MALAYSIA

1999



EFFECTS OF GAMMA IRRADIATION ON IN VITRO CULTURES OF SELECTED ORCHID HYBRIDS

By

ADRAIN LING CHIENG KUANG

Thesis Submitted in Fulfilment of the Requirements for the Degree of Master of Agricultural Science in the Faculty of Agriculture,
Universiti Putra Malaysia

February 1999



To Almighty God for His blessings.

To my parents Mary and Joseph Ling, my sister Susana and my brothers Michael and Vincent for their support, sacrifices and love.



ACKNOWLEDGEMENTS

My sincere thanks to the Chairman, Associate Professor Saleh Kadzimin, Ph.D. of Faculty of Agriculture, Universiti Putra Malaysia (UPM) for his supervision and patience.

My deepest gratitude goes to the members of my Supervisory Committee K. Harikrishna, Ph.D. of Faculty of Food Science and Biotechnology, Professor Marziah Mahmood, Ph.D. of Faculty of Science and Environment Studies, UPM and Mohd. Nazir Basiran, Ph.D. of Malaysia Institute for Nuclear Technology Research (MINT) for their suggestions and understanding.

I am grateful to Miss Haryati Jamhari, Mr. Abdul Rahman Sidam, Mdm. Maininah Tahir, staff and colleague of the Tissue Culture Laboratory and Mr. Ong Choon Hoe of the Seed Technology Laboratory, Faculty of Agriculture, UPM for their help during the course of this project.

To the working team of the Genetic Laboratory, Faculty of Food Science and Biotechnology, UPM especially Mdm. Norliza Hj Mat Sharif, Miss Lim Chin Ching, Mdm. Meilina Ong Abdullah (Mei), Mr. Wong Hann Ling, Miss Ooi Siew Eng, Yoon Soo Lee (Yooni), Ph.D., Mr. M. Sugumaran, Miss N.



Parameswari, Miss Meta Sritua Arief, Mr. Choong Chieh Wean, Miss Chan Pick Kuen, Mdm. Norlia and Mdm. Rogayah, thank you for your kindness, patience and help. It has been an honour working with all of you.

Sincere thanks are extended to Mr. Samson Soon, Mr. Yap Lip Vun and Mdm. Janna Ong Abdullah for their help. To all my friends, thank you for being the source of my strength and making this journey much more meaningful.



TABLE OF CONTENTS

	ABLES
	IGURES LATES
_	BBREVIATIONS
	T
JULIKAN	
HAPTE	R
I	INTRODUCTION
II	LITERATURE REVIEW
	Mutation Induction
	Types of Mutations
	Types of Mutagens
	Effects of Ionising Radiation
	Photons Absorption
	Chemical Changes
	Damage to DNA
	Killing Effect
	Repair of DNA
	Irradiation Dose
	Measurement of Irradiation Dose
	Mutation Induction on In Vitro Cultures
	Measure of Effects of Irradiation
	Random Amplified Polymorphic DNA (RAPD)
	Influence of Different Parameters on the RAPD
	DNA Concentration
	Mg ²⁺ Concentration
	Taq DNA Polymerase Concentration
	Annealing and Denaturing Temperature
III	MATERIALS AND METHOD
	Location
	Plant Materials



	Irradiation	3
	Post-irradiation Treatment	3
	Parameters	3
	Survival	3
	Plantlet Regeneration	34
	Fresh Weight	34
	Dry Weight	3
	DNA Analysis	3
	Subculturing	3:
	Planting and Maintenance of Plantlet	3:
	Genomic DNA Isolation	3
	Polymerase Chain Reaction (PCR)	3
	Recovery of DNA from Agarose Gel	3
	Southern Analysis	3
	Southern Blotting	3
	Southern Hybridisation	3
	Preparation of Labelled Probe	4
	Blocking, Antibody Incubation and Washes	4
	Signal Generation and Detection	4
	Reprobing Blot	4
	Transformation	4
	Plasmid Miniprep	4
	Plasmid DNA Purification	4
	Sequencing and Computer Analysis	4
	Southern Blot Analysis of Genomic DNA	4
	Experimental Design and Statistical Analysis	4
ΙV	RESULTS AND DISCUSSION	4
	Effect of γ-Irradiation on the Survival of <i>In Vitro</i> Cultures	
	of Mokara Chark Kuan	4
	Effect of γ-Irradiation on the Survival of <i>In Vitro</i> Cultures	
	of <i>Dendrobium</i> Jacky	4
	Effect of γ-Irradiation on Plantlet Regeneration of In Vitro	
	Cultures of Mokara Chark Kuan	5
	Effect of γ-Irradiation on Plantlet Regeneration of In Vitro	
	Cultures of Dendrobium Jacky	5
	Effect of γ-Irradiation on the Fresh Weight, Dry Weight	
	and Dry Weight Gain of In Vitro Cultures of Mokara	
	Chark Kuan	5



	and Dry Weight Gain of In Vitro Cultures of Dendrobium Jacky	58
	Effect of γ-Irradiation on the Genomic DNA of <i>Mokara</i> Chark Kuan	60
	Effect of γ-Irradiation on the Genomic DNA of Dendrobium Jacky	76
V	CONCLUSION	77
BIBLIOGRA	РНҮ	82
APPENDICE	s	92
Appendix A	Additional Tables	93
Appendix B	Figures	95
Appendix C	ANOVA Tables	97
Appendix D	Sequences	96
Appendix E	Alignments	109
Appendix F	Possible binding sites of pAL202 over pAL108 by forming secondary structure (loop)	113
X/IT A		114



LIST OF TABLES

Γable		Page
1	The Absorbed Doses as Measured by the Fricke Dosimeter in Set I	32
2	The Absorbed Doses as Measured by the Fricke Dosimeter in Set II	32
3	Mean Percentage in Survival of <i>In Vitro</i> Cultures of <i>Mokara</i> Chark Kuan Adjusted to the Control (Non-irradiated) under Different Doses of γ-Irradiation (Set I)	47
4	Mean Percentage in Survival of <i>In Vitro</i> Cultures of <i>Mokara</i> Chark Kuan Adjusted to the Control (Non-irradiated) under Different Doses of γ-Irradiation (Set II)	47
5	Mean Percentage in Survival of <i>In Vitro</i> Cultures of <i>Dendrobium</i> Jacky Adjusted to the Control (Non-irradiated) under Different Doses of γ-Irradiation (Set I)	49
6	Mean Percentage in Survival of <i>In Vitro</i> Cultures of <i>Dendrobium</i> Jacky Adjusted to the Control (Non-irradiated) under Different Doses of γ-Irradiation (Set II)	50
7	Mean Number of Mokara Chark Kuan Plantlets Regenerated 15 Weeks after Various Doses of γ-Irradiation (Set I)	51
8	Mean Number of <i>Mokara</i> Chark Kuan Plantlets Regenerated 15 Weeks after Various Doses of γ-Irradiation (Set II)	52
9	Mean Number of <i>Dendrobium</i> Jacky Plantlets Regenerated 15 Weeks after Various Doses of γ-Irradiation (Set I)	53
10	Mean Number of <i>Dendrobium</i> Jacky Plantlets Regenerated 15 Weeks after Various Doses of γ-Irradiation (Set II)	53



11	Mean Fresh Weight, Dry Weight and Dry Weight Gain (g/sample) of <i>In Vitro</i> Cultures of <i>Mokara</i> Chark Kuan 15 Weeks after Various Doses of γ-Irradiation (Set I)	55
12	Mean Fresh Weight, Dry Weight and Dry Weight Gain (g/sample) of <i>In Vitro</i> Cultures of <i>Mokara</i> Chark Kuan 15 Weeks after Various Doses of γ-Irradiation (Set II)	56
13	Mean Fresh Weight, Dry Weight and Dry Weight Gain (g/sample) of <i>In Vitro</i> Cultures of <i>Dendrobium</i> Jacky 15 Weeks after Various Doses of γ-Irradiation (Set I)	58
14	Mean Fresh Weight, Dry Weight and Dry Weight Gain (g/sample) of <i>In Vitro</i> Cultures of <i>Dendrobium</i> Jacky 15 Weeks after Various Doses of γ-Irradiation (Set II)	59
15	Modified Vacin and Went's Formulation as Used in Media Preparation	93
16	Primers from Operon Kit AE	94
17	ANOVA for Survival Set I (Mokara Chark Kuan)	97
18	ANOVA for Survival Set II (Mokara Chark Kuan)	97
19	ANOVA for Survival Set I (Dendrobium Jacky)	98
20	ANOVA for Survival Set II (Dendrobium Jacky)	98
21	ANOVA for Plantlet Regeneration Set I (Mokara Chark Kuan)	99
22	ANOVA for Plantlet Regeneration Set II (Mokara Chark Kuan)	99
23	ANOVA for Plantlet Regeneration Set I (Dendrobium Jacky)	100
24	ANOVA for Plantlet Regeneration Set II (Dendrobium Jacky)	100
25	ANOVA for Fresh Weight Set I (Mokara Chark Kuan)	101



26	ANOVA for Dry Weight Set I (Mokara Chark Kuan)	101
27	ANOVA for Dry Weight Gain Set I (Mokara Chark Kuan)	102
28	ANOVA for Fresh Weight Set II (Mokara Chark Kuan)	102
29	ANOVA for Dry Weight Set II (Mokara Chark Kuan)	103
30	ANOVA for Dry Weight Gain Set II (Mokara Chark Kuan)	103
31	ANOVA for Fresh Weight Set I (Dendrobium Jacky)	104
32	ANOVA for Dry Weight Set I (Dendrobium Jacky)	104
33	ANOVA for Dry Weight Gain Set I (Dendrobium Jacky)	105
34	ANOVA for Fresh Weight Set II (Dendrobium Jacky)	105
35	ANOVA for Dry Weight Set II (Dendrobium Jacky)	106
36	ANOVA for Dry Weight Gained Set II (Dendrobium Jacky)	106



LIST OF FIGURES

Figure		Page
1	Graph mean of survival of PLBs (Mokara Chark Kuan) vs. γ-irradiation doses	95
2	Graph mean of survival of PLBs (<i>Dendrobium</i> Jacky) vs. γ- irradiation doses	96



LIST OF PLATES

Plate	
1	Mokara Chark Kuan 'Pink' (Aranda Christine x Ascosenda Cholburi)
2	Dendrobium Jacky (Dendrobium Madam Vipa x Dendrobium Candy Stripe)
3	Agarose gel electrophoresis of amplified sequences from a RAPD reaction directed by primer OPAE-08 using DNA extracted from 7 Mokara Chark Kuan regenerants treated with different doses of γ-irradiation. Two novel bands of approximately 2.5 and 1.1 kb are indicated
4	Southern analysis using the band of approximately 2.5 kb (Plate 3) to probe a Southern blot of the same amplification product, a RAPD reaction directed by primer OPAE-08
5	Southern analysis using the band of approximately 1.1 kb (Plate 3) to probe a Southern blot of the same amplification product, a RAPD reaction directed by primer OPAE-08
6	Agarose gel electrophoresis of amplified sequences from a RAPD reaction directed by primer OPAE-11 using DNA extracted from 7 Mokara Chark Kuan regenerants treated with different doses of γ-irradiation. Two novel bands of approximately 1.2 and 0.9 kb are indicated
7	Southern analysis using the band of approximately 1.2 kb (Plate 6) to probe a Southern blot of the same amplification product, a RAPD reaction directed by primer OPAE-11
8	Southern analysis using the band of approximately 0.9 kb (Plate 6) to probe a Southern blot of the same amplification product, a RAPD reaction directed by primer OPAE-11



9	control (T ₀ ', T ₀) and irradiated (25 Gy, 40 Gy, 50 Gy, 80 Gy) Mokara Chark Kuan	71
10	Gel electrophoresis of Sac1 digested genomic DNA of control (T ₀ ', T ₀) and irradiated (25 Gy, 40 Gy, 50 Gy, 80 Gy) Mokara Chark Kuan	71
11	Southern analysis of EcoR1 digested genomic DNA of Mokara Chark Kuan using probe B (approx. 1.1 kb)	72
12	Southern analysis of Sac1 digested genomic DNA of Mokara Chark Kuan using probe B (approx. 1.1 kb)	72
13	Southern analysis of EcoR1 digested genomic DNA of Mokara Chark Kuan using pAL108 as probe	73
14	Southern analysis of Sac1 digested genomic DNA of Mokara Chark Kuan using pAL108 as probe	73
15	Gel electrophoresis of pAL108 and pAL202	74
16	Southern analysis of pAL108 and pAL202 using pAL202 as probe	75
17	Southern analysis of pAL108 and pAL202 using pAL108 as	75



LIST OF ABBREVIATIONS

AP : 2-amino-purine

bp : base pair

BU 5-bromo-uracil

BUdR : 5-bromo-deoxynridine

°C : degree(s) Celsius

cm : centimetre(s)

⁶⁰Co : cobalt-60

CTAB : cetyltrimethylammonium bromide

CV : coefficient of variation

DES : diethyl sulphate

DF : degree of freedom

DNA : deoxyribonucleic acid

EDTA : ethylenediaminetetraacetic acid

EMS : ethyl methanesulphonate

g : gram(s)

Gy : Gray(s)

Gy/s : Gray(s) per second

HCl hydrochloric acid

HNO₂ nitrous acid

hr(s) hour(s)

IPMS isopropyl methanesulphonate

kb : kilobase

LB : Luria-Bertani

LD lethal dose

M : molar(s)

MeV : megaelectron volt(s)

min : minute(s)

MINT : Malaysia Institute for Nuclear Technology Research

ml : mililitre(s)

mM : milimolar(s)

MNH N-nitroso-N-methyl urea

MNU : N-nitroso urea

ng : nanogram(s)

NaCl : sodium chloride

NaOH : sodium hyroxide

NH₂OH : hydroxylamine

nm : nanometre(s)

nM : nanomolar(s)

NPK sodium:phosphorous:potassium (ratio)

PCR polymerase chain reaction

pH : hydrogen ion concentration

PLBs protocorm-like bodies

psi pound(s) per square inch

PVP polyvinylpyrrolidone

RAPD random amplified polymorphic DNA

RCBD Random Complete Block Design

RNA : ribonucleic acid

rpm : revolution(s) per minute

SCARs sequenced characterised amplified regions

SDS : sodium dodecyl sulphate

sec : second(s)

SSC : saline-sodium citrate

TAE : Tris-acetate-EDTA

TE : Tris-EDTA

Tris-HCl Tris(hydroxymethyl) aminomethane hydrochloride

UV : ultraviolet

V : volt(s)

VW : Vacin and Went's formulation medium (modified)

μg : microgram(s)

μl microlitre(s)

 γ : gamma

Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Agricultural

Science.

EFFECTS OF GAMMA IRRADIATION ON IN VITRO CULTURES OF

SELECTED ORCHID HYBRIDS

By

ADRAIN LING CHIENG KUANG

February 1999

Chairman: Associate Professor Saleh bin Kadzimin, Ph.D.

Faculty: Agriculture

Mutation breeding is an alternative strategy to conventional breeding.

The physical mutagen, gamma (γ) radiation, has long been used in various

mutation breeding programmes. In this connection it is necessary to determine

the optimum dose before starting any mutation breeding programme.

In vitro cultured protocorm-like bodies (PLBs) of 2 orchid hybrids,

Mokara Chark Kuan and Dendrobium Jacky were used to study the effects of γ-

irradiation and to determine the optimum dose. The PLBs were irradiated with

 γ -rays at different doses of 0, 25, 50, 75, 100, 125 and 150 Grays (Gy) in Set I

and 0, 20, 40, 60 and 80 Gy in Set II of the experiment. A series of observations on the effects of γ -irradiation were recorded. Effects on genomic DNA were also observed.

The present study showed that irradiation resulted in various degrees of lethality to *in vitro* cultures of both *Mokara* Chark Kuan and *Dendrobium* Jacky. Decrease in survival with increasing doses of γ -irradiation was significantly observed. Optimum dose of γ -irradiation for *in vitro* cultures of *Mokara* Chark Kuan and *Dendrobium* Jacky were estimated to be in the range of 20-40 Gy and 60-70 Gy respectively.

Regeneration of plantlets was found to decline when higher irradiation doses were applied. Irradiation also has inhibited regeneration of *in vitro* cultures of both hybrids.

In general, fresh weight, dry weight and dry weight gain of *in vitro* cultures of *Mokara* Chark Kuan and *Dendrobium* Jacky decreased with increasing of γ -irradiation doses. *In vitro* cultures of *Mokara* Chark Kuan irradiated at 40 Gy were found to have higher fresh weight than those irradiated at 20 Gy but still lower than the control. Higher fresh weight, dry weight and

dry weight gain were observed in cultures of *Dendrobium* Jacky irradiated at 80 Gy.

In the present study, polymorphic DNA was detected by 2 random primers via polymerase chain reaction (PCR) between 7 DNA samples of *Mokara* Chark Kuan. Variations caused by repeated subculture and irradiation were detected by random amplified polymorphic DNA (RAPD) analysis. Homology search using 2 sequences (pAL108 and pAL202) from polymorphic bands as query sequences revealed close similarity to retrotransposons. No polymorphism could be detected by the primers used in RAPD analysis of *Dendrobium* Jacky DNA samples.

Abstrak tesis ini diserahkan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk penganugerahan ijazah Master Sains Pertanian.

KESAN RADIASI GAMMA KE ATAS KULTUR IN VITRO HIBRID ORKID TERPILIH

Oleh

ADRAIN LING CHIENG KUANG

Februari 1999

Pengerusi: Profesor Madya Saleh bin Kadzimin, Ph.D.

Fakulti : Pertanian

Pembiakan mutasi ialah satu strategi alternatif kepada pembiakan konvensional. Mutagen fizikal, sinaran gamma (γ), telah lama digunakan dalam pelbagai program pembiakan mutasi. Dalam hubungan ini, penentuan dos

optimum adalah penting sebelum memulakan sesuatu program pembiakan

mutasi.

Struktur sel berbentuk protokom (PLBs) daripada 2 hibrid orkid, iaitu

Mokara Chark Kuan dan Dendrobium Jacky yang dikultur secara in vitro telah

digunakan untuk mengkaji dos optimum dan kesan sinaran y ke atas kultur in

vitro. PLBs telah didedahkan kepada sinaran γ pada dos yang berlainan, iaitu 0, 25, 50, 75, 100, 125 dan 150 Gray (Gy) dalam Set I dan 0, 20, 40, 60 dan 80 Gy dalam Set II. Pemerhatian ke atas kesan-kesan radiasi telah direkodkan. Pemerhatian kesan radiasi ke atas DNA juga dilakukan.

Kajian ini menunjukkan bahawa radiasi γ menyebabkan beberapa tahap kematian kepada kultur *in vitro Mokara* Chark Kuan dan *Dendrobium* Jacky. Bilangan PLBs yang hidup terus menurun dengan peningkatan dos radiasi. Dos optimum untuk kultur *in vitro Mokara* Chark Kuan dan *Dendrobium* Jacky adalah masing-masing dianggarkan di antara 20-40 Gy dan 60-70 Gy.

Pertumbuhan semula anak pokok didapati merosot apabila dos radiasi yang lebih tinggi digunakan. Radiasi juga didapati merencatkan pertumbuhan semula anak pokok kedua-dua hibrid.

Secara amnya, berat basah, berat kering dan peningkatan berat kering kultur in vitro Mokara Chark Kuan dan Dendrobium Jacky didapati menurun dengan peningkatan dos radiasi γ. Walau bagaimana pun, rawatan 40 Gy bagi Mokara Chark Kuan didapati memberi berat basah yang lebih tinggi daripada rawatan 20 Gy tetapi masih kurang daripada kawalan. Berat basah, berat kering

dan peningkatan berat kering yang lebih tinggi telah diperhatikan pada rawatan 80 Gy bagi *Dendrobium* Jacky.

Dalam kajian ini, DNA polimorfik telah dikesan oleh 2 primer rambang menerusi 'polymerase chain reaction' (PCR) di antara 7 sampel DNA *Mokara* Chark Kuan. Variasi yang disebabkan oleh subkultur yang berulangkali dan radiasi telah dikesan oleh 'random amplified polymorphic DNA' (RAPD). Pencarian homologi ke atas 2 turutan (pAL108 and pAL202) daripada jalurjalur polimorfik telah menunjukkan darjah persamaan yang tinggi kepada retrotransposon. Tiada polimorfisma dapat dikesan pada sampel DNA *Dendrobium* Jacky dalam RAPD oleh primer yang digunakan.



CHAPTER I

INTRODUCTION

Orchid is one of the major cut flower types grown in numerous tropical countries. It has become an important export commodity in Malaysia, Singapore and Thailand contributing major shares in the floriculture industries of these countries. In Malaysia, orchids have contributed about 40% of the total value of cut flower production in 1995. With the increasing affluence of the local population and that of the importing countries, demand for orchids is expected to increase.

Orchid gains its popularity for its wide range of varieties producing an array of shapes, colours and sizes. However, many of the existing hybrids have been in the market for more than 15 years, with some more than 20. With ever changing tastes and preferences of consumers, there is an urgent need for new and better varieties of orchids. Nevertheless, creation of such varieties through conventional breeding and selection has always been difficult, costly and time-consuming, making it a venture less attractive to many. In addition, improving already established varieties through conventional means is often hampered by genetic sterility.

