



UNIVERSITI PUTRA MALAYSIA

CYTOGENETIC STUDIES OF MALAYAN GAUR (*Bos gaurus hubbacki*), SAHIWAL-FRIESIAN CATTLE AND THEIR HYBRID BACKCROSSES

**MAMAT HAMIDI KAMALLUDIN
FP 2009 29**

**CYTOGENETIC STUDIES OF MALAYAN GAUR (*Bos gaurus hubbaki*),
SAHIWAL-FRIESIAN CATTLE AND
THEIR HYBRID BACKCROSSES**

By

MAMAT HAMIDI KAMALLUDIN

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfilment of the
Requirements for the Degree of Master of Science**

October 2009



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

CYTOGENETIC STUDIES OF MALAYAN GAUR (*Bos gaurus hubbacki*), SAHIWAL-FRIESIAN CATTLE AND THEIR HYBRID BACKCROSSES

By

MAMAT HAMIDI KAMALLUDIN

October 2009

Chairman: Mohamad Hilmi Hj. Abdullah, PhD

Faculty: Agriculture

In early April 1983, a stranded Malaysian gaur (*Bos gaurus hubbacki*) mated with a group of Sahiwal-Friesian dairy heifers and this resulted in the birth of an inter-specific hybrid calf named SELEMBU. In this study, chromosome analysis of four Malayan gaur (*Bos gaurus hubbacki*), ten Sahiwal-Friesian cattle and seven hybrid backcrosses was undertaken to determine their karyotype status and differences.

The gaur and cattle has chromosome complements of $2n = 56$ and $2n = 60$, respectively. Two types of chromosomal arrangements of the backcrosses were observed; namely an intermediate to the parental species ($2n = 58$) and a cattle type ($2n = 60$). The backcrosses with $2n = 60$ shared similar chromosomal arrangement and banding characteristics of the cattle. Backcrosses with $2n = 58$ exhibited the non-homology of two submetacentric

and two acrocentric autosomes, where the variations were inherited from the parental species that possessed different karyotypes.

A comparison of the gaur with cattle, as the model of the ancestors of the modern bovids, showed structural and characteristic differences in their karyotypes. The gaur exhibited two pairs of submetacentric chromosomes and lacked two chromosome pairs, which had resulted from Robertsonian translocations during their karyotype evolution. Banded karyotypes revealed extensive similarities of chromosomes 1 and 2 of the Malayan gaur to the homologous acrocentric chromosomes of cattle. For the first time, chromosome 1 of the gaur was identified to contain an ancient origin inverted segment compared to cattle, which was homozygous in all the studied gaur samples. The intensity of the C-bands on chromosomes 1 and 2 suggested that the translocations occurred at different periods of time during the karyotype evolution.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Sarjana Sains

KAJIAN SITOGENETIK SELADANG (*Bos gaurus hubbacki*), LEMBU SAHIWAL-FRIESIAN DAN KACUKAN SELEMBU

Oleh

MAMAT HAMIDI KAMALLUDIN

Oktober 2009

Pengerusi: Mohamad Hilmi Hj. Abdullah, PhD

Fakulti: Pertanian

Pada awal bulan April 1983, seekor seladang (*Bos gaurus hubbacki*) yang terperangkap telah mengawan dengan sekumpulan lembu tenusu Sahiwal-Friesian dan menghasilkan anak kacukan yang dinamakan SELEMBU. Analisis kromosom empat ekor seladang, sepuluh ekor lembu Sahiwal-Friesian dan tujuh ekor hibrid kacukan balik telah dijalankan untuk mengenalpasti status kariotip dan perbezaan di antara spesies ini.

Seladang dan lembu masing-masing mempunyai pasangan kromosom $2n = 56$ dan $2n = 60$. Dua jenis penyusunan kromosom kacukan balik telah didapati iaitu pertengahan di antara spesies induk ($2n = 58$) dan jenis lembu ($2n = 60$). Kacukan balik dengan $2n = 60$ mempamerkan penyusunan kromosom dan sifat penjaluran yang sama dengan lembu. Kacukan balik dengan $2n = 58$ mempamerkan kromosom yang tidak sepadan iaitu dua submetasentrik dan dua akrosentrik, yang mana variasi tersebut diwarisi dari spesies induk yang mempunyai karotip yang berbeza.

Perbandingan antara seladang dan lembu sebagai model keturunan bovid moden, menunjukkan perbezaan struktur dan ciri-ciri dalam karotip. Seladang mempamerkan dua pasang kromosom submetasentrik dan kekurangan dua pasangan lain akibat translokasi 'Robertsonian' semasa evolusi karotip. Karotip berjalur membuktikan persamaan yang tinggi antara kromosom 1 dan 2 seladang dengan kromosom akrosentrik yang separa pada lembu. Buat pertama kali, kromosom 1 pada seladang dikenalpasti mempunyai segmen terbalik berbanding lembu pada setiap seladang yang dikaji. Intensiti jalur-C pada kromosom 1 dan 2 menandakan kemungkinan translokasi telah berlaku pada jangkamasa yang berbeza semasa evolusi karotip.

ACKNOWLEDGEMENTS

First and foremost, I would like to express my sincere gratitude to my advisors and guru, Assoc. Prof. Dr. Mohamad Hilmi Hj. Abdullah and Assoc. Prof. Dr. Ismail Idris in the Department of Animal Science, Faculty of Agriculture, UPM. Their keen interest, valuable guidance and continuous support have made this study and thesis materialized. I am grateful to Prof. Dino Di Berardino in University of Naples Federico II, Italy for his precious suggestions and assistance. I also indebted to the Universiti Putra Malaysia for allowing me the use of research facilities and to learn. My thanks also goes to Dr. Rafiah Hj. Aini and Dr. Noraihan Mohd Yassin of PTH Lenggong, Dr. Abas Mazni Othman of Agriculture-Biotechnology Institute (ABI), Dr. Zainal Zahari Zainuddin of Department of Wildlife and National Parks, staffs in PTH Ayer Hitam, PTH Lenggong and Pusat Konservasi Seladang Jenderak. My thanks to Cik Norbazlin Md Marham for her expert technical assistance, Assoc. Prof. Dr. Jothi Malar Panandam for being a very good teacher, Professor Zainal Aznam Mohd Jelani for his help and opinion on preparing this thesis. Special thanks to my mother, Zainun Mat Sab, for her continued believe in me and encouraged me to move forward. Last but not least, thanks to my wife, Fezah Othman for being my superwoman, and my daughters, Heidi Yasmine and Kasih Darwisyah for coloring the wall of my house.

I certify that an Examination Committee has met on 29th October 2009 to conduct the final examination of Mamat Hamidi Kamalludin on his Master of Science thesis entitled " Cytogenetic Studies of Malayan Gaur (*Bos gaurus Hubbacki*), Sahiwal-Friesian Cattle and Their Hybrid Backcrosses" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the student be awarded the Masters of Science.

Members of the Examination Committee were as follows:

Jothi Malar Panandam, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Siti Shapor Siraj, PhD

Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Tan Soon Guan, PhD

Professor
Faculty of Biotechnology and Biomolecule Science
Universiti Putra Malaysia
(Internal Examiner)

Mahani Mansor Clyde, PhD

Professor, Dato'
School of Environmental & Natural Resource Sciences,
Faculty of Science and Technology
Universiti Kebangsaan Malaysia
(External Examiner)

BUJANG BIN KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 15 January 2010

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Masters of Science. The members of the Supervisory Committee were as follows:

Mohamad Hilmi Hj. Abdullah, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Ismail Idris, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11th February 2010



DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

MAMAT HAMIDI KAMALLUDIN

Date: 18 December 2009

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xv
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	5
Zoological Position of the Gaur and Cattle	5
The Gaur	7
Karyotype of Cattle, Gaur and Their Hybrids	10
Chromosome Abnormalities	12
Translocation	12
Inversions	14
Interspecific Hybridization	15
Cytogenetics	18
Chromosome Banding	19
High Resolution Banding	23
3 MATERIALS AND METHODS	25
Animals	25
Lymphocyte Culture Method	26
Lymphocytes Synchronization and Late	
Incorporation of BrdU	27
Harvesting the culture	28
Slide Preparation	28
Giemsa (G) Banding Technique	29
R-banding (RBG) Technique	29
Constitutive Heterochromatin (C) Banding Technique	30
Construction of Karyotypes	31
4 RESULTS	32
Chromosome Number and Conventional Karyotype	32
G- and R-banded Karyotypes	40
C-banded Karyotype	49
Phenotypic observation of the selembu backcrosses	53



5	DISCUSSION	57
	Diploid Chromosome Distribution	57
	The Malayan Gaur	57
	Selembu Backcrosses	58
	Banded Karyotype Comparison	60
	Cytogenetic Homology Between Gaur And Cattle	62
	Robertsonian Translocation	62
	Inversions	63
	Identification of Sex Chromosomes / C banding	64
	Different Karyotype of a Female Malayan Gaur	67
	Interspecific Hybrids and Fertility	69
6	CONCLUSION	74
	REFERENCES	77
	APPENDIX I – Preparation of Chemicals	87
	APPENDIX II – Related Pictures	89
	APPENDIX III – Animal’s Identity and Their Chromosome Number	93
	BIODATA OF STUDENT	96
	LIST OF PUBLICATIONS	97

LIST OF TABLES

Table	Page
1. Distribution of the Malayan gaur.	10
2. Banding technique described by Sumner (1990) based on ISCN (1985).	21
3. Chromosome classification of Malayan gaur, Sahiwal-Friesian cattle and gaur x cattle backcrosses.	34
4. Possible number of meiotic elements and chromosome rearrangement in a nucleus of F ₁ hybrid gamete after chromosome segregation during the meiotic division.	73

LIST OF FIGURES

Figure	Page
1. Zoological Position of the Gaur and Cattle.	6
2. A family of Malayan Gaur (<i>Bos gaurus hubbacki</i>).	8
3. Representative Karyotype of Malayan Gaur (<i>Bos gaurus hubbacki</i>); $2n = 56$.	35
4. Representative Karyotype of Sahiwal-Friesian Cattle; $2n = 60$.	36
5. Representative Karyotype of Gaur X Cattle Backcross; $2n = 60$.	37
6. Representative Karyotype of Gaur X Cattle Backcross; $2n = 58$.	38
7. Karyotype of a Gaur with $2n = 58$.	39
8. Representative G-banded Karyotype of Gaur with a) $2n = 56$; b) $2n = 58$.	42
9. Representative G-banded Karyotype of Cattle $2n = 60$.	43
10. Representative G-banded Karyotype of Gaur x Cattle Hybrid Backcross with (a) $2n = 60$; (b) $2n = 58$.	44
11. Comparison of G-banded Chromosome and Ideogram 1, 2, 28 and 29 of Cattle (Courtesy of Leopoldo Iannuzzi) Against Chromosome 1 and 2 of Malayan Gaur.	45
12. Representative R-banded Karyotype of Gaur with $2n = 58$.	46
13. Representative R-banded Karyotype of Sahiwal-Friesian Cattle $2n = 60$.	47
14. Representative R-banded Karyotype of Gaur x Cattle Hybrid with (a) $2n = 60$; (b) $2n = 58$.	48
15. Representative C-banded Karyotype of Gaur with (a) $2n = 56$; (b) $2n = 58$.	50

16. Representative C-banded Karyotype of Sahiwal-Friesian cattle; $2n = 60$.	51
17. Representative C-banded Karyotype of Gaur x Cattle Hybrid Backcross with (A) $2n = 60$; (B) $2n = 58$.	52
18. Phenotype of a male selembu backcross (tag number: SL3) with $2n=60$	54
19. Phenotype of a male selembu backcross (tag number: SL5) with $2n=58$	55
20. Phenotype of a female selembu backcross (tag number:SL1) with $2n=58$	55
21. Phenotype of a female selembu backcross (tag number SL4) with $2n=60$	56
22. G-banded Comparison Between (a) Gaur Chromosome 1, (b) Cattle Chromosome 1, (c) Cattle Chromosome 29 and (d) Gaur Chromosome 1 with Inverted Segment.	64

LIST OF ABBREVIATIONS

BrdU	-	Bromodeoxyuridine (5-bromo-2-deoxyuridine)
C-bands	-	Constitutive Heterochromatin bands
DVS	-	Department of Veterinary Services
DWNP	-	Department of Wildlife and National Park
F1 Hybrids	-	Gaur x cattle hybrids
G-band	-	Giemsa bands
ISCN	-	International System for Human Cytogenetic Nomenclature
ISCNDA	-	International System for Cytogenetic Nomenclature of Domestic Animals
ISCNDB	-	International System for Chromosome Nomenclature of Domestic Bovids
IUCN	-	International Union for Conservation of Nature and Natural Resources
R-band	-	Reverse bands
RPMI 1640	-	Rosewell Park Memorial Institute 1640
SF	-	Sahiwal-Friesian

CHAPTER 1

INTRODUCTION

In early April 1983 at one of the Department of Veterinary Services (DVS) farm, Pusat Ternakan Haiwan Padang Hijau, Kluang, a stranded Malaysian gaur (*Bos gaurus hubbaki*), locally known as seladang mated with a group of Sahiwal-Friesian dairy heifers. This chance mating resulted, for the first time in Malaysian history, the birth of an inter-specific hybrid calf, which was named as SELEMBU (coined from two Malay words seladang meaning gaur and lembu meaning cattle). Soon after a few more hybrids were born until it was observed that the hybrid calves outgrew the dairy calves (Nor Azman, 1989), which indicated the potential of the animals as meat producer.

Malaysia was estimated to be at 121% and 114% self-sufficiency in poultry meat and eggs, respectively, in 2007. However, the country was only at 23.0% and 9.0% self-sufficiency in beef and mutton, respectively. In 2008, local production of beef was only 25.0% provisional of the domestic demand (DVS, 2009). Realizing that Malaysia was not self-sufficient in beef, cattle farming was emphasized so that more beef could be produce and to reduce the national import bill, thus saving the nation's foreign exchange.



Crossbreeding between *Bos indicus* for its adaptability and *Bos taurus* for its productivity is quite common in Malaysia. In 1982, the Malaysian government embarked on an importation program of Sahiwal-Friesian cattle to be used as dual-purpose animal *i.e.* for dairy and beef production. Sahiwal-Friesian bulls are animals with body weight of about 700kg and the females of about 400kg (Nor Azman, 1989). The colour of Sahiwal-Friesians is a blend of brown, yellow, light brown, white and black. The cattle genome is composed of 29 autosome pairs and two sex chromosomes. All autosomes are acrocentrics with small size differences (ISCNDB, 2000).

Hybridization between species is often arranged through human intervention and rarely in a natural manner. However, it may occur if closely related species share an overlapping habitat. Natural interspecies hybridization is considered as an important evolutionary process (Barton, 2001; Dowling and Secor, 1997). Uncontrolled hybridization as it occurs in nature may have had a significant impact on the formation of domestic breeds, but can also affect the genetic integrity of domestic and wild species.

Selembu or the interspecific hybrid of cattle and gaur has phenotypic features intermediate to both of the parents. However, it had been observed that the hybrids at Padang Hijau farm inherited most of the characteristic feature of the sire (gaur) particularly in the coat colour and the gradual changes of coat colour (Nor Azman, 1989). They were reported to have an

intermediate number of chromosomes ($2n=58$) compared to both parents while gaur and cattle chromosome complements are $2n=56$ and $2n=60$, respectively. The hybrid karyotype consists of 54 acrocentric autosomes, 2 submetacentric chromosomes and two sex chromosomes (Bongso *et al.*, 1988).

However, the fertility status of the hybrids was reported to be restricted to only female offspring and the males are sub-fertile (Nor Azman, 1989). Numerous studies provide evidence for the role of Robertsonian translocation or the centric fusion of chromosomes in reduced fertility (Dyrendhal and Gustavsson, 1979; Rangel-Figueiredo and Iannuzzi, 1993). Despite being phenotypically normal, translocation carriers experience a decrease in reproductive potential, predominantly due to the production of chromosomally unbalanced gametes. Due to the fertility problem of the male hybrids, the project of selembu breeding was abandoned and hybrids were kept with the dairy herd. The breeding record of the selembu therefore became unavailable. The remaining group was then transferred to a deer farm (Pusat Ternakan Haiwan) in Lenggong, Perak and kept freely in a fenced paddock in the farm.

A synthetic breed of beef cattle could be produced by controlled breeding involving the female gaur x cattle hybrids, which were reported to be fertile (Nor Azman, 1989) with cattle bulls. The backcrosses should then be

selected by cytogenetic screening to identify a balanced chromosome configuration for further breeding with bulls or heifers. This study was undertaken to investigate the karyotype (number and morphology) of the remaining Selembu and their parental species using conventional, giemsa (G-banding), reverse (R-banding), and constitutive heterochromatin (C-banding).

The main objectives of this study were:

1. To investigate the karyotype status of the remaining Selembu and their parental species.
2. To determine similarities and/or differences of genetic materials possessed by both species and their hybrids via their chromosome banding patterns.

CHAPTER 2

LITERATURE REVIEW

Zoological Position of the Gaur and Cattle

Bovidae and six other family of sub-order ruminantia belong to the zoological order of the artiodactyla or even-number toe ungulates (Vaughan *et al.*, 2000). The family is most diverse with 45 genera and 137 species worldwide. Under the genus *Bos*, there are five species namely *gaurus*; gaur, *grunniens*; yak, *javanicus*; banteng, *taurus* and *indicus*; cattle.

Bos taurus (taurine) and *Bos indicus* (zebu) are the major species of cattle. *Bos taurus* is mainly of temperate origin while *Bos indicus* is of Asian or African origin. There are fundamental anatomical differences justifying their separation. The hump over the withers and the presence of dewlap below the neck are the distinguishing characteristics of the zebu. The zebu also have larger ears, skin folds, shorter hair coat, more pigmented skin, longer legs and more angular body frame as compared to the taurine (Warwick and Legates, 1979). The *Bos taurus* are developed either as dairy or meat animal while the *Bos indicus* was utilized for triple purpose before mechanization. Apart from its purpose as beef and milk resources, the zebu was used as work animal for ploughing paddy fields or pulling carts.

The gaur was reported to have an extensive range from the hilly-forested country of the Indian peninsular, Burma to Indochina and down to Malaysia (Conry, 1989; Walker, 1964). Three subspecies of the gaur were classified by their geographical location, which are India-Nepal, Indochina and Malaysia-Thailand gaur, respectively (Byers *et al.*, 1995; Conry, 1981).

CLASS

Mammalia (warm blooded animals)

SUBCLASS

Eutheria

ORDER

Artiodactyla (mammals with even number toes)

SUBORDER

Ruminantia (true ruminants)

FAMILY

Bovidae (Hollow horn bovine)

SUBFAMILY

Bovinae

GENUS

COMMON NAME

Bison	Bison (Europe, North America)
Bos	Cattle, gaur, banteng, Yak
Boselaphus	Nilgai (Asia)
Bubalus	Asiatic buffalo (Asia)
Syncerus	African Buffalo (Africa)
Taurotragus	Eland (Africa)
Tetracerus	Four-horned antelope (Asia)
Tragelaphus	Bushbuck, Nyala, Kudu, Bongo (Africa)

Figure 1. Zoological Position of the Gaur and Cattle.

(Adapted from Vaughan *et al.*, 2000)

The Gaur

The gaur is a large muscular animal with sturdy limbs and a striking muscular ridge that slope down to the middle of the back. It is also known as Indian bison or locally as Seladang and can be found in Bangladesh, Cambodia, China, India, Malaysia, Myanmar, Nepal, Thailand and Vietnam. They are the native cattle of the rainforests of Southeast Asia, and one of the many wild cattle species currently listed as vulnerable and endangered (IUCN, 2008).

Previously it was deliberated that there are three subspecies of gaur: Indian gaur (*Bos gaurus gaurus*), Indochinese gaur (*Bos gaurus readei*) and Malayan gaur (*Bos gaurus hubbacki*) (Lekagul and McNeely, 1977). Recently, the gaur has been recognized with three subspecies: *Bos gaurus gaurus* (India, Nepal), *Bos gaurus laosiensis* (Myanmar to China) and *Bos gaurus hubbacki* (Malaysia, Thailand) by the Asian Wild Cattle Conservation Assessment and Management Plan (Byers *et al.*, 1995). These animals were classified as endangered species and vulnerable by the International Union for the Conservation of Nature and Natural Resources (Simon, 1966). The global population of wild gaur ranges from 13,000 to 30,000 (IUCN, 2008; Nguyen *et al.*, 2007).



Figure 2. A family of Malayan gaur (*Bos gaurus hubbacki*).

As with other bovidae, chromosome studies indicated that gaur evolved from a wild ancestor from which the domestic taurine and zebu cattle originated (Buckland and Evans, 1978; Wurster and Benirschke, 1968). Centric fusion between chromosomes 2 and 28 of the ancestral cattle karyotype of 58 acrocentric chromosomes gave rise to the Indian gaur karyotype ($2n=58$), consisting of 27 pairs of acrocentric and a pair of submetacentric autosomes (Mastromonaco *et al.*, 2004).

The Malayan Gaur (*Bos gaurus hubbacki*), the second largest animal in Malaysia can weigh up to 1000kg for the male and 500-700kg for the female (Bongso *et al.*, 1988; Ebil, 1982). Huge head, massive body and sturdy limbs are the characterization of vigor and strength of the gaur. It was estimated

that there were 472 individuals in 1980 (DWNP, 1981) and a minimum population of 273-333 based on data collected in 1994 to 2005 (DWNP, 2005). This impressive largest wild cattle which live in isolated and fragmented populations mostly in Pahang, Perak, Kelantan, Terengganu and Taman Negara inhabit forests near to the foothills (Ebil, 1982). Studies indicated that the number was globally declining, may be due to habitat loss, poaching and diseases from domestic cattle (Conry, 1981; Conry, 1989; DWNP, 2005; IUCN, 2008; Nguyen *et al.*, 2007) and Malaysia has the most rapidly declining population (IUCN, 2008).

Developments including agricultural activities have forced the animal from their preferred areas to marginal parts of the habitat (Conry, 1989; Ebil, 1982). This may support the interesting incident in 1983 at southern part of Peninsular Malaysia where a wild gaur was reported to mate with a group of domestic cattle in a farm. The distribution of Malayan gaur is shown in Table 1, with the biggest population being in the state of Pahang.