



UNIVERSITI PUTRA MALAYSIA

**EVALUATING AND TESTING OF A POTENTIAL DNA VACCINE
AGAINST VIBRIO CHOLERAЕ**

LAMA ABDEL QADER MOH'D HAMADNEH

FPSK (M) 2003 2

**EVALUATING AND TESTING OF A POTENTIAL DNA VACCINE AGAINST
*VIBRIO CHOLERA***

BY

LAMA ABDEL QADER MOH'D HAMADNEH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfillment of the Requirements for the Degree of Master of Science**

June 2003



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirements for the degree of Master of Science

**EVALUATING AND TESTING OF A POTENTIAL DNA VACCINE AGAINST
*VIBRIO CHOLERAE***

By

LAMA ABDEL QADER MOH'D HAMADNEH

June 2003

Chairman: Associate Professor Dr. Rozita Rosli

Faculty: Medicine and Health Sciences

Although it has been more than 100 years since the first attempt to produce a cholera vaccine was made, an effective cholera vaccine has yet to be developed. In this study, the level of protection produced by a potential DNA vaccine (pVax/ctxB) was tested against the *ctxB* toxin of *Vibrio cholerae* on Balb/c mice. First, the intramuscular vaccination method was validated using pCMV plasmid that encodes HbsAg, which was detected 5 days after the injection into the tibial muscle. Next, 4 groups of mice were intramuscularly injected with either the pVax/ctxB vaccine construct or pVax1 as the negative control. The first and second groups received 2 injections spaced 3 weeks apart, while the other two groups were given 3 injections spaced 3 weeks apart. This was then followed by challenging the mice with 10^5 or 10^7 cfu/ml/mouse from clinical isolates of *V. cholerae* after 3 weeks of the last injection. Antibody levels for both IgG and serum IgA were monitored using ELISA, and showed high production of IgG after the first booster injection with no significant change of IgA levels. However, after the second



booster injection, the antibody levels for both IgG and IgA declined. This was accompanied by the death of 2 mice in the first vaccinated group, and all the mice in the control group after the bacterial challenge with 10^7 cfu/ml/mouse. In the second group, none of the mice survived in both vaccinated and control groups. The bacterial challenge using 10^5 cfu/ml/mouse failed to induce the death in all the groups.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

PENILAIAN DAN PENGUJIAN POTENSI VAKSIN DNA TERHADAP *VIBRIO CHOLERAE*

Oleh

LAMA ABDEL QADER MOH'D HAMADNEH

Jun 2003

Pengerusi: Profesor Madya Dr. Rozita Rosli

Fakulti: Perubatan dan Sains Kesihatan

Walaupun setelah lebih dari pada 100 tahun semenjak percubaan pertama menghasilkan vaksin taun dilakukan, namun sehingga kini penghasilan vaksin yang benar-benar efektif masih belum menampakkan hasil. Bagi kajian ini pula, tahap ketahanan yang dihasilkan oleh calon vaksin DNA (pVax/ctxB) telah diuji menentang toksin ctxB dari *Vibrio cholerae* dengan tikus Balb/c sebagai hos. Buat permulaan, teknik vaksinasi intramuscular telah di sahkan dengan menggunakan plasmid pCMV yang mengkodkan HbsAg yang mana telah dikesan setelah disuntik ke otot tibial 5 hari kemudian. Empat kumpulan tikus telah disuntik menggunakan teknik di atas samada dengan vaksin pVax/ctxB atau pVax sebagai kawalan negatif. Kumpulan pertama dan kedua menerima 2 suntikan setiap 3 minggu manakala 2 kumpulan lagi 3 suntikan bagi setiap 3 minggu. Langkah berikutnya, setiap kumpulan ditentang dengan 10^5 atau 10^7 cfu yang telah dipencilkan secara klinikal dari *V.cholera*. Paras antibodi bagi kedua-dua IgG dan serum IgA dipantau menggunakan ELISA. Keputusan ELISA menunjukkan paras

antibodi IgG tinggi sementara tiada perubahan ketara penghasilan paras IgA setelah suntikan booster pertama. Setelah suntikan booster kedua, paras antibodi IgG dan IgA didapati menurun. Keadaan ini diikuti dengan kematian 2 ekor tikus dari kumpulan pertama yang divaksin dan kematian kesemua tikus kumpulan kawalan setelah ditentang dengan 10^7 cfu *V. cholerae* yang virulen. Bagi kumpulan kedua, kesemua tikus (vaksin dan kawalan) mati. Tentangan bakteria 10^5 gagal untuk menyebabkan kematian bagi kesemua kumpulan.

Acknowledgements

My greatest and ultimate debt and gratitude is due to Allah, the Most Beneficent and the Most Merciful. May He pardon and forgive my weaknesses and endow me with knowledge and help.

My deepest gratitude to Associate Professor Dr. Rozita Rosli for giving me the opportunity to pursue my master's degree under her supervision, and for her guidance, suggestions and encouragement throughout the project. I am also grateful to my co-supervisors Associate Professor Dr. Mariana Nor Shamsudin and Associate Professor Dr. Abdul Rahman Omar, for their help, support and valuable discussions.

I thank my friends in the Molecular Genetics Laboratory; Shariza, , Suhaili, Thilaka, Nazefah, Rohayu, Shaban, Farouk, Kak Nurma and specially Kak Norshariza for being my family in Malaysia. To Syahrilnizam Abdullah, thank you for all the help and patience. I also thank Kak Wan, Neela, colleagues and staffs of Microbiology Laboratory for their help.

I am grateful to my family for all the support, encouragement and love they gave me and for raising me the way I am today. Finally, I am most grateful to my husband Imad, and to my son Yazan for all the patience and understanding.



I certify that an Examination Committee met on 20th June 2003 to conduct the final examination of Lama Abdel Qader Moh'd Hamadne on her Master of Science thesis entitled "Evaluating and Testing of a Potential DNA Vaccine against *Vibrio cholerae*" in accordance Universiti Pertanian Malaysia (Higher Degree) act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Datin Farida Fatema @ Farida Jamal, Ph.D.

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Rozita Rosli, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Mariana Nor Shamsudin, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Abdul Rahman Omar, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.
Professor/ Deputy Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date: **21 JUL 2003**

This thesis submitted to the Senate of Universiti Putra Malaysia and was accepted as fulfillment of the requirements of the degree of Master of Science. The members of the Supervisory Committee are as follows:

Rozita Rosli, Ph.D.


Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Mariana Nor Shamsudin, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Abdul Rahman Omar, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.
Professor/ Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date: **15** AUG 2003

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



LAMA ABDEL QADER MOH'D HAMADNEH

Date: 26/7/03

TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL SHEETS	vii
DECLARATION	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xv
CHAPTER	
1 INTRODUCTION	1
Objectives	4
2 LITERATURE REVIEW	5
Cholera	5
History	5
Disease	6
Mode of Transmission	7
Incubation Period and Infectious Dose	7
Age and Susceptibility	7
Treatment	8
<i>Vibrio cholerae</i>	9
Properties	9
Reservoir	10
Classification	10
Pathogenesis and Virulence Factors	12
Cholera Toxins	13
Lipopolysaccharides	13
Enterotoxins	13
Immunity to Cholera	15
Cholera Vaccines and their Drawbacks	15
DNA Vaccines	17
Early History of DNA Vaccines	17
DNA Plasmids	18
Routes of Vaccination	19
Adaptive Immune Responses	21
Mechanism of Immunization by DNA Vaccines	22
Induction of Humoral Immune Response	24
Induction of Cellular Immune Response	25
Longevity of Immunity Raised by DNA Vaccines	26



Modification of the Immune Response	27
CpG Motifs	27
Method and Location of Vaccination	28
The Encoded Antigen	29
Immunization Regimen	29
Genetic Adjuvants	30
Additional Adjuvants	31
Advantages of DNA Vaccines	31
3 MATERIALS AND METHODS	
Validation of the Injection Technique	34
HBsAg Reconstitution	35
Transformation	35
Confirming the Transformation	36
Miniprep pCMV-S Purification	36
Plasmid Analysis	37
Spectrophotometer Reading	37
Restriction Digest Analysis	37
Gel Electrophoresis	38
Preparation for the Injection	38
pCMV-S Purification Using Endofree Plasmid Maxi Kits	39
Preparation of the Final Solution of DNA Plasmid for Injection	40
DNA Vaccination Using Intramuscular Injection	41
Experimental Model	41
Anesthesia	41
Intramuscular Injection of the DNA Vaccine	42
Hepatitis B Surface Antigen Detection	42
Testing pVax/ctxB Plasmid <i>in Vivo</i>	43
Plasmid Purification Using EndoFree Plasmid Giga Kits	45
Preparation of Plasmids for Injection	46
Plasmids Injection in the Mice	47
Blood Samples	47
Bacterial Challenge	47
<i>Vibrio cholerae</i> Strengthening in Mice	48
<i>Vibrio cholerae</i> Growth Curve Determination	48
Re-isolation of <i>Vibrio cholerae</i>	49
Enzyme Linked Immunosorbant Assay	49
ELISA Optimization	49
Sera Testing for Total IgG Antibody Production	51
IgA Determination	52
Mice Injection with ctxB Antigen	52
Optimization of ELISA Protocol	52
Sera Analysis for the IgA	53
Statistical Analysis	54



4	RESULTS	55
	Validation of the Intramuscular Injection	57
	Bacterial Challenge	62
	Bacterial Challenge with Clinical Isolates of <i>V. cholerae</i>	63
	ELISA Optimization for IgG Detection	65
	Serum IgA Determination by ELISA	70
5	DISCUSSION	80
	Validation of the Injection Technique	80
	<i>Vibrio cholerae</i> Bacterial Challenge	83
	Enzyme Linked Immunosorbant Assay	84
	Potential Activity of the CtxB Antigen	85
	IgG and IgA Antibody Production	86
	Why not Oral Vaccination?	88
6	FUTURE DIRECTION	90
7	CONCLUSION	92
	REFERENCES	94
	APPENDIX	110
	VITA	111



LIST OF TABLES

Table		Page
1(a).	IgG ELISA optimization for the volume and concentration of ctxB antigen solution using different dilutions of the primary antibody.	66
1(b)	IgG ELISA optimization for the volume and concentration of ctxB antigen solution using washing buffer as the negative control.	67
2(a)	IgG ELISA optimization for the secondary antibody dilution	68
2(b)	IgG ELISA optimization for the secondary antibody dilution using washing buffer as the negative control.	69
3(a)	Serum IgA ELISA optimization for ctxB concentration and volume	71
3(b)	Serum IgA ELISA optimization for ctxB concentration and volume using washing buffer as the negative control.	72
4(a)	Serum IgA ELISA optimization for the secondary antibody dilution.	73
4(b)	Serum IgA ELISA optimization for the secondary antibody dilution.	74



LIST OF FIGURES

Figure		Page
1	Graph of the injection and challenge patterns. (a) mice injected with pVax/ctxB and (b) mice injected with pVax 1, the stars refer to the number of injections	44
2	Restriction enzyme analysis of the transformed cells containing pCMV-S plasmid	56
3	Anesthesia using Sodium Pentobarbital	58
4	Intramuscular injection of plasmid	59
5	Blood sampling by cardiac puncture	60
6	HBsAg detection using Immunocomb® II kit	61
7	Growth curve for <i>Vibrio cholerae</i>	62
8	Re-isolation of <i>Vibrio cholerae</i> from the mice after challenge	64
9	IgG antibody levels in the vaccinated and the control groups after one booster injection	76
10	Serum IgA antibody levels in the vaccinated and the control groups after one booster injection	77
11	IgG antibody levels in the vaccinated and the control groups after two booster injection	78
12	Serum IgA antibody levels in the vaccinated and the control groups after one booster injection	79



LIST OF ABBREVIATIONS

<i>Ace</i>	Accessory cholera enterotoxin
ACF	Accessory colonization factor
AIDS	Acquired immune deficiency syndrome
APC	Antigen presenting cells
BSA	Bovine serum albumin
cAMP	Cyclic adenosine 51-monophosphate
<i>Cep</i>	Core encoded pilin
CpG	Cytosine-phosphate-guanosine
CPS	Capsular polysaccharide
CTL	Cytolytic T lymphocytes
<i>Ctx</i>	Cholera enterotoxins
DNA	Deoxy ribonucleic acid
EDTA	Ethylene Diamine tetra acetic acid
ELISA	Enzyme linked immunosorbant assay
GM-CSF	Granulocyte-monocyte colony stimulating factor
HA	Haemagglutinin
HBsAg	Hepatitis B surface antigen
HIV	Human immunodeficiency virus
HRP	Horseradish peroxidase
H ₂ SO ₄	Sulfuric acid
ID	Intradermal
Ig	Immunoglobulin



IL	Interleukin
IM	Intamuscular
INF	Interferon
KCl	Potassium chloride
LB	Lauria-Bertani
LPS	Lipopolysaccharides
LT	Heat labile enterotoxins
MFRHA	Mannose-fucose-resistant hemagglutinin
MHC	Major histocompatibility complex
mRNA	Messenger ribonucleic acid
MSHA	Mannose-sensitive-hemagglutinin
NaCl	Sodium chloride
NaHCO ₃	Sodium bicarbonate
NK	Natural killer
NP	Nucleoprotein
OMP	Outer membrane proteins
ORS	Oral rehydration salts
PBS	Phosphate buffer saline
PLG	Poly (lactide-coglycolide)
TCBS	Thiosulfate-citrate-bile salts-sucrose.
TCP	Toxin-coregulated pili
Th	T helper
TMB	Tetramethylbenzidine



WHO World Health Organization

Zot Zonula occludens toxin



CHAPTER 1

INTRODUCTION

Around 200 years ago, immunization against infectious diseases was started when Edward Jenner published his method of preventing smallpox in 1798 (1). Although it took nearly 100 years before the appearance of the next vaccine in 1880, vaccine development was rapid, where in 1884, the first attempt to produce a parenteral cholera vaccine of broth cultures of *Vibrio cholerae* was reported by Ferran (2). However, after more than 100 years of this first vaccination attempt, no effective cholera vaccine is yet available to date.

The cholera disease can spread easily by water and food contamination, causing severe diarrhea and death among people living with poor sanitary facilities. Wars and political unrest, climate changes and natural catastrophes, increase the number of people under the threat of cholera epidemics. Furthermore, these conditions are still prevailing in many parts of the world. On the other hand, the increasing number of geographical areas becoming endemic for cholera reflects a failure of implementation of control measures. Water and electrolytes replacement therapy is not fast enough during times of outbreak, and misguided use of antibiotics has led to the emergence of multi-resistant strains, some of which were found to be highly virulent (3).

As of September 2002, 106,547 cases with 3,155 deaths were officially reported to the World Health Organization (WHO) (4). In the year 2001, Malaysia reported 557 cases with 11 deaths. From a total of 184,311 cases 2,728 deaths were reported from the whole world (5).

The need for an effective cholera vaccine is urgent to produce high level of protection in people living in high-risk areas. A parenteral vaccine based on inactivated *V. cholerae* O1 has been available for more than 40 years. Nevertheless, this vaccine is not recommended any more by WHO, since its protective efficacy is modest, of short duration and it does not prevent the transmission of *V. cholerae* (3).

At present, new cholera vaccines are under development, and 2 oral vaccines are already available internationally, WC/rBS and CVD103-HgR, which conferred good protection in adults. But thus far, all the vaccines being used give only 50% protection after 6 months of immunization together with no sustained protection in children under 2 years (3).

In early 1990's, the ability of inserted genes into plasmid vectors was studied to prompt an immune response; surprisingly the immunity elicited was strong enough to protect against infectious diseases. The immune response produced by these vaccines, which are termed as DNA vaccines (unlike that produced by conventional vaccines) can stimulate both humoral and cellular immune responses. The immune response produced was found to be long lasting and it might overcome the deficits of

the conventional vaccines used nowadays. In addition, new protection against diseases like AIDS, malaria, and hepatitis C can be provided.

The field of DNA vaccines is developing rapidly. Today, many clinical trials are conducted to test the efficacy of DNA vaccines against HIV, hepatitis B and some tumors such as B-cell lymphomas. Despite that, any new vaccine should be first tested on small animals to detect the immunogenicity of the antigen expressed, followed by testing its protective properties against the challenge. In addition, before proceeding to human clinical trials, the vaccine should be tested on higher animals to adjust the dose and the boosting pattern.

In an attempt to develop a cholera DNA vaccine, the gene encoding for the B subunit of the enterotoxin AB was cloned in pVax1 plasmid vector in a previous study (6), followed by successful *in vitro* expression of the antigen using COS-7 cell line.

Hence, this study focuses on the next step, in which the ability of this potential vaccine to be expressed and elicit an immune response *in vivo* is tested, or in other words to test the antigen's immunogenicity. *In vivo* tests based on measuring the level of antibody production and the protection level against the challenge with clinical isolates of *Vibrio cholera* after intramuscular immunization of this potential cholera vaccine were conducted using female BALB/c mice.

Objectives

- (1) To validate the intramuscular injection technique using pCMV-S plasmid that carries HBsAg in Balb/c mice.
- (2) To test the cholera DNA vaccine (pVax/ctxB) intramuscularly by measuring the levels of both IgG and IgA produced after each vaccination and compare it to a negative control plasmid (pVax1) without any insert.
- (3) To determine the efficacy of pVax/ctxB in conferring protection or immunity against the challenge with either 10^5 or 10^7 cfu/ml/mouse of clinical isolates of *Vibrio cholerae*.

CHAPTER II

LITERATURE REVIEW

Cholera

History

As early as the time of Hippocrates and Galen, cholera has sporadically affected humans all over the world where some records from this time described cholera-like symptoms. However, modern knowledge about cholera started in the beginning of 19th century, where the English physician John Snow in 1849 indicated the importance of water as the carrier of the disease. In 1883, Robert Koch succeeded to isolate *Vibrio cholerae* from the intestinal discharges of cholera patients.

Cholera has smoldered in an endemic fashion on the Indian subcontinent for centuries, from where the first long-distance spread of the disease to Europe and the Americas began in 1817 and by the early 20th century, six waves of cholera had spread all over the world. The 7th cholera pandemic caused by the *El Tor* biotype was started in 1961 from Indonesia and spread rapidly to Asia, Europe, Africa and finally in 1991 to Latin America, which was free of cholera for more than a century. The 7th pandemic has not receded; on the contrary, cholera has now become endemic in many parts of the world (7). Also, since 1992, *V. cholerae* O139, a new and more virulent serogroup variant of *El Tor* biotype has spread to many parts of Asia (3).

Disease

Cholera is an acute gastrointestinal disease where the production of diarrhea is the main symptom. The disease can range from mild in most cases with less than 1% mortality to a severe life threatening infection termed as cholera gravis with more than 50% mortality (3, 8). The symptoms might appear as sudden with profuse watery diarrhea or there can be some premonitory symptoms like anorexia, abdominal discomfort and simple diarrhea.

Initially the stool passed is brown, but soon it assumes a pale gray color with an inoffensive, slightly fishy odor. Mucus in the stool imparts its characteristic rice water appearance. Resultant water and electrolyte loss leads to thirst, muscle cramps, weakness, and sunken eyes. In cholera gravis, the rate of water loss may reach 1 L/hr leading to tachycardia, hypotension and vascular collapse due to dehydration (9, 10). If untreated severe metabolic acidosis with potassium depletion, anuria, circulatory collapse and cyanosis can occur, leading to death.

Major alterations in mental status are uncommon in adults; the patient usually remains well oriented but apathetic, while hypoglycemia, coma and convulsions might occur in children. After recovery, very small minority of patients (< 1%) continue to carry the pathogen in the gallbladder and excrete it with stools; however, most patients are free after about 2 weeks (10).

Mode of Transmission

Cholera infection usually begins with the ingestion of contaminated food or water. Food can buffer the acidity of the stomach thus decreasing the infecting dose of the bacteria as well as providing an ideal culture medium (11). Furthermore, sea food can acquire the bacteria from the environmental sources and causes outbreaks or sporadic cases specially if it is uncooked or partially cooked (12).

Incubation Period and Infectious Dose

The incubation period can range from a few hours to 5 days, usually 2-3 days and it is dependent in part on the inoculum's size (13), where 10^3 of bacteria in the intestine is enough to start the infection. It is estimated that around 10^{11} bacteria are required as an infective dose in normal gastric acidity individuals (14).

Age and Susceptibility

In endemic areas, the disease is concentrated more in children aged 2 to 9 and in women in their child bearing years (15-35 years) where there is a decrease in immunity and the exposure to the environment is higher. On the other hand, infants under 1 year of age are protected because of breast feeding (15). Individuals with blood group O are at higher risk for cholera gravis due to El Tor biotype and O139 *Vibrio* although the mechanism responsible for this difference is not known (16); also, individuals with gastric achlorhydria have a higher risk for getting the disease.