



UNIVERSITI PUTRA MALAYSIA

**THE EFFICACY OF GARLIC IN ALLEVIATING PHENANTHRENE
AND CHRYSENE INDUCED PULMONARY INJURY IN RATS**

WAN MAZLINA BT. MD. SAAD

FPV 2003 10

**THE EFFICACY OF GARLIC IN ALLEVIATING PHENANTHRENE AND
CHRYSENE INDUCED PULMONARY INJURY IN RATS**

By

WAN MAZLINA BT. MD. SAAD

Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of Requirements for the Degree of Master of Science

November 2003



DEDICATION

This thesis is dedicated with appreciation and love to my late mak (Mawanchik Pawan), late bahamad (Md. Saad Abdullah), Bapak, Mamanah, Inda, Ghoi, Mamanoin, Pak E, Ser, Batah, Bedah, Ain ul, Chek Sha and Kak Siti who are always there for me with full support and courage without which I will not finish this course.

“Thank You”

-Mali-



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

THE EFFICACY OF GARLIC IN ALLEVIATING PHENANTHRENE AND CHRYSENE INDUCED PULMONARY INJURY IN RATS

By

WAN MAZLINA BT. MD. SAAD

November 2003

Chairman: Associate Professor Noordin Mohamed Mustapha Ph.D.

Faculty : Veterinary Medicine

Air pollution specifically referring to haze consists of numerous particles including polycyclic aromatic hydrocarbons (PAH). These particles consist of various compound including phenanthrene (Phen) and chrysene (Chry). Both of these are among hazardous compounds which are found during the haze periods back in 1997 in Malaysia. This situation had a great impact on health and also on the economic status of the country as well. In finding strategies and bringing about a remission of noxious haze effects in humans, acute and chronic exposure to Phen, Chry and their combination were studied in rats.

The acute exposure studies were conducted to evaluate apoptosis in the lung of rats following with Phen, Chry and Phen+Chry. Rats not receiving any treatment served as control while those administered with Phen, Chry and Phen+Chry were instilled at the dose of 14.87ng (7.4 μ l), 7.07ng (3.5 μ l) and 21.94ng (10.9 μ l) respectively. Animals in control and treated group were euthanised at 1, 8 and 24 hours post instillation (p.i.).



Apoptosis detection was made on H&E stained paraffin-embedded sections, terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling analysis (TUNEL) and DNA laddering of lung samples. Selected proteases were also examined from lung lavage of all rats.

Apoptosis were observed in pneumocytes and bronchial epithelium of rats euthanised at the specific time p.i and the percentage of apoptotic cells increased with time, especially in 24 hours p.i. which were highest for group especially the combination group ($p < 0.05$). This was confirmed by TUNEL analysis and DNA laddering. As for proteases, there was also elevation in the content of total protein, number of neutrophil and level of elastase. This activity also showed increased with time as compared to normal.

The chronic exposure study was conducted on changes in the lung of Phen, Chry and Phen+Chry. These rats were kept for three months to assess histopathologic findings, immune response, level of marker enzyme, proteases and the effects of raw garlic as an anti-tumour factor.

Rats from the control group, Phen, Chry and Phen+Chry were daily fed with commercial rat pellet while rats from garlic (G), Phen+G, Chry+G and Phen+Chry+G were fed with rat pellet mixed with garlic at the rate of 80 mg/kg body-weight/rat/day.

From the observation rats from Phen, Chry and Phen+Chry group showed growth disturbances in pneumocytes and bronchial epithelium including pulmonary mechanisms.

Morphological changes including hyperplasia and metaplasia were among the findings in treatment groups. Necrosis was also detected in these groups. Rats induced with Phen and Chry particles showed a low level of IgG and IgA and alveolar macrophage activity in the lung. Furthermore, rats treated with these particles showed an elevation in glutathione-S-transferase and protease activity. Rats treated with garlic had no significant lesions and marked elevation in pulmonary defense mechanism.

Lastly, acute and chronic exposure to environmental pollutants such as Phen and Chry incites injury to the lung. Garlic has shown to be a great potential in alleviating chronic injury induced by Phen and Chry in the treated rats.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**KEBERKESANAN BAWANG PUTIH DALAM MENGATASI KEROSAKAN
PULMONARI AKIBAT INDUS PHENANTHRENE DAN
CHRYSENE PADA TIKUS**

Oleh

WAN MAZLINA BT. MD. SAAD

November 2003

Pengerusi: Prof. Madya Noordin Mohamed Mustapha Ph.D.

Fakulti : Perubatan Veterinar

Pencemaran udara atau lebih dikenali sebagai jerebu mengandungi pelbagai bahan berbahaya termasuk polisiklik aromatik hidrokarbon (PAH). PAH ini terdiri daripada pelbagai komponen termasuk phenanthrene (Phen) dan chrysene (Chry). Kedua-dua komponen ini adalah antara bahan pencemaran berbahaya yang terdapat dalam jerebu tahun 1997 di Malaysia. Jerebu 1997 bukan sahaja memberi kesan terhadap tahap kesihatan malah juga mempengaruhi keadaan ekonomi negara pada masa itu. Dalam mencari strategi dan kesannya terhadap manusia; dua pendedahan iaitu akut dan kronik dijalankan ke atas tikus untuk Phen, Chry dan gabungan kedua-duanya.

Kesan akut dijalankan untuk menilai bilangan apoptosis didalam peparu tikus yang diberi suntikan Phen, Chry dan Phen+Chry. Tikus yang tidak diberi sebarang suntikan dianggap sebagai kawalan manakala kumpulan-kumpulan tikus lain disuntik Phen sebanyak 14.87 ng (7.4 μ l), Chry 7.07 ng (3.5 μ l) dan 21.94 ng (10.9 μ l) untuk kumpulan Phen+Chry.



Tikus-tikus ini diberi pendedahan selama 1, 8 dan 24 jam. Apoptosis dapat dikesan pada peparu tikus yang diberi rawatan diatas melalui keratan histopatologi yang diwarnakan dengan H&E, TUNEL dan pengestrakan DNA. Kesan protease juga dikaji melalui basuhan bronkoalveolar dari kumpulan Phen, Chry dan Phen+Chry.

Apoptosis dapat dikesan pada tisu pneumosit dan epitelium bronkiol pada masa pendedahan tertentu. Kesan apoptosis meningkat bersama masa dan kesannya dapat dilihat dengan jelas pada pendedahan selama 24 jam untuk semua kumpulan ($p < 0.05$). Keputusan ini dikukuhkan lagi melalui analisis TUNEL dan ekstrak DNA. Peningkatan secara keseluruhan untuk kumpulan Phen, Chry dan Phen+Chry juga dapat diperhatikan bagi kandungan protein keseluruhan, neutrofil dan elastase. Peningkatan ini adalah lebih tinggi berbanding kumpulan tikus kawalan.

Bagi pendedahan kronik (tiga bulan) pula, terdapat pelbagai perubahan pada peparu tikus yang disuntik. Tikus-tikus dipelihara selama tiga bulan untuk kajian histopatologi, gerakbalas sistem keimunan, tahap sasaran enzim, tahap protein dan juga kesan bawang putih sebagai faktor anti-tumor.

Tikus dari kumpulan kawalan, Phen, Chry dan Phen+Chry diberi makanan komersial manakala tikus dari kumpulan bawang putih (G), Phen+G, Chry+G dan Phen+Chry+G diberi makan yang dicampur dengan bawang putih pada kadar 80 mg/kg berat/tikus/hari.



Dari kajian yang dijalankan, dapat disimpulkan bahawa terdapat gangguan pertumbuhan pada sel pneumosit dan epitelium bronkial dimana gangguan mekanisme pulmonari juga turut terlibat. Perubahan dari segi morfologi seperti hiperplasia dan metaplasia adalah antara gangguan yang terdapat pada sel epitelium paru tikus. Kesan ini juga dapat dilihat pada keratan histologi. Nekrosis juga kelihatan pada kumpulan Phen, Chry dan Phen+Chry.

Tikus-tikus yang disuntik dengan partikel Phen dan Chry menunjukkan aras IgG dan IgA dan aktiviti alveolar makrofaj yang paling rendah pada paru-paru. Walau bagaimanapun, tikus yang disuntik dengan partikel di atas menunjukkan peningkatan pada glutathione-S-transferase (GST) dan aktiviti protease. Kajian ini juga membuktikan bahawa tikus yang menjadikan bawang putih sebagai diet harian, dapat menghalang gangguan pada pertumbuhan sel dan meningkatkan lagi mekanisme pertahanan pulmonari.

Akhir sekali, pendedahan akut dan kronik terhadap bahan merbahaya dari persekitaran dapat memberi kesan kepada paru. Bawang putih bagaimanapun, telah menunjukkan potensi dapat mengurangkan kesan gangguan kronik Phen, Chry dan Phen+Chry pada tikus-tikus.



ACKNOWLEDGEMENTS

First of all, I would like to convey my deepest appreciation and thanks to Associate Professor Dr. Noordin Mohamed Mustapha, the chairman of my supervisory committee for his endless advise, guidance, patience and encouragement that lead to the completing of this thesis.

I would like to express my thanks and gratitude to Professor Dato' Sheikh Omar Abdul Rahman, a member of my supervisory committee for his study and valuable time throughout this study. I also would like to express my thanks to the other members of my supervisory committee, Dr. Hassan Hj. Mohd. Daud for his opinion, help and permitting me to use his laboratory on my molecular work.

My deeply appreciation and special thanks goes to Dr. Abdul Rahman Omar, who has provided facilities in his laboratory throughout this project. Thanks also go for his kind assistance and invaluable technical guidance. Also thanks go to Dr. Rasedee Abdullah.

My sincere thanks go to UPM for the PASCA foundation that help me got through this study comfortably. A special thanks and appreciation goes to Cik Kamaruddin from the Virology Lab, Cik Hajariah from the Bacteriology Lab and the staff from the Post Mortem Lab.



Here I would like to express my very grateful to an incredible friend Teh Rasyidah Ismail; she is my friend, partner and a family to me as I go through this study. Her encouragement and support really boost my spirit to go on and finish this study. My thanks also go to Imilia, Kak Ina, Kak Zhie (my dear housemate for being around) and my special friend Lil.

Mak, bapak, mamanah, Linda, Ghoi, mamanoin, pak E, Serl, Battah, BEdah, Ainul and Aisyah for being a good family to me all this while and always supporting me regardless every decision I made.

Last but not least, my thanks also due to many others who have one or another aided me on the field as well as in the laboratory for the success of this study. Also special appreciation goes to me for hanging around with the patient build inside throughout the years and this study specifically.



I certify that an Examination Committee met on 13th November 2003 to conduct the final examination of Wan Mazlina Bt. Md. Saad on her Master of Science thesis entitled "The Efficacy of Garlic in Alleviating Phenanthrene and Chrysene Induced Pulmonary Injury in Rats" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Jasni Sabri, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

Noordin Mohamed Mustafa, Ph.D.


Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Dato' Sheikh Omar Abdul Rahman, Ph.D.

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Hassan Hj. Mohd. Daud, Ph.D.

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 27 FEB 2004



The thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment for the degree of Master of Science. The members of the Supervisory Committee are as follow:

NOORDIN MOHAMED MUSTAPHA, Ph.D.

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

DATO' SHEIKH OMAR ABDUL RAHMAN, Ph.D.

Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

HASSAN HJ. MOHD. DAUD, Ph.D.

Lecturer
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)




AINI EDERIS, Ph.D.

Professor/ Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **15 APR 2004**



I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



WAN MAZLINA BT. MD. SAAD

Date: 15 APRIL 2004

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENT	ix
APPROVAL	xii
DECLARATION	xiii
LIST OF TABLES	xvi
LIST OF PLATES	xvii
LIST OF ABBREVIATIONS	xviii
 CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	
Air Pollution	5
Haze	7
Polycyclic Aromatic Hydrocarbon (PAH)	8
The Effects of PAH	9
Phenanthrene	
Physicochemical properties	11
Metabolism and Deposition	12
Distribution	13
Metabolism	13
Excretion	13
Other Routes of Exposure	13
Chrysene	
Physicochemical properties	14
Metabolism and disposition	15
Distribution	16
Metabolism	16
Excretion	17
Target Organs/Critical Effects	18
Carcinogenicity	18
Other Routes of Exposure	19
EPA-Weight of Evidence	20
Garlic	20



III	ACUTE EXPOSURE OF PHENANTHRENE AND CHRYSENE	
	Introduction	23
	Materials and Methods	
	Animals and Management	24
	Inoculum	24
	Experimental Design	25
	Pathology	26
	Detection of DNA Laddering	27
	Protease Activity	28
	Statistical Analysis	29
	Result	29
	Discussion	35
IV	CHRONIC EXPOSURE OF PHENANTHRENE AND CHRYSENE	
	Introduction	42
	Materials and Methods	
	Animals	44
	Inoculum	44
	Garlic	44
	Experimental Design	45
	Immunology	45
	Assay of Alveolar Macrophage Activity	47
	Glutathione-S-Transferase (GST)	48
	Protease Activity	49
	Result	49
	Discussion	62
V	GENERAL DISCUSSION AND CONCLUSION	71
	REFERENCES	75
	APPENDICES	86
	BIODATA OF THE AUTHOR	93



LIST OF TABLES

Table	Page
3.1: The experimental design of acute exposure to Phen and Chry in rats.....	25
3.2: The percentage of apoptosis in lung of rats instilled with Phen, Chry and Phen+Chry at different intervals (Mean \pm SD).....	31
3.3: The percentage of macrophage in lung of rats instilled with Phen, Chry and Phen+Chry at different intervals (Mean \pm SD).....	31
3.4: The percentage of neutrophil in lung of rats instilled with Phen, Chry and Phen+Chry at different intervals (Mean \pm SD).....	32
3.5: The levels (OD) of Protease activity in the lung lavage of rats during acute exposure.....	36
4.1: The experimental design of the chronic exposure study.....	46
4.2: The percentage of lesion scores of lung histology of rats at necropsy (% (Mean \pm SD).....	55
4.3: The levels of IgA and IgG (OD) in the lung lavage of rats at necropsy (OD; Mean \pm SD).....	57
4.4: The levels of IgG in the serum of rats at necropsy (OD; Mean \pm SD).....	58
4.5: The percentage of alveolar macrophage (AM \emptyset) activities in the lung lavage of rats at necropsy (Mean \pm SD).....	59
4.6: The glutathione-S-transferase (GST) enzyme activity in the lung of rats at necropsy (I.U./mg of protein; Mean \pm SD).....	60
4.7: The levels (OD) of Protease activity in the lung lavage of rats during chronic exposure (Mean \pm SD).....	61



LIST OF PLATES

Plates	Page
3.1: Photomicrograph, lung, rat, Chry group at 8 hr p.i. Margination of nuclear chromatin is observed in cells underlying apoptosis (arrow) (x 400, H&E).....	32
3.2: Photomicrograph, lung, rat, Phen + Chry group at 24 hr p.i. Note the presence of pneumocytes undergoing apoptosis (arrows) (x 400, H&E).....	33
3.3: Photomicrograph, lung, rat, Phen group at 24 hr p.i. Note the presence of neutrophil (arrow) and macrophage (dotted arrows)(x 600, H&E).....	33
3.4: Photomicrograph, lung, rat from Phen+Chry group, at necropsy following 24 hrs p.i. [Flourescein stain (H&E, X500)]......	34
3.5: Photograph, agarose gel, all groups.....	35
4.1: Photograph, lung, rat, Phen+Chry group necropsied at 90 days p.i. Note the haemorrhagic spots on the lung surface (dotted circles).....	51
4.2: Photograph, lung, rat from Phen+Chry group necropsied at 90 days p.i. Note the presence of a whitish nodule (arrow) measuring approximately 5mm in diameter on the right side.....	51
4.3: Photomicrograph, lung, rats from Phen+Chry group chronic. Hyperplasia (arrow) of the alveolar epithelium of the lung (x 625, H&E)	53
4.4: Photomicrograph, lung, rat, Phen+Chry group chronic. Necrosis (arrows) of the pneumocyte (x400, H&E).....	53
4.5: Photomicrograph, lung, rat from Phen+Chry group chronic. Neutrophil (arrows) and macrophage (dotted arrows) in the alveolar epithelium of the lung (x 200, H&E).....	54
4.6: The photomicrograph of an AMØ (arrow) in the act of phagocytosis and killing of bacteria. Acridine orange and Crystal violet, X100.....	58



LIST OF ABBREVIATIONS

AMØ	alveolar macrophage
AMT	allylmethyl trisulfide
AO	acridine orange
NH ₄ CL	ammonium chloride
bp	base pair
BAL	bronchoalveolar lavage
B(a)P	benzo(a)pyrene
C	control
CAP	concentrated ambient particles
Chry	chrysene
CO ₂	carbon dioxide
COH	coefficient of haze
CV	crystal violet
DAS	diallyl sulfide
DCNB	1, 2-dichloro-4-nitrobenzene
DNA	deoxyribonucleic acid
EDTA	ethylenediaminetetraacetic acid
ELISA	enzyme-linked immunosorbent assay
EOM	extractable organic material
EPA	Environmental Protection Agency
G	garlic



GST	glutathione-s-transferase
H ₂ O ₂	hydrogen peroxide
H&E	haematoxylin and eosin
IgA	immunoglobulin A
IARC	International Atmosphere Research Centre
IgG	immunoglobulin G
IL-1	interleukin -1
IL-2	interleukin-2
i.p.	intraperitoneal
LAK cell	leukocytes activated K cells
NaCL	sodium chloride
NAS	National Academy of Sciences
N-SLAPN	N-methoxyl-succinyl-ala-ala-pro-val-p-nitroanilide
OD	optical density
PAH	polycyclic aromatic hydrocarbon
PBS	phosphate buffered saline
Phen	phenanthrene
p.i.	post-instillation
SD	standard deviation
SLAPN	succinyl,alanine,alanine-p-nitroanilide
SSC	standard saline citrate
TUNEL	terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling analysis

Th-1	t helper 1
TNF	tumor necrosis factor
TNF- α	tumor necrosis factor-alpha
UPM	Universiti Putra Malaysia
USA	United States of America
UV	ultra violet
WR	working reagent



CHAPTER I

INTRODUCTION

Air pollution is described as impurities in the air due to various substances that endangers the environment or to health of living things. It usually occur due to massive build up of automobile exhaust, industrial and mine smoke. This pollutant may enter the human blood stream through the nose, mouth, skin and digestive tracts which can cause irritation to the eye, throat and lungs respectively. Epidemiology studies indicate that prolonged exposure to high pollution levels is associated with increased risk of cancer especially that of the lung.

The 1997 Indonesian forest fire was an environment disaster with exceptional proportions. Such a disaster may cause massive transboundary air pollution and discriminate destruction of biodiversity in the world. The immediate consequence of these fires was the production of large amounts of haze in the region, causing visibility and health problems within Southeast Asia including Malaysia.

In Kuala Lumpur, a two to three fold increased in the number of respiratory diseases were recorded during the haze (Brauer, 1997; Benedict, 1998). The haze consists mainly of fine particulate matter in the respirable range with predominantly, carcinogenic polycyclic aromatics hydrocarbon (PAHs).

The PAHs are ubiquitous environmental pollutants and include some of the most carcinogenic materials (Anon, 1983). These compounds are usually present in the environment such as components of complex mixtures in tobacco smoke, polluted air, gasoline and diesel engine exhaust and industrial waste from various manufacturing and chemical processes. The PAHs are the most prominent among the genotoxic and carcinogenic agents present in polluted urban air and have been the focus of attention of most air pollution-related biomarker field studies.

Many of these compounds are potent carcinogen (Santodonato *et al.*, 1981) which must be activated metabolically before gaining the ability to attack DNA (Jerina *et al.*, 1981). The PAHs exposed experimental animals shows that the level of PAH-DNA adducts correspond with the tumorigenic response in the lung.

Phenanthrene (Phen) and chrysene (Chry) are among the PAH found to exceed the normal level during the 1997 haze episode. Phenanthrene is ubiquitous in the environment as a product of incomplete combustion of biomass and has been identified in ambient air, surface and drinking water and in foods (Anon, 1983; Anon, 1988). In mammals; this PAH is rapidly metabolized by the liver and excreted primarily in the urine. It is also known to be a human skin photosensitiser and mild allergen (Sandmeyer, 1981).

Phenanthrene is absorbed following an oral and dermal exposure (Storer *et al.*, 1984) and would be absorbed from the lungs (Anon, 1987). Current theories regarding

the mechanism of metabolic activation of PAHs predicts the carcinogenic potential of Phen.

Chrysene accumulates as a result of anthropogenic activities such as coal combustion and gasification, gasoline exhaust, diesel and air craft exhaust and emission from coke ovens, wood burning stoves and waste incineration (Anon, 1983, Anon, 1990). Human are exposed to Chry by oral, inhalation and dermal routes. Significant exposure to Chry also occurs through the inhalation of mainstream cigarette smoke (Anon, 1983).

Chrysene is preferentially distributed to highly lipophilic regions of the body, most notably adipose and mammary tissue (Bartosek *et al.*, 1984). Information on the absorption of Chry in humans is not documented. However, the detection of PAHs, including Chry and its metabolites in the urine of individual who smoke, work in polluted industrial environment with Chry or used therapeutic coal-tar creams (Clonfero *et al.*, 1986) provides indirect evidence of inhalation and dermal absorption. Mixture toxicity did appear to increase slightly when Chry, benzo (a) anthracene and anthracene were added to Phen, pyrene and fluoranthrene (Boxall and Maltby, 1997).

Thus, it is the aim of this study to assess the following acute and chronic effects of phenanthrene and chrysene with the following objectives, to determine:

- i. the manner of cell death during acute exposure to Phen and Chry
- ii. the development and type of chronic changes induced by Phen and Chry
- iii. the sensitive and reliable indicator of chronic changes induced by Phen and Chry
- iv. the effects of Phen and Chry on proteases and protease inhibitors of rat lung
- v. the efficacy of garlic in alleviating Phen and Chry induced injury