

The Validity And Reliability Of The Turkish Version Of The Online Counseling And Face-To-Face Counseling Attitudes Scale

İbrahim Demirci, Ali Haydar Şar¹, Abdullah Manap²

¹Sakarya University, Turkey, ²İnönü University, Turkey

ARTICLE INFO

Article History:

Received 19.06.2014

Received in revised form

01.09.2014

Accepted 04.09.2014

Available online

26.09.2014

ABSTRACT

The purpose of this study is to examine the validity and reliability of the Turkish version of the Online Counseling Attitudes Scale (OCAS) and Face-to-Face Counseling Attitudes Scale (FCAS) (Rochlen, Beretvas, & Zack, 2004). OCAS and FCAS loaded on four factors and the factors were named Value of Online Counseling (OC-V), Discomfort with Online Counseling (OC-D), Value of Face-to-Face Counseling (FC-V) and Discomfort with Face-to-Face Counseling (FC-D). The results of confirmatory factor analysis indicated that the four dimensional model was well fit ($\chi^2=323.61$, $df=157$, $p=0.00000$, $RMSEA=.056$, $NFI=.94$, $CFI=.96$, $IFI=.96$, $RFI=.92$, $GFI=.91$, and $SRMR=.071$). Factor loadings ranged from .47 to .88. The internal consistency coefficients of four subscales were .84, .80, .88 and .78. Overall, findings demonstrated that OCAS and FCAS may be used as a valid and reliable instrument in order to assess attitudes toward online counseling and face to face counseling of individuals.

© 2014 IJPES. All rights reserved

Keywords:

Online counselling, attitudes, validity, reliability, scale adaptation

1. Introduction

Online Counseling is defined as any delivery of mental and behavioral health services, including but not limited to therapy, consultation and psycho-education, by a licensed practitioner to a client in a non-face-to-face setting through distance communication devices for instance, the telephone, fax, asynchronous e-mail, synchronous chat, and videoconferencing (Mallen, & Vogel, 2005; Rochlen, Zach, & Speyer, 2004; Rochlen, Beretvas, & Zack, 2004). Online counseling is different from other types of computer-related services in that it includes direct communication between a counselor and at least one client and the communication is specifically tried to address mental health concerns, analogous to a "talk therapy" procedure that might be conducted face-to-face in a mental health professional's office (Rochlen, Beretvas, & Zack, 2004). We explain simply online counseling in which you would interact with a counselor via the Internet and face-to-face counseling where you would go to a counselor's office.

Since the advent of online counseling, there has been considerable debate over the effectiveness of online counseling versus traditional face-to-face counseling. Some professionals try to argue online counseling out of existence; others seem to have limitless enthusiasm for this latest development in the interface of counseling and technology (Shaw & Shaw, 2006). Disadvantages of online counseling involve maintaining confidentiality over the Internet, delivery of online counseling services by unqualified practitioners as well as important concerns regarding confidentiality, in emergency situations, the inability to directly intervene in a crisis the lack of nonverbal information such as facial expression, tone of voice, and body language, the dangers of offering online services over state jurisdiction lines, the lack of efficacy studies of online counseling services, the difficulty of developing a therapeutic relationship with a client who is never seen face-to-face and the lack

¹ Corresponding author's address: Sakarya University, Faculty of Education, Department of Educational Sciences, 54300 Sakarya, Turkey

Telephone: +90 264 2957145

Fax: +90 264 2957183

e-mail: asar@sakarya.edu.tr (A. H. Şar)

<http://dx.doi.org/10.17220/ijpes.2014.01.002>

of therapeutic control, the lack of visual cues (Barak, 1999; Murphy & Mitchell, 1998; Sampson, Kolodinsky, & Greeno, 1997; Rochlen, Beretvas, & Zack, 2004; Shaw & Shaw, 2006).

As for the Advantages of online counseling, advocates for online counseling emphasize such benefits; potential utility in reaching populations that cannot or will not traditional mental health services the convenience of the service, accessibility of services to clients who are in geographically isolated areas, are physically disabled or seriously ill and cannot leave home, would ordinarily not seek traditional counseling, would feel more comfortable expressing themselves in a written format, the ease of record keeping, and the possible decreased costs related with providing these services (Barak, 1999; Murphy & Mitchell, 1998; Rochlen, Beretvas, & Zack, 2004; Shaw & Shaw, 2006).

An attitude may be explained as a positive or negative evaluation of people, objects, event, activities, ideas, or just about anything in environment this is generally referred to as the attitude object. Social psychologists research attitudes in terms of three parts: cognitive, affective, and behavioral. Attitudes are shaped by our experiences or observing experiences, serve to guide our future conducts and can be used to predict behavior (Kağıtçıbaşı, 1999; Köklü, 1995).

Online counseling services and online mental health services are being accessed in a variety of formats and are expected to developed in the future (Norcross, Hedges, & Prochaska, 2002) and online counseling is a field that has seen increase for ten years (Barak, Hen, Boniel-Nissim, & Shapira 2008). Researchers have tried to reveal clients' attitudes toward seeking help in relation to various demographic, psychological, and cultural variables. One of the most glaring reasons for the omission of this information is a lack of research instruments designed to measure the public's perceptions of online counseling services. This scale exhibits attitudes toward online counseling and a comparable measure of attitudes toward face-to-face counseling services. The public's perceptions of online counseling may help to display or explain clinician's perspectives on the need and demand for such services (Rochlen, Beretvas, & Zack, 2004).

Internet-based treatment programs frequently integrate cognitive-behavior therapy principles, are structured, and have an evidence base. Internet-based treatment programs usually also contain rigorous assessment process prior to commencement of treatment to ensure that the consumer has the clinical disorder that the program treats. Online counseling with online cognitive behavioral therapy deal with; anxiety and depression (Topolovec-Vranic et al., 2010; Graaf et al., 2009; Sethi & Campbell, 2010; Newman, Consoli, & Taylor, 2006; Warmerdam, Stratena, & Jongasma, 2010; Graaf et al., 2011; Beattie, Shaw, Kaur, & Kessler, D2009), obsessive-compulsive disorders (Moritz, Wittekind, Hauschildt, & Timpano, 2011; Wootton et al., 2011), alcohol problem (Postel, De Haan, & De Jong, 2010; Blankers, Koeter, & Schippers, 2011), sexual disorders (McKee, 2004), suicidal ideation (Greidanus & Everall, 2010), eating disorder (Fichter et al, 2012), Panic disorder (Pier et al, 2008; Austin & Kiropoulos, 2007; Advocat & Lindsay, 2009), Posttraumatic stress disorder (Kleina et al., 2010), Social phobia (Berger et al., 2011), Sexual abuse (Littleton et al., 2011) such as various mental problems outcomes.

The Online and Face-to-Face Counseling Attitudes Scale (Rochlen, Beretvas, & Zack, 2004): The result of exploratory factor analyze of OCAS consist of two factors, and factors explained 64.6% of the variance in the responses to the 10 items and an interfactor correlation of-.52. The two FCAS factors together explained 62.2% of the variance in the responses to the 10 items and a correlation between the components of-.59. The factors were named Value of Online Counseling (OC-V) and Discomfort with Online Counseling (OC-D). The OC-V subscale represented the general feelings of participants toward the utility of online counseling, including their perceptions of how they and others would benefit from online counseling. The OC-D subscale represented participants' emotional feelings and reactions regarding their ease and comfort associated with using online counseling services. Example of items assessing each of these two factors include "Using an online counseling would help me learn about myself" and "I would dread explaining my problems to an online counselor" for the OC-V and OC-D subscales, respectively. The same base set of 10 items was selected to assess each of the FCAS factors. These factors were similarly named Value of Face-to-Face Counseling (FC-V) and Discomfort with Face-to-Face Counseling (FC-D). Again, each "Using a face-to-face counseling would help me learn about myself" and "I would dread explaining my problems to a face-to-face counselor". And the result of confirmatory factor analysis (CFA) $\chi^2=265,49$, $p<0.05$, RMSEA=.057, CFI=.96, TLI=95, and SRMR=.57. The pattern coefficients were strong and significant ($p<.001$). Factor loadings ranged from.520 to .921. The Result

of multiple Group CFA $\chi^2=601,52$, $p<0.05$, RMSEA=.041, CFI=.95, TLI=.95, GFI=.96, and SRMR=.77. Internal consistency estimates ranged from .77 to .90 over multiple studies for scores on both the Online Counseling and Face to Face Counseling subscales, and test–retest correlations of .70 to .88 were observed over a 3-week period on both subscales. Responses are made on a 6-point Likert scale (1-strongly disagree, 6-strongly agree).

2. Method

2.1. Participants

Participants were 345 (173 were female and 172 were male) university students.

2.2. Procedure

Primarily the Online Counseling and Face-to-Face Counseling Attitudes Scale were translated into Turkish by six academicians. After that the Turkish form was back-translated into English and examined the consistency between the Turkish and English forms. Turkish form has reviewed by five academicians from educational sciences department. Finally they discussed the Turkish form and along with some corrections this scale was prepared for validity and reliability analyses. In this study confirmatory factor analysis was executed to confirm the original scale's structure in Turkish culture. As reliability analysis internal consistency coefficients and the item-total correlations were examined. Data were analyzed by LISREL 8.8 and SPSS 20.0.

3. Results

3.1. Construct Validity

The results of confirmatory factor analysis indicated that the four dimensional model was well fit and Chi-Square value ($\chi^2=323.61$, $df=157$, $p=0.00000$) which was calculated for the adaptation of the model was found to be significant. The goodness of fit index values of the model were RMSEA=.056, NFI=.94, CFI=.96, IFI=.96, RFI=.92, GFI=.91, and SRMR=.071. Factor loadings ranged from .47 to .88. The correlation between the OC-V and OC-D subscales of the OCAS was $-.40$ ($p<.001$), and the correlation between the FC-V and FC-D subscales was $-.28$ ($p<.001$) for the FCAS. The correlation between the OC-V and FC-V subscales was $.37$, $p<.001$). The correlation between the OC-D and FC-D subscales was $.49$, $p<.001$).




Figure 1: OCAS and FCAS Path Diagram and Factor Loading

3.2. Reliability

The internal consistency coefficients of OCAS and FCAS four subscales were OC-V, .84, OC-D, .80, FC-V, .88 and FC-D, .78. The corrected item-total correlations of OC-V ranged from .59 to .70, OC-D ranged from .50 to .71, FC-V ranged from .71 to .73 and FC-D ranged from .43 to .68.

Table 1: OCAS Item-Total Correlation

OC-V		OC-D	
Mad. No	R	Mad. No	r
1	.69	6	.50
2	.62	7	.62
3	.70	8	.71
4	.63	9	.49
5	.59	10	.61

Table 2: FCAS Item-Total Correlation

FC-V		FC-D	
Mad. No	r	Mad. No	r
1	.73	6	.43
2	.73	7	.61
3	.74	8	.68
4	.71	9	.43
5	.62	10	.64

4. Discussion and Conclusion

The aim of this study was to translate the OCAS and FCAS into Turkish and examine its psychometric properties. Numerous fit indexes are used for the aim of presenting the adequacy of the model tested in Confirmatory factor analysis (CFA). For GFI, CFI, NFI, RFI and IFI indexes, acceptable fit value is considered to be 0.90 and best fit value is 0.95 (Bentler, 1980; Bentler & Bonett, 1980; Marsh, Hau, Artelt, Baumert, & Peschar, 2006). As for the RMSEA, acceptable fit value is considered to be 0.08 and best fit value is 0.05 (Brown & Cudeck, 1993; Byrne & Campbell, 1999). CFA was conducted so as to indicate whether the factor structure of the original form would be confirmed in the Turkish sample. The OCAS and FCAS loaded on four factors. The results of CFA indicated that the model was well fit. Thus, it can be said that the structural model of the OCAS and FCAS which consists of four factors was well fit to the Turkish culture. The internal consistency coefficients of four subscales were high. Overall, findings demonstrated that OCAS and FCAS had high validity and reliability scores (Büyüköztürk, 2004; Tabachnick & Fidell, 1996; Tezbaşaran, 1996) and OCAS and FCAS may be used as a valid and reliable instrument in order to assess attitudes toward online counseling and face to face counseling of individuals and accurate comparisons between the two types of counseling services. Nevertheless, further studies that will use the OCAS and FCAS are important for its measurement force.

References

- Austin, D., & Kiropoulos, L. (2007). An Internet-based investigation of the catastrophic misinterpretation model of panic disorder. *Journal of Anxiety Disorders, 22*(2), 233–242. DOI: 10.1016/j.janxdis.2007.01.013.
- Barak, A. (1999). Psychological applications on the internet: A discipline on the threshold of a new millennium. *Applied and Preventive Psychology, 8*, 231-246. DOI: 10.1016/S0962-1849(05)80038-1.
- Barak, A., Hen, L., Boniel-Nissim, M., & Shapira, N. (2008). A comprehensive review and a meta-analysis of the effectiveness of internet-based psychotherapeutic interventions. *Journal of Technology in Human Services, 26*, 109–60. DOI: 10.1080/15228830802094429.
- Beattie, A., Shaw, A., Kaur, S., & Kessler, D. (2009). Primary-care patients expectations and experiences of online cognitive behavioral therapy for depression: a qualitative study. *Blackwell Publishing Ltd Health Expectations, 12*, 45–59. DOI: 10.1111/j.1369-7625.2008.00531.x.
- Bentler, P. M. (1980). Multivariate analysis with latent variables: Causal modeling. *Annual Review of Psychology, 31*, 419-456. DOI: 10.1146/annurev.ps.31.020180.002223.

- Bentler, P. M., & Bonett, D. G. (1980). Significance tests and goodness of fit in the analysis of covariance structures. *Psychological Bulletin*, 88, 588-606. DOI: 10.1037/0033-2909.88.3.588.
- Berger, T., Caspar, F., Richardson, R., Kneubühler, B., Sutter, D., & Andersson, G. (2011). Internet-based treatment of social phobia: A randomized controlled trial comparing unguided with two types of guided self-help. *Behaviour Research and Therapy*, 49(3), 158-169. DOI: 10.1016/j.brat.2010.12.007.
- Brown, M., & Cudeck, R. (1993). Alternative ways of assessing model fit. In: K. Bollen & J. Long, (Eds), *Testing structural equation models* (pp. 136–162). London: Sage Publications. DOI: 10.1177/0049124192021002005.
- Byrne, B., & Campbell, T. L. (1999). Cross-cultural comparisons and the presumption of equivalent measurement and theoretical structure: A look beneath the surface. *Journal of Cross-Cultural Psychology*, 30(5), 555-574. DOI: 10.1177/0022022199030005001.
- Büyüköztürk, Ş. (2004). *Veri analizi el kitabı*. Ankara: Pegem A Yayıncılık.
- Fichter, M. M., Quadflieg, N., Nisslmüller, K., Lindner, S., Osen, ...Wünsch-Leiteritz, W. (2012). Does internet-based prevention reduce the risk of relapse for anorexia nervosa? *Behaviour Research and Therapy*, 3(11), 103-115. DOI: 10.1016/j.brat.2011.12.003.
- Graaf, L. E., Gerhards, S. A., Arntz, A., Riper, H., Metsemakers, J. F., ...Huibers M. J. (2009). Clinical effectiveness of online computerized cognitive-behavioral therapy without support for depression in primary care: randomized trial. *The British Journal of Psychiatry*, 195, 73–80. DOI: 10.1192/bjp.bp.108.054429.
- Graaf, L. E., Gerhards, S. A., Arntz, A., Riper, H., Metsemakers, J. F., ...Huibers M. J. (2011). One-year follow-up results of unsupported online computerized cognitive behavioral therapy for depression in primary care: A randomized trial. *Journal of Behavior Therapy and Experimental Psychiatry*, 42(1), 89-95. DOI: 10.1016/j.jbtep.2010.07.003.
- Greidanus, E., & Everall, R. D. (2010). Helper therapy in an online suicide prevention community. *British Journal of Guidance & Counseling*, 38(2), 191-204. DOI: 10.1080/03069881003600991.
- Kağıtçıbaşı, Ç. (1999). *Yeni İnsan ve İnsanlar*. İstanbul; Evrim Yayınevi.
- Klein, B., Richards, J. C. & Austin, D. W. (2006). Efficacy of Internet therapy for panic disorder. *Journal of Behavior Therapy and Experimental Psychiatry*, 37(3), 213-238.
- Köklü, N. (1995). Tutumların ölçülmesi ve likert tipi ölçeklerde kullanılan alternatif seçenekler. *Eğitim Bilimleri Fakültesi Dergisi*, 28(2), 81-9.
- Littletona, H., Buck, K., Rosman, L., & Grills-Taquechel, A. (2011). From Survivor to The river: A Pilot Study of an Online Program for Rape Victims. *Cognitive and Behavioral Practice*, 3(4), 147-159. DOI: 10.1016/j.cbpra.2011.04.002.
- Mallen, M. J., & Vogel, D. L. (2005). Introduction to the Major Contribution: Counseling psychology and online counseling. *The Counseling Psychologist*, 33, 761–775. DOI: 10.1177/0011000005278623.

- Marsh, H. W., Hau, K. T., Artelt, C., Baumert, J., & Peschar, J. L. (2006). OECD's brief self-report measure of educational psychology's most useful affective constructs: Cross-cultural, psychometric comparisons across 25 countries. *International Journal of Testing*, 6(4), 311-360. DOI: 10.1207/s15327574ijt0604_1.
- Blankers, M., Koeter, M. W., & Schippers, G. M. (2011). Internet therapy versus internet self-help versus no treatment for problematic alcohol use: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 79 (3), 330-341. DOI: 10.1037/a0023498.
- McKee, H. A. (2004). Always a shadow of hope: Heteronormative binaries in an online discussion of sexuality and sexual orientation. *Computers and Composition*, 21, 315-340.
- Moritz, S., Wittekind, C. E., Hauschildt, M., & Timpano, K. R. (2011). Do it yourself ? Self-help and online therapy for people with obsessive-compulsive disorder. *Current Opinion in Psychiatry*, 24, 541-548. DOI: 10.1097/YCO.0b013e32834abb7f.
- Murphy, L. J., & Mitchell, D. L. (1998). When writing helps to heal: E-mail as therapy. *British Journal of Guidance and Counseling*, 26(2), 1-32. DOI: 10.1080/03069889808253835.
- Newman, M. G., Consoli A., & Taylor, C. B. (1997). Computers in assessment and cognitive behavioral treatment of clinical disorders: Anxiety as a case in point. *Behavior Therapy*, 28, (2), 211-235. DOI: 10.1016/S0005-7894(97)80044-5.
- Norcross, J. C., Hedges, M., & Prochaska, J. O. (2002). The face of 2010: A delphi poll on the future of psychotherapy. *Professional Psychology: Research and Practice*, 3, 316-322. DOI:10.1037/0735-7028.33.3.316.
- Pier, C., Austin, D. W., Klein, B., Mitchell, J., Schattner, P., ...Wade, V. (2008). A controlled trial of internet-based cognitive-behavioral therapy for panic disorder with face-to-face support from a general practitioner or email support from a psychologist. *Mental Health in Family Medicine*, 5, 29-39.
- Postel, M.G., De Haan, H. A., & De Jong, C.A. J. (2010). Evaluation of an E-Therapy Program for Problem Drinkers: A Pilot Study. *Substance Use & Misuse*, 45, 2059-2075. DOI: 10.3109/10826084.2010.481701.
- Rochlen, A.B., Beretvas, S.N., & Zack, S.Z. (2004). The Online and Face-to-Face Counseling Attitudes Scales: A Validation Study. *Measurement and Evaluation in Counseling and Development Journal*, 37(1), 95-111.
- Tabachnick, B. G., & Fidell, L. S. (1996). *Using multivariate statistics*. New York: HarperCollins College Publishers.
- Tezbaşaran, A. A. (1996). *Likert tipi ölçek geliştirme kılavuzu*. Ankara: Türk Psikologlar Derneği Yayınları.
- Topolovec-Vranic, J., Cullen, N., Michalak, A., Ouchterlony, D., Bhalerao, S., ...Cusimano, M. D. (2010). Evaluation of an online cognitive behavioral therapy program by patients with traumatic brain injury and depression. *Brain Injury*, 24(5), 762-772. DOI: 10.3109/02699051003709599.
- Sampson, J. E, Jr., Kolodinsky, R. W., & Greeno, B. E (1997). Counseling on the information highway: Future possibilities and potential problems. *Journal of Counseling and Development*, 75, 203-212. DOI: 10.1002/j.1556-6676.1997.tb02334.x.

- Shaw, H.E., & Shaw, S.F., 2006. Critical Ethical Issues in Online Counseling: Assessing Current Practices With on Ethical Intent Checklist. *Journal of Counseling and Development*, 84, 41-53. DOI: 10.1002/j.1556-6678.2006.tb00378.x.
- Sethi, S., & Campbell, A. J. (2010). The use of computerized self-help packages to treat adolescent depression and anxiety. *Journal of Technology in Human Services*, 28, 144-160. DOI: 10.1080/15228835.2010.508317.
- Warmerdama, L., Stratena, A., & Jongsmas, J. (2010). Online cognitive behavioral therapy and problem-solving therapy for depressive symptoms: Exploring mechanisms of change. *Journal of Behavior Therapy and Experimental Psychiatry*, 41(1), 64–70. DOI: 10.1016/j.jbtep.2009.10.003.