

ZEBRAFISH
Volume 13, Number 5, 2016
© Mary Ann Liebert, Inc.
DOI: 10.1089/zeb.2016.1280

High-Resolution Magic Angle Spinning Nuclear Magnetic Resonance of Intact Zebrafish Embryos Detects Metabolic Changes Following Exposure to Teratogenic Polymethoxyalkenes from Algae

John P. Berry,¹ Upasana Roy,^{2,3} Asha Jaja-Chimedza,¹ Kristel Sanchez,¹ Joerg Matysik,³ and A. Alia^{2,4}

Abstract

Techniques based on nuclear magnetic resonance (NMR) for imaging and chemical analyses of *in vivo*, or otherwise intact, biological systems are rapidly emerging and finding diverse applications within a wide range of fields. Very recently, several NMR-based techniques have been developed for the zebrafish as a model animal system. In the current study, the novel application of high-resolution magic angle spinning (HR-MAS) NMR is presented as a means of metabolic profiling of intact zebrafish embryos. Toward investigating the utility of HR-MAS NMR as a toxicological tool, these studies specifically examined metabolic changes of embryos exposed to polymethoxy-1-alkenes (PMAs)—a recently identified family of teratogenic compounds from freshwater algae—as emerging environmental contaminants. One-dimensional and two-dimensional HR-MAS NMR analyses were able to effectively identify and quantify diverse metabolites in early-stage (≤ 36 h postfertilization) embryos. Subsequent comparison of the metabolic profiles between PMA-exposed and control embryos identified several statistically significant metabolic changes associated with subacute exposure to the teratogen, including (1) elevated inositol as a recognized component of signaling pathways involved in embryo development; (2) increases in several metabolites, including inositol, phosphoryl choline, fatty acids, and cholesterol, which are associated with lipid composition of cell membranes; (3) concomitant increase in glucose and decrease in lactate; and (4) decreases in several biochemically related metabolites associated with central nervous system development and function, including γ -aminobutyric acid, glycine, glutamate, and glutamine. A potentially unifying model/hypothesis of PMA teratogenicity based on the data is presented. These findings, taken together, demonstrate that HR-MAS NMR is a promising tool for metabolic profiling in the zebrafish embryo, including toxicological applications.

Introduction

METHODS BASED ON THE APPLICATION of nuclear magnetic resonance (NMR) for imaging (i.e., magnetic resonance imaging [MRI]) and chemical characterization (i.e., NMR spectroscopy) of *in vivo*, or otherwise intact, biological systems have rapidly evolved within biomedical and related scientific fields. With respect to imaging, MRI techniques utilize, in particular, the abundance of water in biological systems, coupled with the detection of the magnetic nuclear spin of hydrogen (^1H) atoms in H_2O , to collect high-resolution spatial data within biological samples and subse-

quently generate resolved, three-dimensional (3D) images. Conversely, NMR spectroscopy is capable of discerning particular molecules based, likewise, on the spin of ^1H or, in fact, isotopes of several relevant atoms (e.g., ^{13}C , ^{15}N), and more exactly, the corresponding *chemical shift* of a particular atom within the molecule. When applied to biological systems and the detection and measurement of metabolites in particular, NMR spectroscopy can be specifically enabled by a technique called high-resolution magic angle spinning (HR-MAS). Briefly stated, spinning of a sample at the so-called magic angle (54.74°) relative to the applied magnetic field enables averaging of the various interactions (i.e., dipolar,

¹Department of Chemistry and Biochemistry, Florida International University, North Miami, Florida.

²Institute of Medical Physics and Biophysics, University of Leipzig, Leipzig, Germany.

³Institut für Analytische Chemie, University of Leipzig, Leipzig, Germany.

⁴Leiden Institute of Chemistry, Leiden, the Netherlands.

quadrupolar, anisotropic), which otherwise broaden chemical shifts, and thus enables narrowing of signal and consequently sufficient resolution of metabolites within a complex matrix. Accordingly, HR-MAS NMR has emerged as a very promising technique for chemical analysis, including metabolic profiling, in various biological systems.¹

Most recently, various NMR-based techniques for imaging and chemical analysis (including HR-MAS NMR) have been reported for the zebrafish as an increasingly important animal model.^{2–6} To date, however, NMR has been largely applied to adult stages of zebrafish, and has enabled *in vivo* imaging, and localized chemical analyses, in this system. With respect to the former, NMR-based microimaging (μ MRI) was recently demonstrated with live zebrafish and specifically applied to the detection and characterization of melanomas.^{2,3} Localized *in vivo* high-resolution spectroscopy, likewise, was achieved in the brain of adult zebrafish (with a spatial resolution of voxels as small as 3.3 μ L) and specifically utilized, as an example, for metabolic profiling of spatially defined regions of the brain including, most recently, effective lipid profiling.^{4,5} Although the past studies have largely focused on adult stages of the zebrafish, one such study applied NMR, and HR-MAS specifically, for chemical analysis of intact embryonic stages and, in particular, was able to quantify ethanol levels as part of exposure studies.⁶

In the present study, HR-MAS was investigated as a means of metabolic profiling of intact zebrafish embryos. As embryonic stages of the zebrafish represent a particularly versatile system with numerous practical advantages, including capacity for high-throughput analysis, the development of HR-MAS-based metabolic profiling would represent a potentially powerful metabolomics tool that could, indeed, be applicable to a wide range of scientific investigations. Among the potential applications of HR-MAS NMR to the zebrafish model is toxicological utility for understanding metabolic changes related to environmental toxins toward both understanding toxicity (i.e., targets, mechanism of action) and identifying potential biomarkers of exposure. Bolstering this potential, the zebrafish has, indeed, been utilized extensively as a toxicological model for a wide array of toxic environmental contaminants, including heavy metals, endocrine disruptors, and various organic pollutants.^{7–9} In particular, embryonic and subsequent larval stages have proven particularly useful for characterizing teratogenicity (i.e., developmental toxicity).^{8,9} With respect to the current study, the zebrafish embryo teratogenicity assay (ZETA) has been specifically used in the investigation of toxic, and specifically, teratogenic metabolites from marine and freshwater algae, including cyanobacteria,¹⁰ and has enabled both characterization of known algal toxins^{11–13} and identification (through screening, bioassay-guided isolation, and chemical/toxicological characterization) of otherwise unknown toxic metabolites.^{14–16}

In one such study, ZETA was used to identify a family of teratogenic secondary metabolites, namely, the polymethoxy-1-alkenes (PMAs), and subsequently demonstrate a taxonomically widespread distribution of these metabolites

among both prokaryotic cyanobacteria and eukaryotic algae (i.e., Chlorophyta or green algae).^{14,16} Accordingly, it was suggested that PMA may represent an emerging class of environmental toxins specifically associated with increasingly frequent and intense blooms of so-called harmful algae worldwide. In fact, PMAs have been previously described from a wide range of cyanobacteria for more than three decades.^{17–20} However, only recently have they been linked to biological activity and specifically observed teratogenicity in the zebrafish embryo model.¹⁴ As such, no data currently exist with respect to possible mechanisms, or biochemical, molecular, or cellular targets, of these compounds.

Elucidating the effects of toxic compounds, such as the PMAs, at the biochemical and molecular level is not only obviously fundamental to understanding the mode of action but also potentially informative with respect to the assessment of toxic potential through identification of measurable biomarkers of exposure. Accordingly, in the current study, HR-MAS NMR was applied to metabolic profiling of zebrafish exposed to PMAs, at specifically subacute concentrations and exposure times, as a means of identifying relevant metabolic changes toward generating hypothesis regarding mechanisms/targets of these compounds, as well as potential biomarkers, and as a model class of compounds demonstrating, more generally, the potential of the technique as a metabolomics tool for environmental toxicology.

Materials and Methods

Isolation of PMA

The teratogenic PMA variant, 4,6,8,10,12,14,16,18,20-nonamethoxy-1-pentacosene (Fig. 1)—as the typically most abundant congener—was isolated from cultures of the cyanobacterial species, *Cylindrospermopsis raciborskii*, as previously described.^{14,16} Briefly, freeze-dried biomass was twice extracted in chloroform and pooled extracts (following filtration) concentrated to dryness *in vacuo*. Subsequently, the PMA was isolated by silica gel (60 Å, 40–63 μ m) with a gradient of ethyl acetate in hexane (eluting with 100% ethyl acetate) and subsequent reverse-phase HPLC (Phenomenex Luna 5 μ m C18 100 Å, 250 \times 4.6 mm), using a gradient of 50%–100% acetonitrile (and water).

Exposure of zebrafish to PMA

Zebrafish were exposed to PMA by a method modified from previously published protocols.^{10,11,13–16} Briefly, 100 embryos (12 h postfertilization [hpf]) per replicate ($n=4$) were exposed in 35-mm-diameter polystyrene dishes to 20 μ g/mL of PMA (in E3 medium²¹) for 24 h. Exposure concentration was based on previously determined teratogenicity of PMA and specifically represent concentrations that are not acutely toxic (i.e., teratogenic or lethal) to minimize metabolic changes associated with mortality or severe development deformities.¹⁴ Likewise, exposure times <36 hpf (i.e., 12 hpf embryos and 24 h treatment) were selected as




FIG. 1. Structure of 4,6,8,10,12,14,16,18,20-nonamethoxy-1-pentacosene as a common and representative PMA. PMA, polymethoxy-1-alkene.

previous studies with PMA-determined teratogenicity, in the form of morphological deformities, was observed only after 72–96 hpf.¹⁴ Control embryos were exposed to the solvent carrier (i.e., methanol) alone at corresponding (v/v) concentrations. Following exposure, treated and control embryos were washed (to remove any residual PMA) and carefully transferred to a 4-mm zirconium oxide rotor (Bruker BioSpin AG), to which 10 μ L of deuterated phosphate buffer (100 mM; pH 7.0) containing 0.1% (w/v) 3-trimethylsilyl-2,2,3,3-tetradeuteropropionic acid (TSP), as a lock solvent and NMR (i.e., chemical shift) reference, was added. The rotor was immediately transferred to the NMR spectrometer.

HR-MAS NMR studies

All HR-MAS NMR experiments were carried out on a Bruker DMX 600 MHz NMR spectrometer operating with a proton resonance frequency of 600 MHz. The instrument is equipped with a 4-mm HR-MAS dual inverse $^1\text{H}/^{13}\text{C}$ probe with a magic-angle gradient. All measurements were carried out at a magic-angle spinning rate of 6 kHz and a temperature of 277 K. Temperature was controlled by a Bruker BVT3000 control unit. Bruker TOPSPIN software (Bruker BioSpin AG) was used to acquire and process the NMR data.

One-dimensional ^1H HR-MAS NMR spectra were recorded using rotor synchronized Carr–Purcell–Meiboom–Gill (CPMG) pulse sequence with water suppression. Each one-dimensional (1D) spectrum was acquired by applying a spectral width of 8000 Hz, domain data points 16k, number of averages 512 with 8 dummy scans, constant receiver gain of 2048, an acquisition time of 2 s, and a relaxation delay of 2 s. Since NMR measurements were done on intact embryos, the relaxation delay was set to a small value to remove nascent short T_2 components due to the presence of lipids. All spectra were processed by an exponential window function corresponding to a line broadening of 1 Hz and zero-filled before Fourier transformation. ^1H HR-MAS NMR spectra were phased manually and automatically baseline corrected using TOPSPIN 2.1 (Bruker BioSpin AG). The total analysis time (including sample preparation, optimization of NMR parameters, and data acquisition) of ^1H HR-MAS NMR spectroscopy for each sample was \sim 20 min.

Two-dimensional homonuclear correlation spectroscopy (^1H - ^1H COSY) was performed in magnitude mode. The parameters used for COSY were 2048 data points collected in the t_2 domain over the spectral width of 4k; 512 t_1 increments were collected with 16 transients, relaxation delay of 2 s, acquisition time of 116 ms, and presaturated water resonance during relaxation delay. The resulting data were zero-filled with 2048 data points and were weighted with the sine bell window functions in both dimensions before Fourier Transformation.

Data analysis

For quantification of metabolites, NMR data analysis was performed using MestReNova software version 6.0.3-5604 (Mestrelab Research S.L.). The concentrations of the various metabolites in the spectra were determined by comparing the integral peak intensity of the metabolite of interest with that of the TSP peak, after correcting for the number of contributing protons and for tissue weight. All statistical analyses (t -tests and ANOVAs) of the NMR quantification

results were performed with OriginPro v. 8. F -values were calculated, and F -values larger than 2.8 ($p < 0.05$) were considered significant.

Multivariate statistical analysis (i.e., principle component analysis [PCA]) of primary metabolites in the spectra was performed using Bruker software package AMIX (version 3.8.6). The CPMG spectra, collected from embryos, were subdivided in the range between 0.3 and 9 ppm into buckets of 0.04 ppm (total 218 buckets), using Bruker AMIX software (version 3.8.7; Bruker GmbH). The region of 4.80–6.00 ppm was excluded from the analysis to remove the water signal. To compensate for the differences in the overall metabolite concentration between individual samples, the data obtained were mean centered, scaled to unit variance, and then normalized by dividing each integral of the segment by the total area of the spectrum. The resulting data matrix was exported into Microsoft Office Excel (Microsoft Corporation). This was then further imported into SIMCA software (Umetrics AB) for PCA. Correlation coefficients with $p < 0.05$ were considered statistically significant.

Results

A representative 1D ^1H HR-MAS NMR spectrum obtained directly from intact, control zebrafish embryos (12 hpf) is shown in Figure 2. Highly resolved spectra were obtained, and the assignment of the metabolites was further reinstated by comparing the ^1H spectra of reference compounds to existing literature values.^{22,23} The reference compounds in the Biological Magnetic Resonance Data Bank²⁴ were specifically used for the characterization of the metabolites. For further unambiguous assignment of the metabolite resonances in zebrafish embryos, a two-dimensional (2D), homonuclear (^1H - ^1H), dipolar-correlation NMR spectrum was measured from intact embryos (Fig. 3). The 2D spectrum allowed separation of most overlapping peaks for the coupled spin systems that gave rise to off-diagonal peaks. The 2D NMR clearly reveals separate correlation networks of several metabolites and, on the basis of cross-peaks, unambiguous assignments were made (Fig. 3). The detailed description of all the peaks' assignment of metabolites is given in Table 1.

Comparison of 1D HR-MAS spectra obtained from control and PMA-treated zebrafish embryos is depicted in Figure 4. Quantitative analysis of metabolites in embryos exposed to PMA when compared to control embryos shows statistically significant changes in several identified metabolites in exposed embryos (Fig. 5). The major changes include (1) increased levels of glucose (Glc), (*myo*-)inositol (m-Ins), cholesterol (Chol), fatty acids (FA) and phosphoryl choline (PC) and (2) decreased levels of glutathione (GSH), lactose (Lac), γ -aminobutyric acid (GABA), glycine (Gly), glutamate (Glu), and glutamine (Gln). There were no observable effects (data not shown) of the PMA at the concentration and exposure time tested, with respect to mortality, developmental deformities, or other relevant endpoints (e.g., hatching rate) that were previously observed at higher concentrations ($\geq 50 \mu\text{g/mL}$) and longer exposure times (> 72 hpf) for the compound.¹⁴

The ^1H HR-MAS spectra were investigated by PCA, and subsequent partial least square discriminant analysis (PLS-DA) modeling ($R^2 = 0.995$, $Q^2 = 0.999$) to probe if control and PMA-treated embryo can be discriminated, and to determine




FIG. 2. One-dimensional ^1H HR-MAS NMR obtained from intact zebrafish embryos (12 h post-fertilization). All HR-MAS NMR measurements are done at 4°C with a spinning speed of 6 kHz, cycle delay of 2 s, and total number of scan 256. Water suppression was done for obtaining enhanced signals from metabolites. ^1H shift was calibrated using TSP as an internal standard. Glc, glucose; Lac, lactate; Ala, alanine; Gln, glutamine; Glu, glutamate; Gly, glycine; Tau, taurine; m-Ins, (myo)-inositol; Cho, choline; tCr, total creatine; GABA, γ -aminobutyric acid; Asp, aspartate; GSH, glutathione; Ile, isoleucine; Chol, cholesterol; FA, fatty acid; HR-MAS, high-resolution magic angle spinning; NMR, nuclear magnetic resonance; TSP, 3-trimethylsilyl-2,2,3,3-tetra-deuteriopropionic acid.

the spectral regions and corresponding compounds mainly responsible for separation. The PLS-DA scores plot and loading plot are presented in Figure 6. The score plot explains 82.4% of total variance of control sample clustering in the negative PC1 scores and PMA-treated samples in the positive PC2 scores (Fig. 6A). To determine the variables, that is, metabolites assigned to the corresponding buckets that are

mainly responsible for the separation of two groups, the load values or weights of the PC1 were analyzed (Fig. 6B). Signals coming from m-Ins, PC, Glc, Chol, and FA have a positive score in the PC1 loading plot within 95% significance interval ($p < 0.05$), and signals coming from GABA, Glu, and Lac have a negative score in the PC1 loading plot. All of these metabolites have loading within 95% confidence level




FIG. 3. Two-dimensional $[^1\text{H}-^1\text{H}]$ homonuclear correlation HR-MAS NMR spectrum obtained from intact zebrafish embryos (12 h post-fertilization). The ^1H shift was calibrated using TSP as an internal standard. PC, phosphoryl choline; Leu, leucine; Thr, threonine.

TABLE 1. ^1H CHEMICAL SHIFT ASSIGNMENT OF METABOLITES IN ZEBRAFISH EMBRYOS

Metabolite	Abbreviation	Group	Chemical shift (ppm)
Alanine	Ala	^2CH	3.7680
		$^3\text{CH}_3$	1.4655
Aspartate	Asp	^2CH	3.8867
		$^3\text{CH}_2$	2.8021
			2.6508
GABA	GABA	$^2\text{CH}_2$	3.0082
		$^4\text{CH}_2$	2.2828
		$^3\text{CH}_2$	1.8888
Choline	Cho	$\text{N}(\text{CH}_3)_3$	3.1890
Creatine	Cr	$^2\text{CH}_2$	3.9110
		$\text{N}(\text{CH}_3)$	3.0260
β -Glucose	Glc	^1CH	4.630
		$^2\text{CH}_2$	3.230
Glutamate	Glu	^2CH	3.7444
		$^4\text{CH}_2$	2.3354
Glutamine	Gln	^2CH	3.7625
		$^4\text{CH}_2$	2.4350
		$^3\text{CH}_2$	2.1360
Glycine	Gly	^2CH	3.5450
Myo-inositol	m-Ino	$^{4,6}\text{CH}$	3.6114
		^5CH	3.2652
Scyllo-inositol	s-Ino	^{1-6}CH	3.3340
Lactate	Lac	^2CH	4.0908
		$^3\text{CH}_3$	1.3125
Phosphocreatine	PCr	$^2\text{CH}_2$	3.9260
		$\text{N}(\text{CH}_3)$	3.0280
Taurine	Tau	^1CH	3.4190
		^2CH	3.2473
Threonine	Thr	$^4\text{CH}_3$	1.3169
Tyrosine	Tyr	$^{\alpha}\text{CH}$	3.9299

indicating them to be the most significant metabolites for group separation in the score plot.

Discussion

The current study is the first report of HR-MAS NMR as a means of metabolite profiling in the intact zebrafish embryo system (i.e., without any extraction) and generally demonstrates the technique to be an effective and potentially powerful tool for metabolomics studies of early (e.g., ≤ 36 hpf) developmental stages of this particularly versatile system. Building on previous work and prior NMR-based characterization of the most abundant metabolites found in zebrafish embryos, 2D homonuclear (i.e., COSY) HR-MAS NMR was capable of identifying a wide range of relevant metabolites. Subsequent quantitation of peak area, coupled to PCA, enabled statistical analysis of changes in metabolite profiles.

To explore the applicability of HR-MAS NMR metabolite profiling as a toxicological tool, the technique was specifically applied to the algal-derived PMAs as taxonomically widespread potential contaminants of aquatic systems. Although teratogenicity (i.e., morphological deformities) of PMAs in the zebrafish embryo model has been described,¹⁴ the mechanism of this toxicity and possible biochemical,

molecular, and cellular targets remain currently unknown. Consistent with subacute concentrations and exposure times, no discernible effects on development (e.g., mortality, deformities, hatching rates) were observed. However, embryos exposed to PMA were, indeed, characterized by statistically significant changes in several relevant metabolites. Moreover, the changes in metabolic profile, associated with PMA-exposure, notably demonstrate a possible convergence of several interrelated biomarkers, which point to a potentially unifying model of the teratogenicity of these compounds.

As one of the most salient observations, particularly in light of the teratogenicity of the PMAs, inositol levels were found to increase significantly in exposed embryos. Derivatives of m-Ins, and phosphatidylinositol derivatives, in particular, have an established role in *Wnt*-mediated pathways of embryo development (including zebrafish).^{25,26} *Wnt* pathways are a well-described group of signal transduction pathways that regulate crucial aspects of cell-fate determination, cell migration, cell polarity, morphological patterning, and organogenesis during embryonic development. Phosphatidylinositol-based signaling is recognized to serve as a key intracellular function in this regard.²⁷ Consistent with a possible role of the *Wnt* pathway is not only PMA teratogenicity in general but also specific inhibition of organogenesis observed in prior studies; more specifically, exposure to PMAs was found to inhibit the formation of both the eye and heart in developing embryos,¹⁴ and both cardiac and ocular development in the zebrafish are, indeed, linked to *Wnt*.^{28,29} It is tempting, therefore, to hypothesize a role for inositol through *Wnt* pathways in the observed teratogenicity of PMAs.

An alternative hypothesis conversely, however, might implicate a more general interaction of PMA with lipids present in cell membranes. Consistent with this hypothesis, a concurrent increase in FA, PC, and Chol was observed for PMA-exposed embryos. These latter metabolites are clearly associated with lipid composition of cell membranes: PC as the “polar head” of many phospholipids and FAs as “non-polar tails,” whereas cholesterol has an established role in maintaining cell membrane integrity and fluidity, and can comprise, in fact, as much as 50% cell membranes. Similarly, inositols (and m-Ins specifically) are components of—and, in turn, are derived from hydrolytic cleavage of—several phospholipids of membranes, including phosphatidylinositols involved in the *Wnt* pathways.

As such, the combined observation of elevated levels of these four metabolites by HR-MAS may represent a convergence of indicators reflecting possible interactions of PMA with membrane lipids and subsequent disruption of membranes. It has, in fact, been previously proposed based on 3D modeling that PMAs—and the repetitively poly-methoxylated backbone, in particular, which forms a hydrophobic “corkscrew”-like conformation—might interact with lipids and consequently disrupt membranes.³⁰ Accordingly, disruption of cell membranes could be postulated to potentiate consequent hydrolysis of phospholipids (through access to phospholipases) and consequent release (i.e., intracellular elevation) of free PC, m-Ins, and FA. Simultaneously, increased synthesis or, alternatively, release from lipoproteins of cholesterol might, in turn, serve a compensatory response to restore membrane integrity. The increased synthesis of cholesterol as a mechanism to maintain




FIG. 4. Effect of PMA treatment on the metabolic profile of intact zebrafish embryos measured by one-dimensional ^1H HR-MAS NMR. Representative spectra were obtained after treatment of zebrafish embryos (12 h postfertilization) for 24 h with (A) either solvent carrier (control) or (B) PMA, followed by washing three times with buffer. All HR-MAS NMR measurements are done at 4°C with a spinning speed of 6 kHz, cycle delay of 2 s, and total number of scan 256. Water suppression was done for obtaining enhanced signals from metabolites. ^1H shift was calibrated using TSP as an internal standard.

homeostasis has, in fact, precedence in the study of alkylphospholipids which, as proposed for PMA, interact with, and consequently, disrupt membranes.³¹

Another compelling observation in our study is the significant decrease in several interrelated metabolites associated with neuronal function and development. Among these, there is a decrease in Lac concurrent with observed increases

in Glc. Numerous studies suggest that lactate produced from glucose through glycolysis in supporting glial cells (i.e., astrocytes, oligodendrocytes)—and supplied, in turn, to neurons through the so-called astrocyte-neuron lactate shuttle—is the primary energy source for neurons.^{32–35} It has been, furthermore, shown that consumption of energy, in the form of lactate, is particularly important during neuronal development.




FIG. 5. The concentration of metabolites in the zebrafish embryos treated with or without PMA. Zebrafish embryos (12 h postfertilization) were treated for 24 h with either solvent carrier (control) or PMA, followed by washing three times. Representatives of metabolites that show decrease (A) or increase (B) in their concentrations, relative to tCr after PMA treatment. Statistical analyses (*t*-tests and ANOVAs) of the NMR quantification results were performed with OriginPro v. 8. Values calculated from eight spectra (** $p < 0.001$, * $p < 0.05$).




FIG. 6. (A) Partial least square discriminant analysis score plots and (B) loading plot of embryos after treatment with either solvent carrier (control) or PMA for 24 h. Loading plots show assigned peaks from metabolites responsible for clustering in the score plot.

Also, similarly, several lines of evidence suggest that supply of lactate, as a carbon source (as well as energy source), is essential to myelinogenesis in glial cells during embryo development.³⁶

Consistent with reduced glycolysis is decreased levels of GSH. As a key detoxifying molecule, glutathione is particularly important for removing reactive oxygen species and other cytotoxic species (e.g., methylglyoxal³⁷) generated during glycolysis. The reduced levels of GSH could, therefore, reflect a decreased cellular demand, and thus synthesis, due to the reduced levels of glycolysis. However, the observed reduction in GSH could alternatively, or additionally, be explained (as discussed below) by depletion due to decreased levels of key biosynthetic precursors (i.e., glutamate, glycine) of this tripeptide.

Aligned with proposed effects of PMA on central nervous system (CNS) development, simultaneous decreases in Gly, Glu, Gln, and GABA (as key biomarkers of neuronal function and development) were observed. Glutamate is the most abundant and important excitatory neurotransmitter in the brain, whereas GABA and glycine are the most abundant inhibitory neurotransmitters in the brain and spinal cord/brain stem, respectively. Accordingly, the three metabolites are essential mediators of neuronal activity in the adult brain, but known to play, in addition, a key role in CNS development.^{38–40} Glutamine, conversely, is a well-known intermediary of GABA and glutamate recycling by way of the so-called glutamate/glutamine cycle between glia and

neurons.⁴¹ Similarly, glial cells are recognized to play a key role in the cycling of glycine; specifically, astrocytes function both in the uptake of glycine from synapses and, in turn, supply glycine and L-serine (as a biosynthetic precursor of glycine) to neurons.⁴² Furthermore, aside from their well-known roles in CNS function and development, glutamate and glycine are biosynthetic precursors of glutathione, and decreases in the two could, therefore, additionally explain the observed GSH decrease. Indeed, it has been well established that reduction in the precursors (i.e., glutamate, glycine, and cysteine) can, in fact, lead to cellular depletions of GSH in glia and neurons.⁴³

These convergent findings, taken together at the cellular level, may specifically point to a possible role of myelogenic glial cells of the developing zebrafish embryo with respect to the developmental toxicity of PMAs. Implicating a possible involvement of myelogenic glia in the observed PMA toxicity, cholesterol, and phospholipids (containing either PC, e.g., sphingomyelin or inositol), which are, as discussed above, elevated in the PMA-exposed embryos are essential components of myelin. In addition, glial cells—and particularly astrocytes—have an established role in both energy supply (glycolysis) to neurons and recycling of glycine, glutamate, and GABA (as discussed above) to neurons.^{32–35,41,42} Moreover, glial cells are specifically derived from neural crest cells within developing embryos and represent a particularly compelling target for PMAs. In parallel to observed effects on CNS biomarkers, *Wnt* pathways (as a possible target of PMAs,




FIG. 7. Proposed model of the teratogenicity of PMA in relation to the observed changes in metabolic profile. According to this working model, PMA interacts with membrane lipids, leading to hydrolysis of phospholipids, and consequent release (and, thus, elevated levels) of PC, Ins, and FA, as well as a compensatory increase in Chol. Disruption of membrane lipids would, in turn, interfere with signaling involved in *Wnt* developmental pathways. The specific inhibition of neural crest-derived glial cells would, thereby, lead to reduced glycolysis (i.e., decreased Lac, increased Glc) as key energy source for neurons, as well as reduced uptake of Glu, and subsequent synthesis of Gln, leading to impaired recycling of Glu and GABA through glutamate/glutamine cycle.

see above) are well known to control neural crest cell fate (i.e., differentiation and migration),⁴⁴ including subsequent role in the development of eye and heart, which are observably affected in the acute teratogenicity of PMAs.¹⁴ Similarly, one of the additional hallmarks of the developmental toxicity of PMAs is lack of development of melanophores which are, likewise, derived from the neural crest through *Wnt*.⁴⁵

Taken together, therefore, the metabolic profile of PMA-exposed embryos would be consistent with a general unifying model, through which cell-specific effects of the PMAs, and particularly, impairment of neural crest (i.e., myelogenic glial) cells through interference with *Wnt* pathways, could explain observed changes, including (1) elevated free PC, m-Ins, and FA, following proposed disruption of cell membranes and subsequent hydrolysis of phospholipids, along with concomitant compensatory increases in Chol synthesis and/or uptake; (2) increased m-Ins specifically consistent with interference of phosphatidylinositol signaling in *Wnt*-based pathways, including in particular, determination of neural crest cell fate; (3) decreased Lac/increased Glc indicative of reduced glycolysis in glia which, otherwise, support both myelogenesis and energy demands of neurons; and (4) consequently decreased GABA, Gly, Glu, and Gln suggestive, in general, of a loss of glial cell viability and/or function, for example, neurotransmitter recycling, in relation to CNS development (Fig. 7).

In conclusion, the novel application of HR-MAS NMR to intact zebrafish embryos was found to be, in general, an effective technique for metabolic profiling and specifically a promising approach to identifying potential targets/mechanisms and biomarkers related to toxic, or otherwise bioactive, metabolites (as illustrated, in the current study, by the algal-derived PMAs). Although future experimental studies are clearly required to confirm the model/hypotheses generated by the current HR-MAS analysis, these studies do, indeed, demonstrate the considerable potential

of this technique—as a metabolomics tool—for toxicological studies with diverse applicability, in particular, to the zebrafish embryo model.

Acknowledgments

Support for this research was provided by NSF *Catalyzing New International Collaboration* grant (IIA-1427797). The authors sincerely thank the laboratory of Dr. Hermann Spaik at Leiden University and Dr. Stefan Scholz at Helmholtz Centre for Environmental Research (UFZ) Leipzig for providing zebrafish embryos and associated research support used in exposure studies.

Disclosure Statement

No competing financial interests exist.

References

1. Moestue S, Sitter B, Bathen TF, Tessem MB, Gribbestad IS. HR MAS MR spectroscopy in metabolic characterization of cancer. *Curr Top Med Chem* 2011;11:2–26.
2. Kabli S, Alia A, Spaik HP, Verbeek FJ, De Groot HJ. Magnetic resonance microscopy of the adult zebrafish. *Zebrafish* 2006;3:431–439.
3. Kabli S, He S, Spaik HP, Hurlstone A, Jagalska ES, De Groot HJ, *et al.* *In vivo* magnetic resonance imaging to detect malignant melanoma in adult zebrafish. *Zebrafish* 2010;7:143–148.
4. Kabli S, Spaik JP, De Groot HJ, Alia A. *In vivo* metabolite profile of adult zebrafish brain obtained by high-resolution localized magnetic resonance spectroscopy. *J Magn Reson Imaging* 2009;29:275–281.
5. Van Amerongen YF, Roy U, Spaik HP, de Groot HJ, Huster D, Schiller J, *et al.* Zebrafish brain lipid characterization and quantification by ¹H nuclear magnetic

- resonance spectroscopy and MALDI-TOF mass spectrometry. *Zebrafish* 2014;11:240–247.
6. Ali S, Champagne DL, Alia A, Richardson MK. Large-scale analysis of acute ethanol exposure in zebrafish development: a critical time window and resilience. *PLoS One* 2011;6:e20037.
 7. Dai Y, Jia Y, Chen N, Bian W, Li Q, Ma Y, *et al.* Zebrafish as a model system to study toxicology. *Environ Toxicol Chem* 2014;33:11–17.
 8. Nishimura Y, Inoue A, Sasagawa S, Koiwa J, Kawaguchi K, Kawase R, *et al.* Using zebrafish in systems toxicology for developmental toxicity testing. *Congenit Anom* 2016;56:18–27.
 9. Bugel SM, Tanguay RL, Planchart A. Zebrafish: a marvel of high-throughput biology for 21st century toxicology. *Curr Environ Health Rep* 2014;1:341–352.
 10. Berry JP, Gantar M, Gibbs PD, Schmale MC. The zebrafish (*Danio rerio*) embryo as a model system for identification and characterization of developmental toxins from marine and freshwater algae. *Comp Biochem Physiol C Toxicol Pharmacol* 2007;145:61–72.
 11. Berry JP, Gibbs PD, Schmale MC, Saker ML. Toxicity of cylindrospermopsin, and other apparent metabolites from *Cylindrospermopsis raciborskii*. *Toxicol* 2009;53:289–299.
 12. Notch EG, Minuitti DM, Berry JP, Mayer GD. Cyanobacterial LPS potentiates cadmium toxicity in zebrafish (*Danio rerio*) embryos. *Environ Toxicol* 2011;26:498–505.
 13. Jaja-Chimedza A, Gantar M, Mayer GD, Gibbs PD, Berry JP. Effects of cyanobacterial lipopolysaccharides from *Microcystis* on glutathione-based detoxification pathways in zebrafish (*Danio rerio*) embryo. *Toxins* 2012;4:390–404.
 14. Jaja-Chimedza A, Gantar M, Gibbs PD, Schmale MC, Berry JP. Polymethoxy-1-alkenes from *Aphanizomenon ovalisporum* inhibit vertebrate development in the zebrafish (*Danio rerio*) embryo model. *Mar Drugs* 2012;10:2322–2336.
 15. Walton K, Gantar M, Gibbs PD, Schmale MC, Berry JP. Indole alkaloids from *Fischerella* inhibit vertebrate development in the zebrafish (*Danio rerio*) embryo model. *Toxins* 2014;6:3568–3581.
 16. Jaja-Chimedza A, Saez C, Sanchez K, Gantar M, Berry JP. Identification of teratogenic polymethoxy-1-alkenes from *Cylindrospermopsis raciborskii* and taxonomically diverse freshwater cyanobacteria and green algae. *Harmful Algae* 2015;49:156–161.
 17. Mynderse JS, Moore RE. Isotactic polymethoxy-1-alkenes from blue-green alga *Tolypothrix conglutinata* var. *chlorata*. *Phytochemistry* 1979;18:1181–1183.
 18. Mori Y, Kohchi Y, Suzuki M, Carmeli S, Moore RE, Patterson GML. Isotactic polymethoxy-1-alkenes from blue-green algae. Synthesis and absolute stereochemistry. *J Org Chem* 1991;56:631–637.
 19. Mori Y, Kohchi Y, Noguchi H, Suzuki M, Carmeli S, Moore RE, *et al.* Isotactic polymethoxy-1-alkenes from the terrestrial blue-green alga, *Scytonema ocellatum*: structure and synthesis. *Tetrahedron* 1991;47:4889–4904.
 20. Banker R, Teltsch B, Sukenik A, Carmeli S. 7-epicylindrospermopsin, a toxic minor metabolite of the cyanobacterium *Aphanizomenon ovalisporum* from Lake Kinneret, Israel. *J Nat Prod* 2000;63:387–389.
 21. Brand M, Granato M, Nüsslein-Volhard C: Keeping and raising zebrafish. In: *Zebrafish*. Nüsslein-Volhard C and Dahm R (eds), pp. 7–37, Oxford University Press, Oxford, United Kingdom, 2002.
 22. Govindaraju V, Young K, Maudsley AA. Proton NMR chemical shifts and coupling constants for brain metabolites. *NMR Biomed* 2000;13:129–153.
 23. Akhtar MT, Mushtaq MY, Verpoorte R, Richardson MK, Choi YH. Zebrafish as a model for systems medicine R&D: rethinking the metabolic effects of carrier solvents and culture buffers determined by ¹H NMR metabolomics. *OMICS* 2016;20:42–52.
 24. Markley JL, Anderson ME, Qiu C, Eghbalian HR, Lewis IA, Hegeman AD, *et al.* New bioinformatics resources for metabolomics. *Pac Symp Biocomput* 2007;12:157–168.
 25. Gao Y, Wang H. Inositol pentakisphosphate mediates *Wnt/β*-catenin signaling. *J Biol Chem* 2007;282:26490–26502.
 26. MacDonald BT, Tamai K, He X. *Wnt/β*-catenin signaling: components, mechanisms and diseases. *Dev Cell* 2009;17:9–26.
 27. Nusse R, Varmus HE. *Wnt* genes. *Cell* 1992;69:1073–1087.
 28. Fuhrmann S. *Wnt* signaling in eye organogenesis. *Organogenesis* 2008;4:60–67.
 29. Dohn TE, Waxman JS. Distinct phases of *Wnt/β*-catenin signaling direct cardiomyocyte formation in zebrafish. *Dev Biol* 2012;361:364–376.
 30. Jaja-Chimedza A. Contribution of lipophilic secondary metabolites to the toxicity of strains of freshwater cyanobacterial harmful algal blooms, identified using the zebrafish (*Danio rerio*) embryo as a model for vertebrate development [Ph.D. dissertation], Florida International University, Miami, FL, August 2014.
 31. Korade Z, Kenworthy AK. Lipid rafts, cholesterol, and the brain. *Neuropharmacology* 2008;55:1265–1273.
 32. Carrasco MP, Jimenez-Lopez JM, Rios-Marcos P, Segovia JL, Marco C. Disruption of cellular cholesterol transport and homeostasis as a novel mechanism of membrane-targeted alkylphospholipid analogues. *Br J Pharmacol* 2010;160:355–366.
 33. Bogdanowicz P, Pujol JP. Glycophosphatidylinositol (GPI) hydrolysis by transforming growth factor- β 1 (TGF- β 1) as a potential early step in the inhibition of epithelial cell proliferation. *Mol Cell Biochem* 2000;208:143–150.
 34. Grocott T, Johnson S, Bailey AP, Streit A. Neural crest cells organize the eye via TGF- β and canonical *Wnt* signaling. *Nat Commun* 2011;2:265.
 35. Rinholm JE, Bergerson LH. White matter lactate—does it matter? *Neuroscience* 2014;276:109–116.
 36. Belanger M, Allaman I, Magistretti PJ. Brain energy metabolism: focus on astrocyte neuron metabolic cooperation. *Cell Metab* 2011;14:724–738.
 37. Prichard J, Rothman D, Novotny E, Petroff O, Kuwabara T, Avison M, *et al.* Lactate rise detected by ¹H NMR in human visual cortex during physiologic stimulation. *Proc Natl Acad Sci U S A* 1991;88:5829–5831.
 38. Ben-Ari Y. Excitatory actions of GABA during development: the nature of the nurture. *Nat Rev Neurosci* 2002;3:728–739.
 39. Lujan R, Shigemoto R, Lopez-Bendito G. Glutamate and GABA receptor signaling in the developing brain. *Neuroscience* 2005;130:567–580.
 40. Avila A, Nguyen L, Rigo J. Glycine receptors and brain development. *Front Cell Neurosci* 2013;7:184.

41. Bak LK, Schousboe A, Waagepetersen HS. The glutamate/GABA-glutamine cycle: aspects of transport, neurotransmitter homeostasis and ammonia transfer. *J Neurochem* 2006;98: 641–653.
42. Furuya S, Tabata T, Mitoma J, Yamada K, Yamasaki M, Makino A, *et al.* L-Serine and glycine serve as major astroglia-derived trophic factors for cerebellar Purkinje neurons. *Proc Natl Acad Sci U S A* 2000;97:11528–11533.
43. Dringen R, Hamprecht B. Glutathione restoration as indicator for cellular metabolism of astroglial cells. *Dev Neurosci* 1998;20:401–407.
44. Dorsky RI, Moon RT, Raible DW. Control of neural crest cell fate by the Wnt signaling pathway. *Nature* 1998;396: 370–373.
45. Curran K, Raible DW, Lister JA. Foxd3 controls melanophore specification in the zebrafish neural crest by regulation of Mitf. *Dev Biol* 2009;332:408–417.

Address correspondence to:

A. Alia, PhD
*Institute of Medical Physics and Biophysics
University of Leipzig
Härtelstr. 16-18
Leipzig D-04107
Germany*

E-mail: a.alia@chem.leidenuniv.nl

John P. Berry, PhD
*Department of Chemistry and Biochemistry
Florida International University
3000 NE 151st Street
North Miami, FL 33181*

E-mail: berryj@fiu.edu