

Lumbar lordosis and extensor muscle size

1 **The relationship between sagittal curvature and extensor muscle volume in** 2 **the lumbar spine.**

3

4 Judith R. Meakin^{1,2}, Jonathan Fulford³, Richard Seymour⁴, Joanne R Welsman⁵, Karen M.
5 Knapp²

6

7 ¹Biomedical Physics Group, College of Engineering, Mathematics and Physics Sciences,
8 University of Exeter

9 ²Medical Imaging, College of Engineering, Mathematics and Physics Sciences, University of
10 Exeter

11 ³Exeter NIHR Clinical Research Facility, University of Exeter

12 ⁴Torbay Hospital, South Devon Healthcare NHS Foundation Trust

13 ⁴Sport and Health Sciences, University of Exeter

14

15

1 **Summary**

2 A previous modelling study predicted that the forces applied by the extensor muscles to stabilise
3 the lumbar spine would be greater in spines that have a larger sagittal curvature (lordosis). Since
4 the force generating capacity of a muscle is related to its size, it was hypothesised that the size of
5 the extensor muscles in a subject would be related to the size of their lumbar lordosis. MR image
6 data was obtained, together with age, height, body mass and back pain status, from 42 female
7 subjects. The volume of the extensor muscles (multifidus and erector spinae) caudal to the mid-
8 lumbar level was estimated from cross-sectional area measurements in axial T1-weighted MR
9 images spanning the lumbar spine. Lower lumbar curvature was determined from sagittal T1-
10 weighted images. A stepwise linear regression model was used to determine the best predictors
11 of muscle volume. The mean lower lumbar extensor muscle volume was 281 cm³ (sd 49 cm³).
12 The mean lower lumbar curvature was 30° (sd 7°). Five subjects reported current back pain and
13 were excluded from the regression analysis. Nearly half the variation in muscle volume was
14 accounted for by the variables age (standardised coefficient, B = -3.2, P = 0.03) and lower
15 lumbar curvature (B = 0.47, P = 0.002). The results support the hypothesis that extensor muscle
16 volume in the lower lumbar spine is related to the magnitude of the sagittal curvature; this has
17 implications for assessing muscle size as an indicator of muscle strength.

18 **Key Words**

19 Extensor muscle size; Lumbar lordosis; Magnetic resonance imaging

20

1 **Introduction**

2 The extensor muscles of the lumbar spine, located posteriorly to the vertebral bodies play an
3 important role in controlling movement and providing mechanical stability. They comprise two
4 main muscle groups: the transversospinalis (multifidus, rotatores, interspinales and
5 intertransversarii) and the erector spinae (iliocostalis and longissimus). The transversospinalis
6 are located deeply, attaching to the lumbar vertebrae, and are considered responsible for small
7 movements stabilising the spine (Macintosh & Bogduk, 1986; Hansen et al., 2006; Ward et al.,
8 2009; Cornwall et al., 2011). The erector spinae, situated more superficially, spanning larger
9 sections of the spine, are considered to have a greater role in producing spinal movement
10 (Macintosh & Bogduk, 1991; Hansen et al., 2006).

11 The mechanical stability of the lumbar spine is achieved when the resultant forces in the spine
12 travel tangentially to the sagittal spinal curvature (Aspden, 1989); these allow the lumbar spine
13 to support the weight of the upper body without buckling and have been termed ‘follower loads’
14 (Patwardhan et al., 2000). Several modelling studies have investigated the mechanisms by which
15 follower loads can be achieved and have demonstrated that the extensor muscles could play a
16 role in providing the forces needed to generate a follower load, maintaining the stability of the
17 spine with little deformation, shear or bending (Kim et al., 2007; Kim & Kim, 2008; Han et al.,
18 2011).

19 A recent modelling study has investigated how the forces required to produce a follower load
20 would be affected by differences in the magnitude of the curvature (Meakin & Aspden, 2012).

21 The range of lumbar sagittal curvature, lordosis, in the normal population is considerable
22 (Berthonnaud et al., 2005; Boulay et al., 2006; Meakin et al., 2008) and although the curvature is

Lumbar lordosis and extensor muscle size

1 modified by changes in posture (Meakin et al., 2009), it has been suggested that a proportion of
2 the inter-subject variability is due to intrinsic factors such as the bony anatomy (Meakin et al.,
3 2009). The model considered the spine as a two-dimensional structure with forces applied at
4 each vertebral level in a simplified representation of the extensor muscles and did not investigate
5 individual muscle groups. It predicted, however, that a lumbar spine with a large lordosis would
6 require larger forces to produce a follower load in the standing posture than a spine with a small
7 lordosis (Meakin & Aspden, 2012). This was particularly so in the lower regions of the spine
8 where the majority of the lumbar lordosis is usually located and the predictions have implications
9 for understanding how the muscles provide biomechanical stability to the lumbar spine in a range
10 of individuals.

11 The maximum force that a muscle can apply is limited by its size; this is because the force
12 generating capacity of a muscle is, in theory, directly proportional to the number and size of the
13 muscle fibres within it (Brinckmann et al., 2000). In practice, this relationship is modified
14 (positively or negatively) by various factors including the type of muscle fibres present, their
15 length, and their arrangement in the tissue (Brinckmann et al., 2000; Jones et al., 2008) together
16 with the neural control of the muscle and the presence of non-contractile tissue such as fat (Jones
17 et al., 2008). It is generally accepted, however, that muscle strength is proportional to muscle
18 size (Brinckmann et al., 2000; Folland & Williams, 2007; Jones et al., 2008) and many studies
19 have found the size of a muscle (measured as an area or a volume) to be highly correlated with
20 its ability to apply force (Akagi et al., 2009; Blazevich et al., 2009).

21 Measurements of lumbar extensor muscle size show that, similarly to spine shape, there is
22 considerable variability in size in the normal population, especially in the lower lumbar regions
23 (Danneels et al., 2000; Kamaz et al., 2007; Hides et al., 2008; Wallwork et al., 2009). This

Lumbar lordosis and extensor muscle size

1 variation in muscle size indicates a concomitant variation in muscle strength such that an
2 individual with small extensor muscles will not be able to generate the same magnitude of spine
3 stabilisation force as an individual with large extensor muscles. The relationship between muscle
4 size and lumbar lordosis has not previously been investigated but, based on the predictions of the
5 recent modelling study (Meakin & Aspden, 2012), we hypothesised that subjects with larger
6 lumbar lordosis, particularly in the lower half, would have more muscle (thus enabling them to
7 stabilise their spine appropriately). The aim of the study reported here was to test this hypothesis
8 in a sample of volunteer subjects.

9 **Methods**

10 *Data*

11 Magnetic Resonance Imaging (MRI) data from forty-two female subjects was acquired together
12 with the age, height, body mass and back pain status of the subjects. The images, which were
13 collected for a previous study (Knapp et al., 2012), were acquired using a Philips Gyroscan
14 Intera 1.5 T MRI scanner (Philips Medical Systems, The Netherlands). Back pain status was
15 ascertained from the answer to the question “Do you, or have you in the past suffered from back
16 pain?” that was administered via a written questionnaire: Approval for the study was given by
17 Devon and Torbay Research Ethics Committee (08/H0202/109) and all subjects gave their
18 written informed consent.

19 *Muscle size*

20 Images acquired in the axial plane were used to determine extensor muscle size in the lower part
21 of the lumbar spine. These images were obtained using a gradient echo sequence with a
22 repetition time (TR) of approximately 55 ms and an echo time (TE) of 1.9 ms. Forty 8 mm thick

Lumbar lordosis and extensor muscle size

1 slices, with a 1 mm gap were acquired; the in-plane resolution was between 1.76 and 1.95
2 mm/pixel, depending on subject size.

3 The MR images were saved in TIFF format and regions of interest corresponding to the extensor
4 muscles were segmented, by one observer, using GIMP software (GNU Image Manipulation
5 Programme, version 2.6.10, www.gimp.com). The regions of interest (Figure 1) included the
6 iliocostalis, longissimus and multifidus muscles and may have included the rotatores,
7 interspinales and intertransversarii muscles. Macroscopic infiltrations of fat were excluded. No
8 attempt was made to segment the muscles separately as the demarcation between them was not
9 clear on all the images. The number of segmented pixels was multiplied by the in-plane
10 resolution to give the cross-sectional area.

11 The extensor muscle volume caudal to the level of L3/L4 was estimated by multiplying the
12 cross-sectional areas by the effective slice thickness (acquired slice thickness plus slice gap = 0.9
13 cm) and summing across the relevant number of slices. The first of these slices was defined as
14 that in which the presence of the L3/L4 disc was observed; the last was defined as the most
15 caudal slice in which extensor muscle was observed. The L3/L4 disc was chosen to define the
16 upper boundary of the muscle volume because it provided an identifiable anatomical plane that
17 was approximately parallel to the axial slices in all the subjects (thus minimising inter-subject
18 artefacts in the volume measurement). The variation in the height of the subjects meant that the
19 number of slices varied from 10 to 15 (mean = 13).

20 The segmentation of the slice at the level of L3/L4 was performed twice by one observer so that
21 intra-observer reliability and measurement error could be assessed. The segmentation of this
22 slice was also performed by a second observer so that inter-observer reliability could be
23 assessed.

1 ***Spine shape***

2 Images acquired in the sagittal plane were used to determine the curvature of the lower lumbar
3 spine. These images were acquired using a T1-weighted turbo-spin echo sequence with a TR of
4 approximately 198 ms and a TE of 7 ms. Slices were 4 mm thick with an in-plane pixel size
5 between 0.59 and 0.68 mm, depending on body size. The slice that was as close as possible to the
6 mid-line of the lumbar spine was identified (by observing the presence of the conus medullaris
7 and/or the spinous processes) and saved in BMP format (Figure 2). For eight subjects, these
8 sagittal images did not show the full length of the lumbar spine and so survey images, used for
9 positioning the axial slices, were used instead. The survey images included 10 slices, acquired in
10 the sagittal plane, with a slice thickness of 10 mm and an in-plane pixel size of 1.76 mm. The
11 survey slice closest to the mid-line of the lumbar spine was identified as described above.

12 The curvature was characterised as the angle between the superior endplates of L4 and S1
13 (θ_{L4S1}). This angle was determined using a procedure described in a previous paper (Ali et al.,
14 2012), where output from an active shape model is used to determine the end-plate angles. The
15 active shape model was generated using freely available software tools from the University of
16 Manchester, UK in the same way as described previously (Meakin et al., 2008). This involved
17 one observer placing 168 landmark points around the periphery of the vertebral bodies from L1
18 to S1 in each image. The software then transformed the data, to remove scale and rigid body
19 movement, and performed principle component analysis to give ‘modes of variation’ describing
20 patterns of variation in the spine shape. The output from this model was subsequently processed
21 using in-house Matlab code (version R2010a, The MathWorks Inc. Natick MA, USA) to
22 determine the superior end-plate angles. Based on our previous work (Meakin et al., 2008; Ali et

1 al., 2012), and the resolution of the MRI data, we estimate that the angle θ_{L4S1} could be
2 determined to within 1 or 2 degrees.

3 *Data analysis and statistics*

4 Statistical analysis of the data was performed using SPSS software (version 18, IBM
5 Corporation, New York, USA). The Kolmogorov–Smirnov test was used to establish whether
6 data followed a normal distribution. The inter- and intra-observer reliability of the cross-
7 sectional area measurements was assessed by determining the single measures intra-class
8 correlation coefficient (using a one-way random model). Measurement error was determined as
9 2.77 times the within subject standard deviation calculated using a one-way analysis of variance
10 of the repeat measurements made by the first observer. The volume measurement error was
11 estimated by multiplying the cross-sectional area measurement error by the square root of 13
12 (mean number of slices measured). Analysis of variance was used to assess differences between
13 groups; post-hoc testing of differences was performed with the Sidak correction for multiple
14 comparisons. The strength of the relationship between variables was assessed from the Pearson
15 correlation coefficient. Stepwise linear regression was used to determine the best model for
16 muscle volume using input variables age, height, body mass and θ_{L4S1} .

17 **Results**

18 The 42 subjects had a mean age of 50 years (sd 13 years), a mean height of 165 cm (sd 7 cm) and
19 a median mass of 70 kg (inter quartile range 18 kg). Their mean lower lumbar curvature, θ_{L4S1} ,
20 was 30° (sd 7°) and their mean extensor muscle volume caudal to L3/L4 was 281 cm^3 (sd 49
21 cm^3). The reliability of the cross-sectional area measurements was excellent with an intra-

Lumbar lordosis and extensor muscle size

1 observer ICC of 0.97 (95% CI 0.94 – 0.98) and an inter-observer ICC of 0.77 (95% CI 0.61 –
2 0.87). Measurement error was 3 cm² for cross-sectional area and 10 cm³ for volume.

3 Eleven subjects reported no back pain, twenty-six subjects reported back pain in the past, and
4 five subjects reported current back pain. Table 1 shows the subject characteristics broken down
5 by back pain status. There was a statistically significant difference in the muscle volume between
6 the three groups but not for the other parameters (Table 1); post-hoc testing showed that the
7 muscle volume of the group with current back pain was significantly smaller (95% CI 11 – 135
8 cm³) than the group that reported no back pain.

9 Excluding the subjects with current back pain, the muscle volume was found to be negatively
10 correlated with age and positively correlated with height, body mass and lower lumbar curvature
11 (Table 2). The relationships were statistically significant with the exception of that between
12 volume and body mass. Of the variables included in the stepwise linear regression (age, height,
13 body mass and lower lumbar curvature), only age and lower lumbar curvature were found to be
14 significant predictors of muscle volume (Table 3). These two parameters accounted for 46 % of
15 the variation in muscle volume. The relationship between the variables is visually demonstrated
16 in Figure 2, which shows the sagittal lumbar curvature for two subjects with differing lumbar
17 muscle volumes.

18 **Discussion**

19 The aim of this study was to test the hypothesis that there would be a relationship between the
20 size of the extensor muscles and the magnitude of the sagittal curvature of the lumbar spine. The
21 hypothesis that a relationship would exist resulted from a model that investigated the effect of
22 lumbar spine shape on the forces required to produce a stabilising follower load in standing

Lumbar lordosis and extensor muscle size

1 (Meakin & Aspden, 2012). The model predicted that larger muscle forces would be required to
2 provide stability in lumbar spines that had larger curvatures and, since the force generating
3 capacity of a muscle is related to its physical size, this then suggested that variation in extensor
4 muscle size (Danneels et al., 2000; Kamaz et al., 2007; Hides et al., 2008; Wallwork et al., 2009)
5 might be related to variation in lumbar curvature (Berthonnaud et al., 2005; Boulay et al., 2006;
6 Meakin et al., 2008). The results of current study support the hypothesis in that larger extensor
7 muscles caudal to L3L4 were associated with subjects who had a greater degree of lumbar
8 curvature between L4 and S1. Although, in the current study, the subjects were supine, the
9 results are relevant to the predictions of the model since the shape of the lumbar spine in
10 standing is highly correlated to that in the supine posture (Meakin et al., 2009) and changes in
11 curvature at the lower lumbar levels have been shown to be only a few degrees (Wood et al.,
12 1996; Andreasen et al., 2007).

13 The correlations between lower lumbar muscle volume and the age, height and body mass of the
14 subjects (Table 2) are consistent with other studies reported in the literature. Ageing is known to
15 be associated with a reduction in the quantity of muscle throughout the body (Janssen et al.,
16 2000; Lang et al., 2010) due to various cellular and molecular changes (Ryall et al., 2008).
17 Skeletal muscle mass is known to be greater in subjects who are taller and heavier (Janssen et al.,
18 2000) and previous studies focusing on lumbar muscles have found a positive correlation with
19 physical size (Chaffin et al., 1990; Jorgensen et al., 2003). Of these factors, however, subject size
20 (i.e. height and body mass) were not found to be significant predictors of lower lumbar muscle
21 volume in the current study. This may be because, in our sample, there was a correlation between
22 age and height (Table 2) which had a p value of 0.06; although not statistically significant at the
23 5 % level, this suggests some covariance between the two variables.

Lumbar lordosis and extensor muscle size

1 The smaller muscle volume in the subjects that reported current back pain (Table 1) is also
2 consistent with the literature. A number of studies have shown that patients with chronic low
3 back pain have small lumbar muscles (usually defined by their cross-sectional area) in
4 comparison to normal volunteers (Danneels et al., 2000; Hides et al., 2008; Kamaz et al., 2007;
5 Wallwork et al., 2009) and patients with non-chronic low back pain (Stokes et al., 1992). This
6 difference in size is particularly prevalent in the multifidus muscles at the lower lumbar levels
7 (Danneels et al., 2000; Hides et al., 2008; Wallwork et al., 2009), although it has also been found
8 in the erector spinae muscles (Kamaz et al., 2007; Stokes et al., 1992). The subjects who reported
9 current back pain in the current study were excluded from the analysis of the relationship
10 between muscle size and lumbar curvature since they formed a very small ($n = 5$) sub-group, and
11 no information was available as to whether their back pain was acute or chronic. Future
12 investigation into the relationship between muscle volume and lumbar curvature in subjects with
13 and without chronic back pain could be useful, however, in furthering our understanding of the
14 relevance of these observations.

15 It is impossible to establish the nature of the causal relationship between spine curvature and
16 muscle size from the current study; however, there are plausible mechanisms that can be inferred
17 from the literature. The first, which was the hypothesis of the current study, is that the curvature
18 of an individual's spine is an intrinsic property and that inter-subject variation in this intrinsic
19 shape then requires a variety of stabilising forces, which manifest as a variety of muscle sizes.
20 Previous studies have suggested that an individual's spine shape is an intrinsic property
21 (Stagnara et al., 1982; Meakin et al., 2009) and the causes for this include the morphology of the
22 vertebrae (Meakin et al., 2009) which is known to vary in terms of size and shape of the vertebral
23 bodies and the size and angulation of the posterior elements (Berry et al., 1987). The relationship

Lumbar lordosis and extensor muscle size

1 between vertebral morphology and spinal curvature has not been previously explored and so we
2 intend to investigate this in a future study. This is an important further step since vertebral
3 morphology may also affect the line of action of the extensor muscles, modifying the forces they
4 apply to the spine (Nussbaum et al., 1995).

5 An alternative mechanism to explain the relationship between spine curvature and muscle size is
6 that the size of an individual's muscles is dictated by various factors including their body size,
7 gender, and lifestyle. This variation in muscle size gives rise to a variation in the ability of the
8 muscles to provide stabilisation to the spine and this causes the spine to conform to a shape that
9 can be successfully stabilised by the available muscle forces. The effect of age on the
10 relationship can then be explained by the age-related loss of muscle mass, described above,
11 having the effect of reducing the strength of the spinal muscles and leading to a reduction in
12 lumbar curvature. A reduction in lumbar lordosis with age has been observed in cross-sectional
13 (Hammerberg & Wood, 2003) and longitudinal studies (Takeda et al., 2009); however, the
14 mechanism of this phenomenon may also be explained by other age-related changes in the spine
15 such as an increase in spinous process height (Aylott et al., 2012) or a decrease in disc height and
16 anterior wedging (Takeda et al., 2009).

17 In the study reported here, extensor muscle size was defined as the volume caudal to the level of
18 L3/L4 as estimated from measurements of cross-sectional area. Most previous studies concerned
19 with lumbar muscle size have considered the cross-sectional area: either in the form of the
20 anatomical cross-sectional area (perpendicular to the long axis of the muscle) or the
21 physiological cross-sectional area (perpendicular to the muscle fibres). In our study, cross-
22 sectional area was measured from images acquired in the axial plane and, due to the varying
23 orientations of the extensor muscles being considered, was not a true anatomical or true

Lumbar lordosis and extensor muscle size

1 physiological cross-sectional area. One problem in defining muscle size by the cross-sectional
2 area, whether it be anatomical or physiological, is that it is influenced by a number of extrinsic
3 factors. Active contraction of a muscle will increase the cross-sectional area from the relaxed
4 state and passive elongation (due to movement of the muscle attachment sites with joint
5 movement) will decrease the area. This second issue is particularly relevant to the spine, since it
6 has a large degree of flexibility, and it has been shown that trunk flexion leads to a reduction in
7 the measured cross-sectional area of the extensor muscles (Jorgensen et al., 2003). Many studies
8 that have measured muscle cross-sectional area have therefore tried to ensure that their subjects
9 were placed in the same posture (Stokes et al., 1992) or have attempted to position their subjects
10 so as to minimise the lordosis (Wallwork et al., 2009). In the current study, every subject was
11 imaged in the supine position with her legs extended; this still resulted in a range of lower
12 lumbar curvatures from 28° to 47° . Attempting to equalise this curvature would have required
13 subjects to adopt different postures and may have been an unrealistic goal in some subjects. By
14 measuring volume, we believe that the problem of passive elongation will be minimised since it
15 is generally considered that muscle tissue is incompressible (Ehret et al., 2011) because of its
16 high water content. This means that although passive elongation would be expected to increase
17 length and reduce area, it would not be expected to alter volume. Although this hypothesis has
18 not been tested in the spine, it is supported by measurements of gastrocnemii volume and length
19 at a range of ankle flexion angles (Barber et al., 2009). In addition to the issue of passive
20 elongation, a further advantage of measuring volume is that it may provide a better indicator of
21 lumbar muscle strength than cross-sectional area (Akagi et al., 2009; Blazevich et al., 2009) thus
22 having more relevance for indicating the ability of the extensor muscles to apply force.

Lumbar lordosis and extensor muscle size

1 One of the limitations of the study was that the resolution of the axial MRI images, although
2 sufficient to segment the muscles as a whole, would ideally have been higher to allow individual
3 muscle groups to be identified. This would have allowed us to establish whether the relationship
4 found between muscle volume and curvature was the same in both the multifidus and erector
5 spinae. A second limitation is that our subjects were exclusively female and we cannot be certain
6 that our results would be the same in male subjects since the size of the lumbar muscles is
7 generally larger in male subjects than female (Marras et al., 2001; Jorgensen et al., 2003; Stokes
8 et al., 2005) and previous studies have found the anthropometric predictors of lumbar muscle
9 size to differ between the sexes (Marras et al., 2001).

10 Despite these limitations, the results of the study show the quantity of extensor muscle present in
11 a subject is related to the magnitude of their lumbar lordosis. This supports the predictions of a
12 recent model that suggested that the muscle forces required for biomechanical stability via a
13 follower load depend on the spinal curvature (Meakin & Aspden, 2012). These results may have
14 implications for assessing the size of muscles in back pain patients. Several studies have found
15 the extensor muscles of back pain patients to be smaller than in normal volunteers (Danneels et
16 al., 2000; Kamaz et al., 2007; Hides et al., 2008; Wallwork et al., 2009) but with some overlap
17 between the two groups. Taking the lumbar curvature into account (together with the other
18 factors that affect muscle size, such as age) may allow the relevance of muscle size to be
19 evaluated, providing a better assessment of whether their muscles are adequate for their needs or
20 would benefit from intensive therapy.

1 **Concluding remarks**

2 The conclusions from this study are that the volume of the lumbar extensor muscles in the lower
3 half of the lumbar spine (caudal to the level of the L3/L4 disc) has a positive correlation with the
4 magnitude of the sagittal lumbar curvature over the same region.

5 **Acknowledgements**

6 We thank the following students, who were on the Medical Imaging (Diagnostic Radiography)
7 programme at the University of Exeter, for helping to collect the data that was used in this study:
8 Susan Hopkins, Sophie Holl, Andrew Bartlett and Soukina May.

9 **Conflict of interest**

10 There are no conflicts of interest to declare.

11 **Author contributions**

12 **Judith Meakin:** developed the hypothesis underlying the study, performed the measurements of
13 muscle size and spine curvature, analysed the data, and wrote the initial drafts of the manuscript.

14 **Jonathan Fulford:** acquired the MRI data, was involved with interpretation of the results and
15 edited and approved the manuscript.

16 **Richard Seymour:** was involved in the data collection and the interpretation of the results and
17 edited and approved the manuscript.

18 **Joanne Welsman:** was involved in the data collection and the interpretation of the results and
19 edited and approved the manuscript.

Lumbar lordosis and extensor muscle size

- 1 **Karen Knapp:** was involved in the data collection and the interpretation of the results and edited
- 2 and approved the current manuscript.
- 3

1 **References**

- 2 Akagi R, Takai Y, Ohta M, Kanehisa H, Kawakami Y, Fukunaga T (2009) Muscle volume
3 compared to cross-sectional area is more appropriate for evaluating muscle strength in
4 young and elderly individuals. *Age and Ageing*, **38**, 564-569.
- 5 Ali A, Cowan A-b, Gregory J, Aspden R, Meakin J (2012) The accuracy of active shape
6 modelling and end-plate measurements for characterising the shape of the lumbar spine in
7 the sagittal plane. *Computer Methods in Biomechanics and Biomedical Engineering*, **15**,
8 167-172.
- 9 Andreassen ML, Langhoff L, Jensen TS, Albert HB (2007) Reproduction of the lumbar lordosis:
10 a comparison of standing radiographs versus supine magnetic resonance imaging
11 obtained with straightened lower extremities. *Journal of Manipulative and Physiological*
12 *Therapeutics*, **30**, 26-30.
- 13 Aspden RM (1989) The spine as an arch. A new mathematical model. *Spine*, **14**, 266-274.
- 14 Aylott CE, Puna R, Robertson PA, Walker C (2012) Spinous process morphology: the effect of
15 ageing through adulthood on spinous process size and relationship to sagittal alignment.
16 *European Spine Journal*, **21**, 1007-1012.
- 17 Barber L, Barrett R, Lichtwark G (2009) Validation of a freehand 3D ultrasound system for
18 morphological measures of the medial gastrocnemius muscle. *Journal of Biomechanics*,
19 **42**, 1313-1319.
- 20 Berry JL, Moran JM, Berg WS, Steffee AD (1987) A Morphometric Study of Human Lumbar
21 and Selected Thoracic Vertebrae. *Spine*, **12**, 362-367.
- 22 Berthonnaud E, Dimnet J, Rousouly P, Labelle H (2005) Analysis of the sagittal balance of the
23 spine and pelvis using shape and orientation parameters. *Journal of Spinal Disorders and*
24 *Techniques*, **18**, 40-7.
- 25 Blazeovich A, Coleman D, Horne S, Cannavan D (2009) Anatomical predictors of maximum
26 isometric and concentric knee extensor moment. *European Journal of Applied*
27 *Physiology*, **105**, 869-878.
- 28 Boulay C, Tardieu C, Hecquet J, et al. (2006) Sagittal alignment of spine and pelvis regulated by
29 pelvic incidence: standard values and prediction of lordosis. *European Spine Journal*, **15**,
30 415-22.
- 31 Brinckmann P, Frobin W, Leivseth G (2000) *Musculoskeletal Biomechanics*, Thieme, Stuttgart.
- 32 Chaffin DB, Redfern DR, Erig M, Goldstein SA (1990) Lumbar muscle size and locations from
33 CT scans of 96 women of age 40 to 63 years. *Clinical Biomechanics*, **5**, 9-16.
- 34 Cornwall J, Stringer MD, Duxson M (2011) Functional Morphology of the Thoracolumbar
35 Transversospinal Muscles. *Spine*, **36**, E1053-E1061.
- 36 Danneels LA, Vanderstraeten GG, Cambier DC, Witvrouw EE, De Cuyper HJ (2000) CT
37 imaging of trunk muscles in chronic low back pain patients and healthy control subjects.
38 *European Spine Journal*, **9**, 266-72.
- 39 Ehret AE, Bül M, Itskov M (2011) A continuum constitutive model for the active behaviour of
40 skeletal muscle. *Journal of the Mechanics and Physics of Solids*, **59**, 625-636.
- 41 Folland J, Williams A (2007) The Adaptations to Strength Training: Morphological and
42 Neurological Contributions to Increased Strength. *Sports Medicine*, **37**, 145-168.
- 43 Hammerberg EM, Wood KB (2003) Sagittal profile of the elderly. *Journal of Spinal Disorders*
44 *and Techniques*, **16**, 44.

- 1 Han K-S, Rohlmann A, Yang S-J, Kim BS, Lim T-H (2011) Spinal muscles can create
2 compressive follower loads in the lumbar spine in a neutral standing posture. *Medical*
3 *Engineering and Physics*, **33**, 472-8.
- 4 Hansen L, de Zee M, Rasmussen J, Andersen TB, Wong C, Simonsen EB (2006) Anatomy and
5 biomechanics of the back muscles in the lumbar spine with reference to biomechanical
6 modeling. *Spine*, **31**, 1888-99.
- 7 Hides J, Gilmore C, Stanton W, Bohlscheid E (2008) Multifidus size and symmetry among
8 chronic LBP and healthy asymptomatic subjects. *Manual Therapy*, **13**, 43-9.
- 9 Janssen I, Heymsfield SB, Wang Z, Ross R (2000) Skeletal muscle mass and distribution in 468
10 men and women aged 18–88 yr. *Journal of Applied Physiology*, **89**, 81-88.
- 11 Jones E, Bishop P, Woods A, Green J (2008) Cross-sectional area and muscular strength: a brief
12 review. *Sports Medicine*, **38**, 987-994.
- 13 Jorgensen MJ, Marras WS, Gupta P (2003) Cross-sectional area of the lumbar back muscles as a
14 function of torso flexion. *Clinical Biomechanics*, **18**, 280-286.
- 15 Kamaz M, Kireşi D, Oğuz H, Emlik D, Levendoğlu F (2007) CT measurement of trunk muscle
16 areas in patients with chronic low back pain. *Diagnostic and Interventional Radiology*,
17 **13**, 144-148.
- 18 Kim K, Kim YH (2008) Role of trunk muscles in generating follower load in the lumbar spine of
19 neutral standing posture. *Journal of Biomechanical Engineering*, **130**, 041005-041005.
- 20 Kim K, Kim YH, Lee S (2007) Increase of load-carrying capacity under follower load generated
21 by trunk muscles in lumbar spine. *Journal of Engineering in Medicine*, **221**, 229-235.
- 22 Knapp KM, Welsman JR, Hopkins SJ, Fogelman I, Blake GM (2012) Obesity increases
23 precision errors in dual-energy x-ray absorptiometry measurements. *Journal of Clinical*
24 *Densitometry*, **Epub ahead of print**.
- 25 Lang T, Streeper T, Cawthon P, Baldwin K, Taaffe D, Harris T (2010) Sarcopenia: etiology,
26 clinical consequences, intervention, and assessment. *Osteoporosis International*, **21**, 543-
27 559.
- 28 Macintosh JE, Bogduk N (1986) The biomechanics of the lumbar multifidus. *Clinical*
29 *Biomechanics*, **1**, 205-213.
- 30 Macintosh JE, Bogduk N (1991) The attachments of the lumbar erector spinae. *Spine*, **16**, 783-
31 792.
- 32 Marras WS, Jorgensen MJ, Granata KP, Wiand B (2001) Female and male trunk geometry: size
33 and prediction of the spine loading trunk muscles derived from MRI. *Clinical*
34 *Biomechanics*, **16**, 38-46.
- 35 Meakin JR, Aspden RM (2012) Modeling the effect of variation in sagittal curvature on the force
36 required to produce a follower load in the lumbar spine. *Journal of Mechanics in*
37 *Medicine and Biology*, **12**, 1-10.
- 38 Meakin JR, Gregory JS, Aspden RM, Smith FW, Gilbert FJ (2009) The intrinsic shape of the
39 human lumbar spine in the supine, standing and sitting postures: characterization using an
40 active shape model. *Journal of Anatomy*, **215**, 206-11.
- 41 Meakin JR, Gregory JS, Smith FW, Gilbert FJ, Aspden RM (2008) Characterizing the shape of
42 the lumbar spine using an active shape model: reliability and precision of the method.
43 *Spine*, **33**, 807-13.
- 44 Nussbaum MA, Chaffin DB, Rechten CJ (1995) Muscle lines-of-action affect predicted forces
45 in optimization-based spine muscle modeling. *Journal of Biomechanics*, **28**, 401-409.

Lumbar lordosis and extensor muscle size

- 1 Patwardhan AG, Havey R, Meade KP, et al. (2000) A Follower Load Stabilizes the Lumbar
2 Spine with Minimal Changes in Sagittal Plane Mobility.), pp. 0389-0389. Orlando,
3 Florida.
- 4 Ryall JG, Schertzer JD, Lynch GS (2008) Cellular and molecular mechanisms underlying age-
5 related skeletal muscle wasting and weakness. *Biogerontology*, **9**, 213-228.
- 6 Stagnara P, Claude de Mauroy J, Dran G, et al. (1982) Reciprocal angulation of vertebral bodies
7 in a sagittal plane: approach to references for the evaluation of kyphosis and lordosis.
8 *Spine*, **7**, 335-342.
- 9 Stokes M, Rankin G, Newham DJ (2005) Ultrasound imaging of lumbar multifidus muscle:
10 normal reference ranges for measurements and practical guidance on the technique.
11 *Manual Therapy*, **10**, 116-126.
- 12 Stokes MJ, Cooper RG, Morris G (1992) Selective changes in multifidus dimensions in patients
13 with chronic low back pain. *European Spine Journal*, **1**, 38-42.
- 14 Takeda N, Kobayashi T, Atsuta Y, Matsuno T, Shirado O, Minami A (2009) Changes in the
15 sagittal spinal alignment of the elderly without vertebral fractures: a minimum 10-year
16 longitudinal study. *Journal of Orthopaedic Science*, **14**, 748-753.
- 17 Wallwork TL, Stanton WR, Freke M, Hides JA (2009) The effect of chronic low back pain on
18 size and contraction of the lumbar multifidus muscle. *Manual Therapy*, **14**, 496-500.
- 19 Ward SR, Kim CW, Eng CM, et al. (2009) Architectural analysis and intraoperative
20 measurements demonstrate the unique design of the multifidus muscle for lumbar spine
21 stability. *Journal of Bone and Joint Surgery*, **91**, 176-185.
- 22 Wood KB, Kos P, Schendel M, Persson K (1996) Effect of patient position on the sagittal-plane
23 profile of the thoracolumbar spine. *Journal of Spinal Disorders*, **9**, 165-169.
- 24
- 25

Lumbar lordosis and extensor muscle size

- 1 Table 1. Age, height, mass, lower lumbar curvature (θ_{L4S1}) and muscle volume for all 42
 2 subjects. Normally distributed values reported as a mean (standard deviation) and range; non-
 3 normally distributed values as a median [inter-quartile range] and range.

	Back pain status				ANOVA
	All subjects	None	Previous	Current	
	(n = 42)	(n = 11)	(n = 26)	(n = 5)	
Age (years)	50 (13) 20 – 75	44 (11) 23 - 59	51(13) 20 – 75	53(12) 36 – 69	F = 1.5
Height (cm)	165 (7) 151 - 178	167 (6) 158 – 177	164 (7) 151 – 178	163 (4) 157 – 167	F = 0.8
Mass (kg)	68 [18] 52 - 128	73 (18) 57 – 106	70 (13) 52 – 107	76 (30) 55 – 128	F = 0.1
θ_{L4S1} (degrees)	39 (4) 28 – 47	40 (3) 35 – 44	38 (4) 28 – 47	39 (4) 33 – 43	F = 1.1
Volume (cm ³)	281 (49) 169 - 358	303 (34) 237 - 348	281 (48) 195 - 358	230 (57) 169 - 299	F = 4.4 **

- 4 Statistical significance is noted as * $p < 0.05$ and ** $p < 0.01$.

5

Lumbar lordosis and extensor muscle size

1 Table 2. Correlation between age, height, mass, lower lumbar curvature (θ_{L4S1}), and muscle
 2 volume. The Pearson correlation coefficients (**R**) are given together with the statistical
 3 significance. Only the data for the 37 subjects who did not have current back pain were included
 4 in this analysis.

	Age	Height	T_mass	θ_{L4S1}	Volume
Age	1.00	-0.31	0.14	-0.44 **	-0.53 **
Height		1.00	0.24	0.17	0.38 *
T_mass			1.00	0.25	0.15
θ_{L4S1}				1.00	0.61 **
Volume					1.00

5 Statistical significance is noted as * $p < 0.05$ and ** $p < 0.01$.

6

Lumbar lordosis and extensor muscle size

- 1 Table 3. Predictors of muscle volume determined from stepwise linear regression model. Only
- 2 the data for the 37 subjects who did not have current back pain were included in this analysis.

	Unstandardized coefficients	Standardized coefficients	P
Constant	128.617		0.107
θ_{L4S1}	5.522	0.47	0.002
Age	-1.103	-0.32	0.030

3

Lumbar lordosis and extensor muscle size

1 Figure legends

2 Figure 1. Segmentation of the lumbar extensor muscles at the level of L3/L4.



4

5 Figure 2. Sagittal images showing the curvature of the lumbar spine for two subjects. Subject A

6 aged 47 years: θ_{L4S1} = 47 degrees, muscle volume = 352 cm². Subject B aged 58 years: θ_{L4S1} = 36

7 degrees, muscle volume = 236 cm².



8 **A**

B