

1 **Usefulness of species traits in predicting range shifts**

2 Alba Estrada<sup>1, 2</sup>, Ignacio Morales-Castilla<sup>3</sup>, Paul Caplat<sup>4</sup> and Regan Early\*<sup>5</sup>

3

4 1: CIBIO/InBIO, Universidade de Évora, Portugal

5 2: Department of Integrative Ecology. Estación Biológica de Doñana (EBD-CSIC). Sevilla,

6 Spain

7 3: Department of Biology, McGill University, Montréal, QC, Canada

8 4: Centre for Environmental and Climate Research, Lund University, Sweden

9 5: Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Cornwall, UK.

10 Twitter: @ReganEarly

11

12 \*Corresponding author: Early, R. ([r.early@exeter.ac.uk](mailto:r.early@exeter.ac.uk)).

13

14

## 15 **Abstract**

16 Information on species' ecological traits might improve predictions of climate-driven range  
17 shifts. However, the usefulness of traits is usually assumed rather than quantified. We  
18 present a framework to identify the most informative traits, based on four key range-shift  
19 processes: (i) emigration of individuals or propagules away from the natal location, (ii) the  
20 distance a species can move, (iii) establishment of self-sustaining populations, and (iv)  
21 proliferation following establishment. We propose a framework that categorises traits  
22 according to their contribution to range-shift processes. We demonstrate how the  
23 framework enables the predictive value of traits to be evaluated empirically, how this  
24 categorisation can be used to better understand range shift processes, and illustrate how  
25 range shift estimates can be improved.

26

## 27 **Predictive traits as a new paradigm in climate change ecology**

28 Mitigating the threat from climate change to biodiversity and ecosystems requires a robust  
29 understanding of how species will respond to new climatic conditions. The most common  
30 method for estimating a species' exposure to climate change is to compare future climatic  
31 conditions against the conditions in which a species currently lives [1]. While there is  
32 disagreement about the accuracy of these techniques, they are well explored, and there is  
33 literature on best practice [2, 3]. Species vulnerability to exposure is less well understood. A  
34 major uncertainty is whether species are able to colonise newly climatically suitable areas as  
35 current geographic ranges become unsuitable. Such "range shifts" (see Glossary) could  
36 mitigate threats from climate change.

37 Predictive traits have been suggested as a simple way to improve estimation of species'  
38 range-shift capacities, identifying how well species are likely to cope with climate change [4-  
39 7]. 'Range-shift capacity' could be interpreted as the likelihood of range shifts occurring or  
40 the timescale over which range shifts might occur. The emerging approach is to integrate  
41 information on both a species' exposure to climate change and the traits expected to drive  
42 range-shift capacity, producing a relative metric of species risk [4-6]. Quantitative evidence  
43 is rarely used to inform the choice of predictive traits in these approaches, and trait  
44 selection often relies on expert opinion and data availability [4, 6]. There is no consensus as  
45 to the traits that should be considered in range shift forecasts, meaning that studies are not  
46 comparable, and that trait data could obscure rather than clarify climate change threats.  
47 Consequently, there is a timely need to quantify how different traits contribute to species  
48 range-shifts, and to identify potentially informative traits for which we do not have  
49 sufficient information.

#### 50 **A framework to evaluate and employ predictive traits**

51 The framework we propose permits the use of existing evidence bases to identify the most  
52 important traits and range shift processes for a given taxa, improved testing of the  
53 relationship between traits and range shifts, and superior assessments of the range-shift  
54 capacities of large numbers of species. Given the hundreds of traits that could be analysed  
55 for different taxa, this framework would permit future studies to be comparable. And while  
56 the evidence bases we discuss are correlative, choosing traits to analyse based on our  
57 framework would generate testable hypotheses as to causal mechanisms.

58 In addition to climate change ecology, bodies of theory within metapopulation, invasion,  
59 life-history, restoration and reintroduction ecology deal with range shifts, i.e. the

60 establishment and expansion of new populations. We draw from these disciplines to  
61 identify four key range shift processes (Box 1). Despite widespread acceptance of these  
62 range shift processes, their importance for any given taxa or range-shift scenario is poorly  
63 known. Traits could be used to indicate success at each range shift process, but the relevant  
64 traits are numerous and diverse, will differ between taxa, and some are poorly quantified  
65 for many species. This presents difficulties for evaluating the importance of a given trait. We  
66 therefore propose a trait categorisation (Box 2) that corresponds to range shift processes.  
67 This categorisation results in testable hypotheses as to causal mechanisms underlying the  
68 relationship between traits and range shifts. The categorisation also allows trade-offs and  
69 interactions between traits to be recognised and accounted for. For example, migratory  
70 status could affect range-shift capacity positively by conferring high movement ability, or  
71 negatively by limiting emigration because migrants show fidelity to breeding and over-  
72 wintering sites between years [8, 9, Box 2]. Indeed species with migratory ability appear to  
73 have low range-shift potential in [10], possibly because migrants show site fidelity (Table 1).  
74 We demonstrate how the framework could be applied to improve range-shift predictions in  
75 box 3.

76 In addition to traits, various range shift stages might be affected by species' exposures to  
77 climate change and thus by species' climatic tolerances. Here we deal exclusively with how  
78 traits can be employed to improve range-shift predictions, but raise the aforementioned  
79 issues in box 4 (Outstanding Questions).

#### 80 **Evidence bases for the relevance of predictive traits to range-shift capacities**

81 There are multiple metrics for which a wealth of data exists that can be used to evaluate the  
82 predictive value of traits. Traits that correlate with biogeographical patterns and their

83 changes through time are likely to be the same traits that correspond to the processes of  
84 climate-driven range shifts. Below we discuss potential metrics for evaluating predictive  
85 traits, summarize their advantages and disadvantages and the major lessons learned for the  
86 use of predictive traits in estimating range-shift potential. We illustrate these points with  
87 selected examples from the literature (Table 1). We focus on terrestrial systems, for which  
88 we found the most evidence as to the processes that drive range shifts.

### 89 *Recent range shift*

90 Detecting the traits shared by species that have undergone the greatest range shifts in  
91 recent decades is the most direct way to infer traits that will promote or hinder modern  
92 range shifts (which must also take place in tens to hundreds of years due to the current pace  
93 of climate change). However, analyses of traits that correspond to recent range shifts have  
94 yielded equivocal results [11-15] (Table 1). One explanation might be that the drivers of  
95 range shift are so complex that a few decades worth of data are insufficient to draw  
96 generalisations. A second explanation is that these analyses have rarely considered species'  
97 exposure to climate change. If no areas become newly climatically suitable then no  
98 colonisation can occur, and if species can tolerate new conditions then no range contraction  
99 should occur. It has often been difficult to calculate exposure due to challenges in obtaining  
100 accurate data on climatic tolerances for the species that have undergone range shifts  
101 (though see [15]). Where this is not possible, measuring shifts along climatic gradients  
102 should yield more insights than using only latitudinal or altitudinal shifts [16]. The  
103 requirement for data at multiple time points restricts measurement of recent range shifts to  
104 a handful of very well-studied taxa and geographic regions, reinforcing biases that already  
105 exist in the climate change literature [17]. Despite these limitations, the few trends that

106 have emerged correspond to results obtained using other biogeographical metrics (range  
107 size or range filling, see below, Glossary and Table 1). We therefore suggest that, while  
108 monitoring ongoing range shifts is highly important for understanding the predictive value  
109 of traits, proxies such as range filling and range size are also useful.

#### 110 *Range size*

111 Widespread species with large ranges have colonised and maintained persistent populations  
112 over large areas [18], and should therefore be able to colonise and proliferate in newly  
113 climatically-suitable areas. This is particularly the case in regions where species ranges are  
114 substantially affected by historic climate change. For instance, in Europe, species' current  
115 ranges are largely determined by the degree to which species have been able to expand  
116 since the Last Glacial Maximum [19]. The traits that correlate with the range size of  
117 European species are therefore expected to be traits that have facilitated post-glacial range  
118 expansion [20]. Traits that correlate with Extent of Occupancy (EOO) would indicate the  
119 furthest distance at which new populations might be established, whereas traits that  
120 correlate with Area of Occupancy (AOO) would indicate the distance at which most  
121 populations will be established. A major caveat is the difficulty in distinguishing the degree  
122 to which range size is determined by species' capacities to colonise and persist in suitable  
123 areas, or by the availability of suitable environmental conditions. Range sizes could also be  
124 affected by historical biogeographical processes, and analytical techniques are emerging to  
125 quantify these effects [21]. These caveats also apply to range filling, below.

#### 126 *Range filling*

127 The rationale behind the use of range filling to inform predictive traits is that, when  
128 potential range is calculated using suitable climatic conditions (using an SDM or

129 physiological data [2, 3, 15]) unoccupied portions of the potential range must be due to non-  
130 climatic range limitations (e.g. edaphic conditions, dispersal limitations, biotic interactions).  
131 Traits associated with range filling would therefore indicate vulnerability to historic non-  
132 climatic range limitations [22, 23], and thus inform species' vulnerability to similar  
133 limitations during modern range shifts [20]. However, range-filling reflects processes that  
134 occurred over thousands of years, some of which might not be at play in modern range-  
135 shifts, due to the faster pace of current climate change. For example, biotic interactions like  
136 competitive exclusion are often more observable over long timescales [24, 25]. Therefore,  
137 traits that correspond to competitive ability might be more important to range filling than to  
138 modern range shifts; a hypothesis that could be tested using available evidence bases and  
139 our framework.

#### 140 *Naturalisation potential*

141 Predictive traits are widely investigated as indicators of species' potential to naturalise,  
142 spread, and impact ecosystems following introductions by humans [26-28]. Similar  
143 ecological and evolutionary processes occur during climate change in the native region and  
144 during naturalisation, as species encounter novel climates and biotic communities [29].  
145 Most of the world's many thousands of human-mediated naturalisations have occurred  
146 within the last 200 years, many within the last 50-100 years [30], i.e. a similar timescale to  
147 that over which the impacts of climate change on native species are normally considered  
148 [30]. However, many studies of predictive traits in naturalisation involve species classified as  
149 invasive [31]. Few introduced species are thought to become invasive (though estimates  
150 vary from, e.g. 1% [32] to 25% [33]). Thus, invasive species may not be representative of  
151 most species, in particular species that are rare as natives and thus of particular concern

152 under climate change. Indeed, when traits related to invasion success are inverted they do  
153 not indicate extinction risk [34]. However, the vast majority of naturalised species do not  
154 become problematic or widespread invasives and are thus broadly representative,  
155 suggesting that tapping this information source would be valuable.

#### 156 *Abundance shifts*

157 Examining changes in abundance across geographic space offers insights into population  
158 growth rate that cannot be gained by studying range (i.e. occurrence) shifts alone [16, 17].  
159 Abundance changes might occur before range shifts can be observed, and could give specific  
160 insight into the traits associated with establishment. The relationship between predictive  
161 traits and trends in abundance have been fairly well studied [35-37]. Traits related to  
162 abundance declines suggest susceptibility to anthropogenic stressors, which could in turn  
163 limit range-shift capacity, particularly in human dominated areas [38]. The heavy data  
164 demands for calculating population trends limit these data to a few countries and for  
165 conspicuous groups of species (Table 1). Recent research has investigated spatial abundance  
166 dynamics at the community level rather than focusing on individual species [e.g. 39]. This is  
167 particularly useful when individual species are rare or show only small abundance changes.  
168 However, community indices might mask hidden drivers or differences between species  
169 [40], and to our knowledge no study has yet linked them to predictive traits.

#### 170 *Threat status*

171 Examining the traits that correspond to species' threat status is commonly done thanks to  
172 large, standardised datasets such as the IUCN Red List [41]. However, threat status is driven  
173 by a plethora of environmental stressors, of which climate change is a relatively recent  
174 factor. We consider threat status to be a weak proxy for species' range-shift capacities



175 under climate change. Nonetheless this line of enquiry has generated much expertise and  
176 understanding of the relationship between traits and species vulnerability.

177

### 178 **Traits explaining distribution changes: what is done, and what could be done**

179 Table 1 summarises 40 studies that are excellent examples of the above evidence bases  
180 because they analysed a large number of species from different taxa and a variety of traits  
181 from different regions. The studies were selected following criteria in Appendix S1 (section  
182 A), and include all metrics discussed above. There are two major hurdles to interpreting  
183 existing evidence bases. First, the suite of traits analysed rarely represent all processes that  
184 could drive either the response variable (metric of trait predictive value) or climate-driven  
185 range shifts (Box 1). This has limited the mechanistic insight that can be obtained, as links  
186 between traits and the response variable might be dependent on trade-offs and interactions  
187 with unmeasured traits [9]. Second, traits employed by different studies are not  
188 standardized between taxa, and often not within taxa [11]. The literature therefore does not  
189 yet permit a meta-analysis, but it is possible to identify high quality studies with broad  
190 taxonomic representation from which we can assess evidence for the importance of trait  
191 categories (Box 2). See box 2 and table for traits included in each category.

192 The taxa for which most trait-based analyses have been undertaken are plants, terrestrial  
193 vertebrate endotherms, and butterflies (Table 1). Geographic areas analysed are mainly in  
194 Europe and North America. Notably, fewer than half of the studies specify hypotheses for  
195 the ecological processes that could underlie the relationship between traits and the  
196 response variable [e. g. 20, 23, 42, 43].

197 Movement ability, ecological generalization and reproductive strategy were the most widely  
198 studied trait categories (Table 1). Movement and ecological generalization were identified  
199 as important more frequently than any other category. However, movement ability had a  
200 significant and positive effect in fewer than half of the cases it was studied. Thus, while the  
201 movement process is clearly a major determinant of range shift capacity, other range-shift  
202 processes also play an important role. Ecological generalization was found to have a positive  
203 effect in the majority of studies that assessed it, whereas reproductive strategy had a  
204 positive effect in a quarter of the cases in which it was studied. The equivocal support for  
205 reproductive strategy might be the result of trade-offs with other traits such as competitive  
206 ability or persistence in unfavourable climatic conditions [44].

207 Persistence in unfavourable climatic conditions is rarely studied, but the proportion of  
208 positive results in Table 1 suggests it should be considered in the future. For instance,  
209 Estrada *et al.* [20] found that seed bank persistence in plants was even more important than  
210 movement ability or ecological generalization in predicting range size and filling for plant  
211 species in Europe. For animals, hibernation has been hypothesised as improving range shifts  
212 by enabling individuals to avoid unfavourable conditions [11] (Box 2), but showed no  
213 predictive power (Table 1). No support was found for longevity (Box 2), possibly because  
214 longevity trades off with age of first reproduction, reflecting slower colonization [44]. Thus,  
215 traits that correspond clearly to persistence might be difficult to define for animals.

216 Although rarely accounted for in range-shift forecasts, competitive ability was studied in 14  
217 cases, and had a positive effect in nine. For plants there are well-established frameworks to  
218 evaluate competitiveness (e.g. the Competitor-Stress tolerator-Ruderal framework [44]) but  
219 there is as yet no corresponding framework for terrestrial animals. For animals therefore,

220 we included traits in this category that correspond to dominance (e.g. local abundance),  
221 intra-generic co-occurring species richness (indicating the number of similar, potential  
222 competitor, species), and brain size (which corresponds to innovation, a key component of  
223 competition avoidance [27]). Given the surprisingly high importance for these traits, we  
224 recommend more work towards understanding the impacts of competition on species  
225 ranges.

226 The least studied and least supported trait categories in our sample were avoidance of small  
227 population effects, which corresponds to colonisation ability, and site (in)fidelity, which  
228 corresponds to emigration (Box 2). Categorising avoidance of small population effects in  
229 plants is fairly straightforward: self-fertilization and vegetative regeneration are key traits in  
230 this respect and are widely measured. However, in the papers we sampled for animals,  
231 relevant traits analysed were population or social group size, which could be too simplistic  
232 to capture complex outcomes of animal behaviour for small population sizes. We note that  
233 traits that correspond to reproductive strategies, which were important in some cases,  
234 could also contribute to avoidance of small population effects. The limited support for traits  
235 related to avoidance of small population effects might imply that the colonisation process  
236 does not strongly limit range shifts for the majority of taxa studied (Box 1). However, we  
237 suggest that the importance of the avoidance of small population effects should not be  
238 precluded until a broader range of relevant traits is examined. With respect to site  
239 (in)fidelity, although there is little evidence for the predictive value of relevant traits in Table  
240 1 modelling studies support the importance of emigration to climate-driven range shifts  
241 [45]. We therefore suggest that site (in)fidelity traits merit further investigation.

#### 242 **Limitations of existing evidence bases**

243 A lack of support in the evidence bases we sampled should not be taken to suggest that a  
244 given trait is not important for any species. Rather, positive or negative effects in Table 1  
245 indicate evidence across many species that a given trait is sufficiently tightly linked to a  
246 range-shift process that it could be used to inform relative estimates of range-shift  
247 likelihood in multi-species analyses. A caveat to all correlative approaches we outline is that  
248 trait plasticity and evolution (past or ongoing) might obscure the relevance of traits to range  
249 shifts.

250 While trait data are becoming increasingly available, potentially important traits are often  
251 unavailable, e.g. movement distance in animals. Alternatives are to use values for  
252 congeneric species, or morphological proxies [9, 14, 46]. Increasing awareness of the  
253 usefulness of traits (particularly non-movement traits) to inform responses to climate  
254 change should stimulate collection and curation of potentially informative traits.

255 Macro-ecological analyses of predictive traits typically use a single trait value for the whole  
256 species (usually the mean of all recorded values) [14, 20], despite increasing evidence of  
257 substantial intraspecific variation [47, box 4: Outstanding Questions]. Whenever possible,  
258 the use of trait variance together with the mean can provide new insights and more  
259 accurate separation of species' range shift capacities. However, information on intraspecific  
260 variation is rarely available, so it is not yet possible to develop a comprehensive macro-  
261 ecological analysis for a large number of traits. Nevertheless, the findings in Table 1 indicate  
262 that characterizing species with single trait values does detect relative interspecific variation  
263 in metrics that correspond to range-shift capacity.

264 **Concluding remarks**

265 We encourage the use of predictive traits in assessment of species potential to colonise new  
266 areas in response to climate change. We recommend that all range-shift processes should  
267 be represented in the choice of traits, and we demonstrate how traits can be categorised in  
268 terms of their contribution to these processes and used to inform range shift potential.  
269 While correlative methods are still the main option for assessing predictive traits for large  
270 numbers of species, employing our framework would formalise hypotheses on the  
271 mechanistic underpinning of the relationship between trait and range shift. This would  
272 permit synthesis across studies and meta-analysis, ultimately leading to a broad  
273 understanding of how traits mediate range shift. While we encourage new analyses to  
274 improve understanding for different global regions, taxonomic groups, and under-studied  
275 range-shift processes, there is already sufficient evidence that traits corresponding to  
276 movement, ecological generalization, persistence in unfavourable climatic conditions,  
277 reproductive strategy, and competitive ability should be considered for inclusion in range-  
278 shift evaluations. Further experimental studies could use our proposed framework to better  
279 make mechanistic linkages between traits and range shifts and site persistence.

280

## 281 **Acknowledgments**

282 This research is funded by the ERA-Net BiodivERSA, with the national funder FCT, through  
283 the project BIODIVERSA/0003/2011. AE was funded by the project  
284 IF/00266/2013/CP1168/CT0001. IM-C is funded by the Fonds de Recherches du Québec -  
285 Nature et Technologies (FQRNT) programme. PC is funded by the Swedish strategic research  
286 environment Biodiversity and Ecosystem Services in a Changing Climate (BECC).

287

288 **Boxes**

289

290 **Glossary**

291 **AOO:** area of occurrence, the geographic area that is actually occupied by a species [48],  
292 often defined as the number of occupied grid cells (which vary in size between atlases) [20].

293 **Competitive ability:** the ability of an individual of one species to reduce the availability of  
294 contested resources to an individual from another species, and to tolerate or avoid  
295 reduction in contested resource availability by an individual from another species [49].

296 **Ecological generalization:** the ability to use a wide variety of a given resource type, e.g.,  
297 ecological generalists could breed in a wide variety of land cover types, have a broad diet, or  
298 tolerate a broad range of soil types.

299 **Emigration:** first range-shift process, in which an individual embarks on a journey  
300 (movement) outside its natal location.

301 **EOO:** extent of occurrence, the area within the outer limits of the species geographic  
302 distribution [43, 48].

303 **Establishment:** range-shift process following movement, in which one or more individuals  
304 reproduce and found a self-sustaining population.

305 **Indicative traits:** species' characteristics related to environmental tolerance, habitat  
306 specialization, geographical boundaries or spatial distribution [5, 7, 50]. These traits can be  
307 measured at the individual or population level, so they are not life-history traits in strict  
308 sense.

309 **Life-history traits:** those morphological, physiological or phenological characteristics  
310 measurable at the individual level that have an effect on individual performance [51].

311 **Movement ability:** ability of an individual or propagule to travel outside its natal location.  
312 This ability is often represented as the average or upper end of the distance moved in the  
313 lifetime of an individual or propagule. Note that this can be informed by, but is not  
314 restricted to, natal dispersal distances (movement from natal to breeding site). We  
315 specifically use 'movement ability' instead of 'dispersal ability' to avoid confusion, as the  
316 latter term is widely used to include emigration, movement, and establishment [52]. The  
317 movement process in dispersal has also been called transience, transport, and transfer [52].

318 **Persistence in unfavourable climatic conditions:** a population's capacity to survive during  
319 periods in which poor climate conditions leads to zero or negative population growth rate.  
320 Unfavourability could occur through climate change directly, or as a result of climatically  
321 induced changes in other elements of habitat suitability.

322 **Potential range:** the geographic area in which environmental conditions are suitable for a  
323 given species, even if the species is present or not. Potential range is often calculated using  
324 SDMs.

325 **Predictive traits:** we define predictive traits as any species' feature that can be used to  
326 predict (a) the likelihood and extent of range shift given exposure to climate change, or (b)  
327 the species' interactions with other species and non-climatic elements of the environment,  
328 which directly or indirectly affect range shifts. Predictive traits might include 'intrinsic'  
329 ecophysiological, life-history and demographic traits [51], as well as broader, non-  
330 organismal 'indicative' traits such as habitat requirements or spatial distribution [5, 7, 11].

331 **Proliferation:** fourth range-shift process in which established populations become more  
332 than self-sustaining, producing individuals that will in turn disperse and cause further  
333 population spread.

334 **Range filling:** The proportion of its potential range that a species occupies.

335 **Range shift:** expansion of one part of the range margin following colonisation events. Range  
336 shift may or may not be accompanied by a contraction in another part of the range margin.

337 **Reproductive strategy:** the number, timing and degree of investment in each reproductive  
338 event, which are related to demography, fecundity and speed of life history. Species with an  
339 'r' strategy reproduce early, have small body mass, and many offspring per year. Species  
340 with a 'K' strategy are older at first reproduction, have larger body mass and fewer  
341 offspring.

342 **SDM:** species distribution model. SDMs relate a species distribution with the environmental  
343 conditions in which species are found, in order to calculate environmentally suitable areas  
344 for that species.

345 **Site (in)idelity:** reflects the likelihood that an individual will embark on a dispersal event to  
346 emigrate away from the natal patch. High site fidelity corresponds to a low likelihood of  
347 emigration and thus low range-shift capacity. We therefore use 'site (in)idelity' in line with  
348 the other six trait categories, for which the terms correspond to a positive effect on range-  
349 shift capacity.

350 **Small population effects:** factors that make it difficult for small populations to grow, and  
351 thus hinder population establishment. These include Allee effects, genetic drift, and  
352 susceptibility to demographic or environmental stochasticity.



353

354 **Box 1. Range-shift processes.**

355 The first range-shift process (see Figure 1) is that individuals embark on a journey away from  
356 their natal location (**emigration**). For animals the motivations involved include high  
357 population density, low resource availability, and harassment from prospective mates [45].  
358 For sessile taxa, such as plants, some aspects of reproductive behaviour can affect the  
359 probability that propagules disperse away from the adults' location, for example timing of  
360 seed shed to maximise dispersal by animals [53]. Species that have a physical or behavioural  
361 mechanism that promotes emigration are more likely to respond to poor environmental  
362 conditions (as driven by climate change) by leaving the natal location than those that do not  
363 have these mechanisms. Recent applications of metapopulation modelling approaches to  
364 climate change highlight the importance of emigration for range shifts [45, 54]. **Movement**  
365 itself, i.e. the transfer of individuals or propagules away from the location in which they  
366 originated [54], is the second and most widely studied range-shift process. The upper limit  
367 of a species' dispersal distance is one of the strongest limitations on metapopulation  
368 persistence, invasive spread, and population recovery [55-57]. The third range-shift process  
369 is **establishment**, i.e. the ability of dispersing individuals to reproduce and found new  
370 populations following a dispersal event. While probability of establishment is affected by  
371 the number of arriving propagules at a site, dispersal, invasion and reintroduction ecology  
372 demonstrate that non-movement traits and their interaction with local conditions also  
373 mediate ease of establishment [58]. The fourth range-shift process is **proliferation**, i.e. the  
374 growth of established populations to become more than self-sustaining, producing  
375 individuals that will in turn disperse and cause further population spread [59]. In the short

376 term, spreading outwards to locations immediately surrounding the newly colonised  
377 location will improve population robustness. In the longer term, a large number of  
378 emigrating individuals will sustain the range-shift itself. Life-histories that permit rapid  
379 proliferation contribute greatly to the long term success of naturalised and reintroduced  
380 populations [47]. Predictive traits employed in climate-change risk assessments should  
381 correspond to all of the above processes.

## 382 **Figure I**

383 Figure I. Schematic representation of range-shift processes and their relationship with the  
384 trait categories we suggest (Box 2). We do not suggest that there is an exclusive  
385 correspondence between a given trait category and range-shift process. Rather, we identify  
386 the links between trait categories and range-shift process that evidence suggests are the  
387 most directly informative. Note that we are considering species traits and not their  
388 interaction with the environment, e.g. we do not include the effect that climate-driven  
389 resource limitation could have on emigration.

390

## 391 **Box 2. Categorisation of predictive traits**

392 We propose seven trait categories related to range-shift processes (Figure I). Trade-offs and  
393 interactions between traits mean that some categories cannot be tied exclusively to one  
394 range-shift process. We demonstrate how traits addressed by studies in Table 1 could be  
395 categorised. First, **site (in)fidelity** corresponds to emigration, and can be informed by  
396 migration (migrants show fidelity to breeding and over-wintering sites between years [8]),  
397 and breeding behaviour (social or territorial) that encourages individuals to remain at natal

398 locations or to disperse to new breeding territories [60]. Sessile taxa can also have traits  
399 that correspond to site (in)fidelity, for example plants can time seed shed to maximise  
400 dispersal by animals [53]. Second, **movement ability** corresponds directly to the movement  
401 process, and for plants can be informed by properties related to dispersal vectors [20]. In  
402 animals, movement can be informed by natal dispersal [14] and by behaviours such as  
403 migratory status [61], home ranging area [37], or flight period length [62], and morphologies  
404 such as wing or leg length [63]. Third, **small population effects** hinder establishment, and  
405 particularly occur in animals that rely on group behaviour for breeding, resource acquisition,  
406 or predator avoidance [41], and in plants that cannot reproduce vegetatively or self-  
407 pollinate [20]. Fourth, **persistence in unfavourable climatic conditions** aids establishment  
408 under climate change because newly colonised locations can fluctuate in climatic suitability  
409 before becoming consistently suitable for a given species [64, 65]. This category can be  
410 informed in plants by seed-bank persistence [20], and in animals by 'sleep' (e.g. hibernation,  
411 dormancy) and 'hide' (use of burrows, caves, tree-holes) behaviours that might allow  
412 species to 'wait out' unfavourable periods [66] or by traits indicating the importance of each  
413 reproductive event to lifetime reproduction (e.g. longevity). Fifth, **ecological generalisation**  
414 aids establishment and proliferation by increasing resource availability, and includes diet  
415 breadth, land-cover types occupied, and breadth of diurnal cycle [11]. Sixth, species with an  
416 'r' **reproductive strategy** will rapidly achieve high local abundances (aiding establishment),  
417 thus driving emigration (aiding proliferation). Reproductive strategy can be informed by  
418 traits linked to the number, timing and degree of investment in each reproductive event [11,  
419 27]. Seventh, low **competitive ability** hinders establishment when competition with  
420 incumbent individuals occurs, and hinders proliferation by slowing the growth of already-  
421 established populations. Competitive ability could be informed by population traits (e.g.

422 local density [34]) the number of co-occurring congeners [67]), by brain size in animals [27]  
423 and by leaf-economic traits in plants [44, 68].

424 **Box 3. Case studies applying the predictive trait framework to evaluate range-shift**  
425 **capacities**

426 We examine pairs of species that are predicted to potentially undergo climate-driven range  
427 shift in the 21<sup>st</sup> century (Table I). We show how different traits can be used to inform the  
428 likelihood or speed of range shifts. There are multiple ways in which trait categories could  
429 be combined to determine range-shift ability. Here we classify trait values for each species  
430 as High, Moderate or Low depending on the species' trait value relative to values for related  
431 species. We used only the best-supported traits for a given taxa (Table 1). For the sake of  
432 illustration, we consider that a trait category is supported when at least three studies in  
433 Table 1 found a significant relationship between the trait category and a response variable.  
434 Other approaches are possible, such as i) assigning a numeric score to the results in each  
435 trait category and summing across all, or the best-supported categories, ii) two species can  
436 be compared by summing the number of differences (positive and negative) between  
437 results in each, or the best-supported, trait category. We make no recommendations as to  
438 best practice as insufficient information exists on the relative importance of each trait  
439 category, but urge research that compares the importance of trait categories amongst  
440 species. Both *Populus nigra* and *Carpinus betulus* are predicted to gain climatically suitable  
441 areas to the north of their range (Figure II), but *Carpinus betulus* is more likely to colonise  
442 this area. *Sylvia cantillans* is predicted to gain proportionally more climatically suitable area  
443 than *Corvus monedula* but has less ability to colonise this area, changing relative  
444 assessments of climate change effects on these species.

445 **Table I.** Case studies illustrating the application of our framework to improve range-shift  
 446 evaluations under climate change.

Taxa	Plants (trait data from [20])		Median values for native European trees in [20]
Species	<i>Carpinus betulus</i>	<i>Populus nigra</i>	
Movement ability	High (estimated 1500m, bird and mammal seed dispersal, seed mass 53 mg)	Moderate (estimated 15m, wind seed dispersal, low seed mass 0.81 mg)	Dispersal 500m Seed mass 25mg
Persistence in unfavourable climatic conditions	Moderate (seed bank persistence 1-5 years)	Low (seed bank persistence < 1 year)	< 1 year
Ecological generalisation	High (2.19 vegetation types occupied)	Low (1.2 vegetation types occupied)	1.86 vegetation types
Reproductive strategy	High (perennial tree species, age of first flowering 15 years)	Moderate (perennial tree species, age of first flowering 6 years)	8 years
Competitive ability	High/moderate (mean height 15m, SLA 24 mm <sup>2</sup> /mg)	High/moderate (mean height 50m, SLA 11 mm <sup>2</sup> /mg)	Height 20 m SLA 10 mm <sup>2</sup> /mg
<b>Resulting range-shift ability</b>	<b>High</b>	<b>Moderate / Low</b>	
Taxa	Birds (trait data from [69])		Median values for native European passerines in [69]
Species	<i>Corvus monedula</i>	<i>Sylvia cantillans</i>	
Movement ability	High/Moderate (natal dispersal 8.6km, seasonal migrant)	Moderate (seasonal migrant)	8.45 km
Ecological generalisation	High (omnivorous and opportunistic, high habitat breadth 5.4)	Low (omnivorous, low habitat breadth 1.25)	3.4 vegetation types
Reproductive strategy	Low (large-bodied: 248g, late reproducing: year 2, average clutch size 4.46, 1 brood per year)	Moderate (small-body: 9.8 g, early reproducing: year 1, small clutches: 3.86, 2 broods per year)	Body mass 21 g Sexual maturity year 1 Clutch size 4.6 Clutches/year 1.7
Competitive ability	High (large brain size : body mass ratio)	Moderate (small brain size : body mass ratio)	
<b>Resulting range-shift ability</b>	<b>High/Moderate</b>	<b>Moderate/Low</b>	

447  
 448 **Figure II.** Current distributions and areas predicted climatically suitable for case study  
 449 species in 2071-2100 (see section C in Appendix S1 for details of distributions and forecasts).

450 Black dots are currently occupied, blue areas are climatically suitable in the future, and  
451 yellow areas are climatically unsuitable.

452 **References**

- 453 1 Garcia, R.A. *et al.* (2014) Multiple dimensions of climate change and their implications for  
454 biodiversity. *Science (New York, N.Y.)* 344, 1247579-1247579
- 455 2 Anderson, R.P. (2013) A framework for using niche models to estimate impacts of climate  
456 change on species distributions. *Annals of the New York Academy of Sciences* 1297, 8-  
457 28
- 458 3 Márquez, A.L. *et al.* (2011) Combining climate with other influential factors for modelling the  
459 impact of climate change on species distribution. *Clim. Chang.* 108, 135-157
- 460 4 Foden, W.B. *et al.* (2013) Identifying the world's most climate change vulnerable species: a  
461 systematic trait-based assessment of all birds, amphibians and corals. *PLoS ONE* 8,  
462 e65427
- 463 5 Triviño, M. *et al.* (2013) Risk assessment for Iberian birds under global change. *Biol. Conserv.*  
464 168, 192-200
- 465 6 Garcia, R.A. *et al.* (2014) Matching species traits to projected threats and opportunities from  
466 climate change. *J. Biogeogr.* 41, 724-735
- 467 7 Pearson, R. *et al.* (2014) Life history and spatial traits predict extinction risk due to climate  
468 change. *Nat. Clim. Change* 4, 217-221
- 469 8 Bensch, S. (1999) Is the range size of migratory birds constrained by their migratory  
470 program? *J. Biogeogr.* 26, 1225-1235
- 471 9 Laube, I. *et al.* (2013) Towards a more mechanistic understanding of traits and range sizes.  
472 *Global Ecol. Biogeogr.* 22, 233-241
- 473 10 Forsyth, D.M. *et al.* (2004) Climatic suitability, life-history traits, introduction effort, and the  
474 establishment and spread of introduced mammals in Australia. *Conserv. Biol.* 18, 557-  
475 569
- 476 11 Angert, A.L. *et al.* (2011) Do species' traits predict recent shifts at expanding range edges?  
477 *Ecol. Lett.* 14, 677-689

- 478 12 Mattila, N. *et al.* (2011) Ecological correlates of distribution change and range shift in  
479 butterflies. *Insect Conserv. Diver.* 4, 239-246
- 480 13 Auer, S.K. and King, D.I. (2014) Ecological and life-history traits explain recent boundary  
481 shifts in elevation and latitude of western North American songbirds. *Global Ecol.*  
482 *Biogeogr.* 23, 867-875
- 483 14 Bradshaw, C.J.A. *et al.* (2014) Predictors of contraction and expansion of area of occupancy  
484 for British birds. *P. Roy. Soc. B-Biol. Sci.* 281, 20140744
- 485 15 Sunday, J.M. *et al.* (2015) Species traits and climate velocity explain geographic range shifts  
486 in an ocean warming hotspot. *Ecol. Lett.* 18, 944-953
- 487 16 Tayleur, C. *et al.* (2015) Swedish birds are tracking temperature but not rainfall: evidence  
488 from a decade of abundance changes. *Global Ecol. Biogeogr.* 24, 859-872
- 489 17 Lenoir, J. and Svenning, J.C. (2015) Climate-related range shifts – a global multidimensional  
490 synthesis and new research directions. *Ecography* 38, 15-28
- 491 18 Kunin, W.E. and Gaston, K.J. (1997) *The biology of rarity: causes and consequences of rare-*  
492 *common differences.* Springer
- 493 19 Svenning, J.-C. *et al.* (2008) Postglacial dispersal limitation of widespread forest plant  
494 species in nemoral Europe. *Ecography* 31, 316-326
- 495 20 Estrada, A. *et al.* (2015) Species' intrinsic traits inform their range limitations and  
496 vulnerability under environmental change. *Global Ecol. Biogeogr.* 24, 849–858
- 497 21 Warren, D.L. *et al.* (2014) Mistaking geography for biology: inferring processes from species  
498 distributions. *Trends Ecol. Evol.* 29, 572-580
- 499 22 Svenning, J.-C. and Skov, F. (2004) Limited filling of the potential range in European tree  
500 species. *Ecol. Lett.* 7, 565-573
- 501 23 Dullinger, S. *et al.* (2012) Post-glacial migration lag restricts range filling of plants in the  
502 European Alps. *Global Ecol. Biogeogr.* 21, 829-840



503 24 Hortal, J. *et al.* (2011) Ice age climate, evolutionary constraints and diversity patterns of  
504 European dung beetles. *Ecol. Lett.* 14, 741-748

505 25 Wisz, M.S. *et al.* (2013) The role of biotic interactions in shaping distributions and realised  
506 assemblages of species: implications for species distribution modelling. *Biol. Rev.* 88,  
507 15-30

508 26 Bucharova, A. and Van Kleunen, M. (2009) Introduction history and species characteristics  
509 partly explain naturalization success of North American woody species in Europe. *J.*  
510 *Ecol.* 97, 230-238

511 27 Sol, D. *et al.* (2012) Unraveling the life history of successful invaders. *Science* 337, 580-583

512 28 Caplat, P. *et al.* (2013) Movement, impacts and management of plant distributions in  
513 response to climate change: insights from invasions. *Oikos* 122, 1265-1274

514 29 Moran, E.V. and Alexander, J.M. (2014) Evolutionary responses to global change: lessons  
515 from invasive species. *Ecol. Lett.* 17, 637-649

516 30 Mack, R.N. *et al.* (2000) Biotic invasions: Causes, epidemiology, global consequences, and  
517 control. *Ecol. Appl.* 10, 689–710

518 31 Richardson, D.M. and Pysek, P. (2012) Naturalization of introduced plants: ecological drivers  
519 of biogeographical patterns. *New Phytol.* 196, 383-396

520 32 Williamson, M. (1996) *Biological Invasions*. Chapman & Hall

521 33 Jeschke, J.M. and Strayer, D.L. (2005) Invasion success of vertebrates in Europe and North  
522 America. *PNAS* 102, 7198-7202

523 34 Blackburn, T.M. and Jeschke, J.M. (2009) Invasion success and threat status: two sides of a  
524 different coin? *Ecography* 32, 83-88

525 35 Jiguet, F. *et al.* (2007) Climate envelope, life history traits and the resilience of birds facing  
526 global change. *Global Change Biol.* 13, 1672-1684

527 36 Seoane, J. and Carrascal, L.M. (2008) Interspecific differences in population trends of  
528 Spanish birds are related to habitat and climatic preferences. *Global Ecol. Biogeogr.* 17,  
529 111-121

530 37 Collen, B. *et al.* (2011) Predicting how populations decline to extinction. *Phil. Trans. R. Soc.*  
531 *B* 366, 2577-2586

532 38 Gill, J.A. *et al.* (2001) The buffer effect and large-scale population regulation in migratory  
533 birds. *Nature* 412, 436-438

534 39 Devictor, V. *et al.* (2008) Birds are tracking climate warming, but not fast enough. *P. Roy.*  
535 *Soc. B-Biol. Sci.* 275, 2743-2748

536 40 Clavero, M. *et al.* (2011) Climate change or land use dynamics: Do we know what climate  
537 change indicators indicate? *PLoS ONE* 6, e18581

538 41 Cardillo, M. *et al.* (2008) The predictability of extinction: biological and external correlates  
539 of decline in mammals. *P. Roy. Soc. B-Biol. Sci.* 275, 1441-1448

540 42 Schurr, F.M. *et al.* (2007) Colonization and persistence ability explain the extent to which  
541 plant species fill their potential range. *Global Ecol. Biogeogr.* 16, 449-459

542 43 Van der Veken, S. *et al.* (2007) Life-history traits are correlated with geographical  
543 distribution patterns of western European forest herb species. *J. Biogeogr.* 34, 1723-  
544 1735

545 44 Grime, J.P. (1977) Evidence for the existence of three primary strategies in plants and its  
546 relevance to ecological and evolutionary theory. *Am. Nat.* 111, 1169-1194

547 45 Altwegg, R. *et al.* (2013) Density-dependent dispersal and the speed of range expansions.  
548 *Divers. Distrib.* 19, 60-68

549 46 Catullo, R.A. *et al.* (2015) Extending spatial modelling of climate change responses beyond  
550 the realized niche: estimating, and accommodating, physiological limits and adaptive  
551 evolution. *Global Ecol. Biogeogr.* 24, 1192-1202

552 47 Jongejans, E. *et al.* (2011) Importance of individual and environmental variation for invasive  
553 species spread: a spatial integral projection model. *Ecology* 92, 86-97

554 48 Gaston, K.J. and Fuller, R.A. (2009) The sizes of species' geographic ranges. *J. Appl. Ecol.* 46,  
555 1-9

556 49 Aarssen, L. (1984) On the distinction between niche and competitive ability: Implications for  
557 coexistence theory. *Acta Biotheor* 33, 67-83

558 50 Bartomeus, I. *et al.* (2013) Historical changes in northeastern US bee pollinators related to  
559 shared ecological traits. *PNAS* 110, 4656-4660

560 51 Violle, C. *et al.* (2007) Let the concept of trait be functional! *Oikos* 116, 882-892

561 52 Travis, J.M.J. *et al.* (2012) Modelling dispersal: an eco-evolutionary framework  
562 incorporating emigration, movement, settlement behaviour and the multiple costs  
563 involved. *Methods Ecol. Evol.* 3, 628-641

564 53 Warren, R.J. *et al.* (2014) Competition as a mechanism structuring mutualisms. *J. Ecol.* 102,  
565 486-495

566 54 Bocedi, G. *et al.* (2014) Mechanistic modelling of animal dispersal offers new insights into  
567 range expansion dynamics across fragmented landscapes. *Ecography* 37, 1240-1253

568 55 Baguette, M. (2003) Long distance dispersal and landscape occupancy in a metapopulation  
569 of the cranberry fritillary butterfly. *Ecography* 26, 153-160

570 56 Coutts, S. *et al.* (2011) What are the key drivers of spread in invasive plants: dispersal,  
571 demography or landscape: and how can we use this knowledge to aid management?  
572 *Biol. Invasions* 13, 1649-1661

573 57 Brederveld, R.J. *et al.* (2011) Dispersal as a limiting factor in the colonization of restored  
574 mountain streams by plants and macroinvertebrates. *J. Appl. Ecol.* 48, 1241-1250

575 58 Hampe, A. (2011) Plants on the move: The role of seed dispersal and initial population  
576 establishment for climate-driven range expansions. *Acta Oecol.* 37, 666-673

577 59 Blackburn, T.M. *et al.* (2011) A proposed unified framework for biological invasions. *Trends*  
578 *Ecol. Evol.* 26, 333-339

579 60 Møller, A.P. (2006) Sociality, age at first reproduction and senescence: comparative  
580 analyses of birds. *J. Evol. Biol.* 19, 682-689

581 61 Cassey, P. *et al.* (2004) Global patterns of introduction effort and establishment success in  
582 birds. *P. Roy. Soc. B-Biol. Sci.* 271, S405-S408

583 62 Pöyry, J. *et al.* (2009) Species traits explain recent range shifts of Finnish butterflies. *Global*  
584 *Change Biol.* 15, 732-743

585 63 Bohning-Gaese, K. *et al.* (2006) Range size: Disentangling current traits and phylogenetic  
586 and biogeographic factors. *Am. Nat.* 167, 555-567

587 64 Early, R. and Sax, D.F. (2011) Analysis of climate paths reveals potential limitations on  
588 species range shifts. *Ecol. Lett.* 14, 1125-1133

589 65 Jackson, S.T. *et al.* (2009) Ecology and the ratchet of events: climate variability, niche  
590 dimensions, and species distributions. *PNAS* 106, 19685-19692

591 66 Liow, L.H. *et al.* (2009) Lower extinction risk in sleep-or-hide mammals. *Am. Nat.* 173, 264-  
592 272

593 67 Laube, I. *et al.* (2013) Intra-generic species richness and dispersal ability interact to  
594 determine geographic ranges of birds. *Global Ecol. Biogeogr.* 22, 223-232

595 68 Hamilton, M.A. *et al.* (2005) Life-history correlates of plant invasiveness at regional and  
596 continental scales. *Ecol. Lett.* 8, 1066-1074

597 69 Estrada, A. *et al.* (In prep) What can life-history traits tell us about species' range-shift  
598 responses to climate change? A multi-taxon approach.

599 70 Zhu, K. *et al.* (2012) Failure to migrate: lack of tree range expansion in response to climate  
600 change. *Global Change Biol.* 18, 1042-1052

601 71 Grossenbacher, D. *et al.* (2015) Geographic range size is predicted by plant mating system.  
602 *Ecol. Lett.* 18, 706-713

603 72 Gove, A.D. *et al.* (2009) Dispersal traits linked to range size through range location, not  
604 dispersal ability, in Western Australian angiosperms. *Global Ecol. Biogeogr.* 18, 596-  
605 606

606 73 Thompson, K. *et al.* (1999) Range size, dispersal and niche breadth in the herbaceous flora  
607 of central England. *J. Ecol.* 87, 150-155

608 74 Nogués-Bravo, D. *et al.* (2014) Phenotypic correlates of potential range size and range filling  
609 in European trees. *Perspect. Plant Ecol. Evol. Syst.* 16, 219-227

610 75 Walker, K.J. and Preston, C.D. (2006) Ecological predictors of extinction risk in the Flora of  
611 Lowland England, UK. *Biodivers. Conserv.* 15, 1913-1942

612 76 Munguía, M. *et al.* (2008) Dispersal limitation and geographical distributions of mammal  
613 species. *J. Biogeogr.* 35, 1879-1887

614 77 Reif, J. *et al.* (2010) Population changes in Czech passerines are predicted by their life-  
615 history and ecological traits. *Ibis* 152, 610–621

616 78 Hoare, J.M. *et al.* (2012) Can correlated population trends among forest bird species be  
617 predicted by similarity in traits? *Wildlife Research* 39, 469-477

618 79 Kotiaho, J.S. *et al.* (2005) Predicting the risk of extinction from shared ecological  
619 characteristics. *PNAS* 102, 1963-1967

620

621

Predictor trait category

Taxon	Number of species studied	Location	Response variable	Predictor trait category							Reference
				Site (in) fidelity	Movement ability	Avoidance of small population effects	Persistence in unfavourable climatic conditions	Ecological generalization	Reproductive strategy	Competitive ability	
Alpine plants	133	Switzerland	RS		x	x			x		[11] <sup>p</sup>
Forest plants	92	USA	RS		x						*[70]
Plants	734	Global	RZ			+					*[71] <sup>p</sup>
Plants	1276	Europe	RZ		+	x	+	+	-	+	[20] <sup>p</sup>
Angiosperms	524	Australia	RZ		x				x		[72] <sup>p</sup>
Herbaceous flora	263	Central England	RZ		+			+			[73] <sup>p</sup>
Forest plants	273	W Europe	RZ		+		+				[43] <sup>p</sup>
Plants	1276	Europe	RF		+	x	+	+	-	+	[20] <sup>p</sup>
Plants	183	Austrian Alps	RF		+			+		x	*[23] <sup>p</sup>
Plants	37	Fynbos biome	RF		+		+			+	[42] <sup>p</sup>
Trees	48	Europe	RF		+						[74] <sup>p</sup>
Trees	55	Europe	RF						x	x	[22] <sup>p</sup>
Plants	150	E Australia	IS		+					+	[68] <sup>p</sup>
Woody plants	278	N America and Europe	IS		+	x	x	+	-	+	[26]
Plants	898	Lowland England	ER		x			+	x	x	[75]
Mammals	28	NW America	RS				x	x	x		[11] <sup>p</sup>
Mammals	23	Australia	RZ	x	x		-	+	+		[10] <sup>p</sup>
Endemic mammals	89	Mexico	RF		x			x	x		*[76]
Mammals	40	Australia	IS	+	-		x	+	x		[10] <sup>p</sup>
Mammals	292	USA	AC		x			x	x		[37] <sup>p</sup>
Mammals	372	Global	ER		-	x		+	+	+	[41] <sup>p</sup>
Birds	254	N America	RS		x			-	x		[11] <sup>p</sup>
Birds	97	UK	RS		+				+		[14] <sup>p</sup>
Songbirds	40	N America	RS	x	x			-	-		[13] <sup>p</sup>
Sylvia warblers	26	Global	RZ		+				x		[63] <sup>p</sup>
Passerines	165	Global	RZ	-	+			+	+		[9] <sup>p</sup>
Birds	23	Palaeartic <sup>a</sup>	RF		+			x		+	[67] <sup>p</sup>
Birds	1813	N America and Europe	IS				+	+	+	+	[34]
Birds	416	Global	IS	x	x			+	x		[61] <sup>p</sup>
Birds	428	Global	IS	x	x	x	x	+	x	+	[27] <sup>p</sup>
Birds	71	France	AC	x	x			+	+		[35] <sup>p</sup>
Passerines	57	Spain	AC	x	x			+		x	[36]
Passerines	68	Czech Republic	AC	x	x			-	-		[77] <sup>p</sup>
Forest birds	18	New Zealand	AC					x	x		[78]
Herptiles	36	USA	ER		x			+	+	+	[7]
Butterflies	48	Finland	RS		+			x			[62] <sup>p</sup>
Butterflies	95	Finland	RS		+			+			*[12]
Odonata	24	UK	RS	x	x			+	x		[11] <sup>p</sup>
Bees	187	NE USA	AC			x		+			[50] <sup>p</sup>
Butterflies	95	Finland	ER		+			+			*[79]
			Total +	1	16	1	5	20	7	9	
			Total -	1	2	0	1	3	5	0	
			Total x	8	16	7	4	7	13	5	

622

623

624 **Table 1. Selected studies testing relevance of predictive traits to range-shift capacities, and a**  
625 **summary of the results.** An \* before the reference indicates that predictive traits were tested  
626 individually, not in concert with other traits. A 'P' after the reference indicates that the study  
627 accounts for the phylogeny of the species. 'a': the study area of this reference is Palaearctic,  
628 Afrotropics and Indo-Malaya. References are sorted by taxa (plants, mammals, birds, herptiles  
629 and invertebrates) and then by the response variable. Response variables are as follows. RS:  
630 range shift, RZ: range size, RF: range filling, IS: introduction (naturalisation) or invasion success,  
631 AC: abundance change, ER: extinction risk. + indicates positive relationship, - indicates negative  
632 relationship, x indicates no relationship, an empty cell means that the trait was not tested. In  
633 all cases '+' indicates a greater capacity to undergo range shifts. Thus, for AC '+' corresponds to  
634 positive population growth, for ER '+' corresponds to low extinction risk.

635 For each of the trait categories, a '+' sign indicates that the relationship between trait and  
636 metric indicates the following increases range-shift capacities. Site (in)fidelity: a high likelihood  
637 of leaving the natal location; movement: strong movement ability; avoidance of small  
638 population sizes: ability to avoid small population effects; persistence in unfavourable climatic  
639 conditions: ability to persist in unfavourable conditions; ecological generalization: ecological  
640 generalists; reproductive strategy: an 'r' strategy; competitive ability: strong competitors (see  
641 Box 2 for more details). Predictive traits and further rationales for each of the seven categories  
642 are detailed in Appendix S1 (section B).