

ACTA ARITHMETICA
161.4 (2013)

Mean value theorems for L -functions over prime polynomials for the rational function field

by

JULIO C. ANDRADE (Providence, RI)
and JONATHAN P. KEATING (Bristol)

1. Introduction. It is a much studied problem in analytic number theory to obtain asymptotic formulae for the moments of families of L -functions. For the family of quadratic Dirichlet L -functions $L(s, \chi_d)$, where χ_d is a real primitive Dirichlet character modulo $|d|$ defined by the Kronecker symbol $\chi_d(n) = \left(\frac{d}{n}\right)$, the problem is to establish asymptotics for

$$(1.1) \quad \sum_{d \leq X}^* L(1/2, \chi_d)^k,$$

in the limit as $X \rightarrow \infty$ and where the sum is over positive fundamental discriminants d (the corresponding sums over negative values of d are also of interest). For $k = 1, 2$, Jutila [8] established the asymptotic formulae

$$(1.2) \quad \sum_{d \leq X}^* L(1/2, \chi_d) \sim c_1 X \log X$$

and

$$(1.3) \quad \sum_{d \leq X}^* L(1/2, \chi_d)^2 \sim c_2 X (\log X)^3,$$

where c_1 and c_2 are computable constants given in terms of Euler products and factors involving the Riemann zeta function. For $k = 3$, Soundararajan [12] proved that

$$(1.4) \quad \sum_{d \leq X}^* L(1/2, \chi_{8d})^3 \sim c_3 X (\log X)^6,$$

2010 *Mathematics Subject Classification*: Primary 11G20; Secondary 11M38, 11M50, 14G10.

Key words and phrases: finite fields, function fields, hyperelliptic curves, moments of quadratic Dirichlet L -functions, prime polynomials.

where d is an odd, square-free and positive number, so that χ_{8d} is a real, even primitive Dirichlet character with conductor $8d$, and c_3 is a constant. Recently, Soundararajan and Young [13] established under the Generalized Riemann Hypothesis an asymptotic formula for the fourth power moment for this family of L -functions, i.e.,

$$(1.5) \quad \sum_{d \leq X}^* L(1/2, \chi_{8d})^4 \sim c_4 X (\log X)^{10},$$

where c_4 is a computable constant. No other asymptotic values are known for the mean values of quadratic Dirichlet L -functions at the centre of the critical strip.

Using results from Random Matrix Theory, Keating and Snaith [9] have put forward a conjecture for the leading order asymptotic for all moments of quadratic Dirichlet L -functions which agrees with the results listed above.

CONJECTURE 1.1 (Keating–Snaith). *For k fixed with $\Re(k) \geq 0$, as*

$$(1.6) \quad \frac{1}{X^*} \sum_{0 < d \leq X}^* L(1/2, \chi_{8d})^k \sim a_{k,Sp} \frac{G(k+1)\sqrt{\Gamma(k+1)}}{\sqrt{G(2k+1)\Gamma(2k+1)}} (\log X)^{k(k+1)/2}$$

as $X \rightarrow \infty$, where

$$a_{k,Sp} = 2^{-k(k+2)/2} \prod_{p \geq 3} \frac{(1-1/p)^{k(k+1)/2}}{1+1/p} \left(\frac{(1-1/\sqrt{p})^{-k} + (1+1/\sqrt{p})^{-k}}{2} + \frac{1}{p} \right)$$

and $G(z)$ is Barnes’ G -function.

Conjectures for the lower order terms are presented in [5] and [6].

A similar problem involving moments of quadratic Dirichlet L -functions was considered by Goldfeld and Viola [7], who have conjectured an asymptotic formula for

$$(1.7) \quad \sum_{\substack{p \leq X \\ p \equiv 3 \pmod{4}}} L(1/2, \chi_p),$$

where $\chi_p(n) = \left(\frac{n}{p}\right)$ is defined by the Legendre symbol. In this context Jutila [8] established the following asymptotic formula:

$$(1.8) \quad \sum_{\substack{p \leq X \\ p \equiv 3 \pmod{4}}} (\log p) L(1/2, \chi_p) = \frac{1}{4} X \log X + O(X(\log X)^\epsilon).$$

It is natural to ask about higher moments for the family of quadratic Dirichlet L -functions associated to χ_p . This problem has the same flavour as that involving the mean values of quadratic Dirichlet L -functions over fundamental discriminants and we formulate it as follows:

PROBLEM 1.2. *Establish asymptotic formulas for*

$$(1.9) \quad \sum_{\substack{p \leq X \\ p \equiv 3 \pmod{4}}} L(1/2, \chi_p)^k$$

when $X \rightarrow \infty$ and $k > 1$.

In this paper we study the function field analogue of this problem in the same spirit as the recent result obtained in [3] for the first moment of quadratic Dirichlet L -functions over the rational function field $\mathbb{F}_q(T)$. Our aim is to obtain asymptotic formulae for the first and second moments for the function field analogue of Problem 1.2 as developed in the next section. Higher moments are studied in [2].

2. Statement of results. Before stating our main results we establish some notation and some preliminary facts about quadratic Dirichlet L -functions for function fields.

2.1. Zeta function of curves. We start with \mathbb{F}_q denoting a finite field of odd cardinality, $A = \mathbb{F}_q[T]$ the polynomials in the variable T with coefficients in \mathbb{F}_q , and $k = \mathbb{F}_q(T)$ the rational function field over \mathbb{F}_q . Let C be any smooth, projective, geometrically connected curve of genus $g \geq 1$ defined over the finite field \mathbb{F}_q . Artin [4] defined the zeta function of the curve C as

$$(2.1) \quad Z_C(u) := \exp\left(\sum_{n=1}^{\infty} N_n(C) \frac{u^n}{n}\right), \quad |u| < \frac{1}{q},$$

with $N_n(C) := \text{Card}(C(\mathbb{F}_{q^n}))$ the number of points on C where the coordinates are in a field extension \mathbb{F}_{q^n} of \mathbb{F}_q of degree $n \geq 1$. It turns out that, as shown by Weil [14], the zeta function associated to C is a rational function of the form

$$(2.2) \quad Z_C(u) = \frac{L_C(u)}{(1-u)(1-qu)},$$

where $L_C(u) \in \mathbb{Z}[u]$ is a polynomial of degree $2g$ that satisfies the functional equation

$$(2.3) \quad L_C(u) = (qu^2)^g L_C\left(\frac{1}{qu}\right).$$

The Riemann Hypothesis for curves over finite fields, established by Weil [14], asserts that the zeros of $L_C(u)$ all lie on the circle $|u| = q^{-1/2}$, i.e.,

$$(2.4) \quad L_C(u) = \prod_{j=1}^{2g} (1 - \alpha_j u) \quad \text{with } |\alpha_j| = \sqrt{q} \text{ for all } j.$$

2.2. Essential facts about $\mathbb{F}_q[T]$. In this paper we denote the *norm* of a polynomial $f \in A$ by $|f| := q^{\deg(f)}$ for $f \neq 0$ and $|f| = 0$ for $f = 0$, and we call a monic irreducible polynomial $P \in A$ a *prime polynomial*.

The zeta function of $A = \mathbb{F}_q[T]$ will be denoted by $\zeta_A(s)$ and is defined in the following way:

$$(2.5) \quad \zeta_A(s) := \sum_{\substack{f \in A \\ f \text{ monic}}} \frac{1}{|f|^s} = \prod_{\substack{P \text{ monic} \\ \text{irreducible}}} (1 - |P|^{-s})^{-1}, \quad \Re(s) > 1.$$

In this case the zeta function $\zeta_A(s)$ is simply given by

$$(2.6) \quad \zeta_A(s) = \frac{1}{1 - q^{1-s}}.$$

The fact that this has a simple pole and no zeros leads to the analogue of the Prime Number Theorem for polynomials in $A = \mathbb{F}_q[T]$:

THEOREM 2.1 (Prime Polynomial Theorem). *If $\pi_A(n)$ denotes the number of monic irreducible polynomials in A of degree n , then*

$$(2.7) \quad \pi_A(n) = \frac{q^n}{n} + O\left(\frac{q^{n/2}}{n}\right).$$

2.3. Quadratic Dirichlet L -function for χ_P . Let $P \in A$ be a monic irreducible polynomial. We denote by χ_P the quadratic character defined in terms of the quadratic residue symbol for $\mathbb{F}_q[T]$:

$$(2.8) \quad \chi_P(f) = \left(\frac{P}{f}\right),$$

where $f \in A$. For more details see [10, Chapters 3, 4]. We will make use of the quadratic reciprocity law for polynomials in A :

THEOREM 2.2 (Quadratic reciprocity). *Let $A, B \in \mathbb{F}_q[T]$ be relatively prime and $A \neq 0$ and $B \neq 0$. Then*

$$(2.9) \quad \left(\frac{A}{B}\right) = \left(\frac{B}{A}\right) (-1)^{((q-1)/2) \deg(A) \deg(B)} = \left(\frac{B}{A}\right) (-1)^{((|A|-1)/2)((|B|-1)/2)}.$$

The L -function attached to the character χ_P is defined by

$$(2.10) \quad L(s, \chi_P) := \sum_{\substack{f \in A \\ f \text{ monic}}} \frac{\chi_P(f)}{|f|^s} = \prod_{\substack{Q \text{ monic} \\ \text{irreducible}}} \left(1 - \frac{\chi_P(Q)}{|Q|^s}\right)^{-1}, \quad \Re(s) > 1.$$

Henceforth we consider P to be a monic irreducible polynomial such that $\deg(P)$ is odd and $q \equiv 1 \pmod{4}$. Then [10, Propositions 4.3, 14.6 and 17.7] $L(s, \chi_P)$ is a polynomial in $u = q^{-s}$ of degree $\deg(P) - 1$ and

$$(2.11) \quad L(s, \chi_P) = \mathcal{L}(u, \chi_P) = L_{C_P}(u),$$

where $L_{C_P}(u)$ is the numerator of the zeta function associated to the hyperelliptic curve given in affine form by

$$(2.12) \quad C_P : y^2 = P(T),$$

with

$$(2.13) \quad P(T) = T^{2g+1} + a_{2g}T^{2g} + \dots + a_1T + a_0$$

a monic irreducible polynomial in A of degree $2g + 1$.

The following proposition is quoted from Rudnick [11] and the main ingredient to establish it is the Riemann Hypothesis for curves,

PROPOSITION 2.3. *If $f \in A$ is monic, $\deg(f) > 0$ and f is not a perfect square then*

$$(2.14) \quad \left| \sum_{\substack{P \text{ prime} \\ \deg(P)=n}} \left(\frac{f}{P} \right) \right| \ll \frac{\deg(f)}{n} q^{n/2}.$$

2.4. The main results. We now present the main results of this paper.

THEOREM 2.4. *Let \mathbb{F}_q be a fixed finite field of odd cardinality with $q \equiv 1 \pmod{4}$. Then for every $\varepsilon > 0$ we have*

$$(2.15) \quad \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} (\log_q |P|) L(1/2, \chi_P) = \frac{|P|}{2} (\log_q |P| + 1) + O(|P|^{3/4+\varepsilon}).$$

This theorem also appears as part of the Ph.D thesis [1] of the first author. This is the exact function field analogue of Jutila’s result (1.8) for number fields. Note that the function field theorem above has a saving in the error term when compared with the number field result (1.8).

THEOREM 2.5. *Using the same notation as before, for a fixed finite field \mathbb{F}_q we have*

$$(2.16) \quad \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} L(1/2, \chi_P)^2 = \frac{1}{24} \frac{1}{\zeta_A(2)} |P| (\log_q |P|)^2 + O(|P| (\log_q |P|)).$$

We have the following corollary:

COROLLARY 2.6.

$$(2.17) \quad \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1 \\ L(1/2, \chi_P) \neq 0}} 1 \gg \frac{|P|}{(\log_q |P|)^2}.$$

Proof. From Theorems 2.4 and 2.5 we have

$$(2.18) \quad \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} L(1/2, \chi_P) \sim k_1 |P|$$

and

$$(2.19) \quad \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} L(1/2, \chi_P)^2 \sim k_2 |P| (\log_q |P|)^2,$$

where k_1 and k_2 are the constants given in the above theorems. By the Cauchy–Schwarz inequality it follows that the number of monic irreducible polynomials P with $\deg(P) = 2g + 1$ such that $L(1/2, \chi_P) \neq 0$ exceeds the ratio of the square of the quantity in (2.18) to the quantity in (2.19). ■

3. The first moment. Setting $D = P$ in Lemma 3.3 from [3], we may write $L(1/2, \chi_P)$ as

$$(3.1) \quad L(1/2, \chi_P) = \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n}} \chi_P(f_1) q^{-n/2} + \sum_{m=0}^{g-1} \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2)=m}} \chi_P(f_2) q^{-m/2}.$$

We need to average both double sums on the right-hand side of (3.1) over monic irreducible polynomials of degree $2g + 1$. However, they are clearly related and we will only need to calculate one of them to obtain the result for the other. Therefore we will focus on the average of the first double sum in (3.1). We can write this as

$$(3.2) \quad \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n}} \chi_P(f_1) q^{-n/2} \\ = \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 \neq \square}} \chi_P(f_1) q^{-n/2} + \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 = \square}} \chi_P(f_1) q^{-n/2}.$$

3.1. Square contributions—the main term. In this section we focus our attention on the average of the first double sum on the right-hand side of (3.2). The main result is

PROPOSITION 3.1.

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 = \square}} \chi_P(f_1) q^{-n/2} = \frac{|P|}{\log_q |P|} \left(\left[\frac{g}{2} \right] + 1 \right) + O\left(\frac{\sqrt{|P|}}{\log_q |P|} g \right),$$

where $[x]$ denotes the integer part of x .

Proof. We have

$$\begin{aligned}
 \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1=\square}} \chi_P(f_1)q^{-n/2} &= \sum_{\substack{n=0 \\ 2|n}}^g q^{-n/2} \sum_{\substack{l \text{ monic} \\ \deg(l)=n/2}} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \chi_P(l^2) \\
 &= \sum_{\substack{n=0 \\ 2|n}}^g q^{-n/2} \sum_{\substack{l \text{ monic} \\ \deg(l)=n/2}} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1 \\ (P,l)=1}} 1 \\
 &= \sum_{\substack{n=0 \\ 2|n}}^g q^{-n/2} \sum_{\substack{l \text{ monic} \\ \deg(l)=n/2}} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} 1,
 \end{aligned}$$

where we obtain the last line from the fact that $\deg(P) = 2g + 1 > \deg(l)$. Making use of the Prime Polynomial Theorem 2.1 we can write

$$\begin{aligned}
 \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1=\square}} \chi_P(f_1)q^{-n/2} &= \sum_{\substack{n=0 \\ 2|n}}^g q^{-n/2} \sum_{\substack{l \text{ monic} \\ \deg(l)=n/2}} \left(\frac{q^{2g+1}}{2g+1} + O\left(\frac{q^{g+1/2}}{2g+1}\right) \right) \\
 &= \frac{q^{2g+1}}{2g+1} \sum_{m=0}^{\lfloor g/2 \rfloor} 1 + O\left(\frac{q^{g+1/2}}{2g+1} \sum_{m=0}^{\lfloor g/2 \rfloor} 1\right) \\
 &= \frac{|P|}{\log_q |P|} \left(\left[\frac{g}{2} \right] + 1 \right) + O\left(\frac{\sqrt{|P|}}{\log_q |P|} g\right). \blacksquare
 \end{aligned}$$

In an analogous way we can prove

PROPOSITION 3.2.

$$\begin{aligned}
 \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{m=0}^{g-1} \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2)=m \\ f_2=\square}} \chi_P(f_2)q^{-m/2} &= \frac{|P|}{\log_q |P|} \left(\left[\frac{g-1}{2} \right] + 1 \right) + O\left(\frac{\sqrt{|P|}}{\log_q |P|} g\right).
 \end{aligned}$$

3.2. Contributions of non-squares. In this section we prove the following result.

PROPOSITION 3.3.

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 \neq \square}} \chi_P(f_1)q^{-n/2} = O\left(\frac{q^{3g/2+1/2}}{\log_q |P|} g\right).$$

Proof. Let $f_1 \in \mathbb{F}_q[T]$ be a fixed monic non-square polynomial such that $\deg(f_1) < \deg(P) = 2g + 1$. By the quadratic reciprocity law, Theorem 2.2, we have

$$(3.3) \quad \left(\frac{P}{f_1}\right) = (-1)^{\frac{q-1}{2}(2g+1)\deg(f_1)} \left(\frac{f_1}{P}\right).$$

Note that the sign $(-1)^{\frac{q-1}{2}(2g+1)\deg f_1}$ is the same for all monic irreducible polynomials P of degree $2g + 1$, so

$$(3.4) \quad \left| \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \left(\frac{P}{f_1}\right) \right| = \left| \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \left(\frac{f_1}{P}\right) \right|.$$

Thus we can write

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 \neq \square}} \chi_P(f_1)q^{-n/2} \ll \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 \neq \square}} q^{-n/2} \left| \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \left(\frac{f_1}{P}\right) \right|$$

and using the bound for character sums over prime polynomials given in Proposition 2.3 we have

$$\begin{aligned} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n \\ f_1 \neq \square}} \chi_P(f_1)q^{-n/2} &\ll \sum_{n=0}^g q^{-n/2} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n}} n \frac{q^{g+1/2}}{2g+1} \\ &\ll \frac{\sqrt{|P|}}{\log_q |P|} gq^{g/2}. \blacksquare \end{aligned}$$

We can prove a corresponding estimate for the dual sum in (3.1) using the same approach. In the end we have

PROPOSITION 3.4.

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{m=0}^{g-1} \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2)=m \\ f_2 \neq \square}} \chi_P(f_2)q^{-m/2} = O\left(\frac{q^{3g/2+1/2}}{\log_q |P|} g\right).$$

3.3. Proof of the theorem for the first moment. We are now in a position to prove Theorem 2.4.

Proof of Theorem 2.4. We can write

$$\begin{aligned} & \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} (\log_q |P|) L(1/2, \chi_P) \\ = & \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} (\log_q |P|) \left(\sum_{n=0}^g \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n}} \chi_P(f_1) q^{-n/2} + \sum_{m=0}^{g-1} \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2)=m}} \chi_P(f_2) q^{-m/2} \right). \end{aligned}$$

Making use of Propositions 3.1–3.4 we establish that

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} (\log_q |P|) L(1/2, \chi_P) = |P| \left(\left[\frac{g}{2} \right] + \left[\frac{g-1}{2} \right] + 2 \right) + O(q^{3g/2+1/2}g).$$

and using

$$(3.5) \quad \left[\frac{g}{2} \right] + \left[\frac{g-1}{2} \right] = g - 1$$

and

$$(3.6) \quad g + 1 = \frac{\log_q |P|}{2} + \frac{1}{2}$$

we conclude the proof of the theorem. ■

4. The second moment. In this section we prove Theorem 2.5.

4.1. Secondary lemmas. We will need some auxiliary lemmas before we proceed to the proof of Theorem 2.5.

The starting point is a representation for $L(1/2, \chi_P)^2$ which can be viewed as the analogue of the approximate functional equation for a quadratic Dirichlet L -function (Lemma 3 in [8]). In this case there is no error term and the formula is exact.

LEMMA 4.1. *Let χ_P be the quadratic Dirichlet character associated to the monic irreducible polynomial $P \in A$. Then*

$$(4.1) \quad L(1/2, \chi_P)^2 = \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g}} \frac{\chi_P(f_1) d(f_1)}{|f_1|^{1/2}} + \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2) \leq 2g-1}} \frac{\chi_P(f_2) d(f_2)}{|f_2|^{1/2}},$$

where $d(f)$ is the divisor function for polynomials $f \in A$ (see [10, p. 15]).

Proof. We have $L(s, \chi_P) = L_{C_P}(u)$. So

$$(4.2) \quad L_{C_P}(u)^2 = ((qu^2)^g)^2 L_{C_P}\left(\frac{1}{qu}\right)^2.$$

Writing $L_{C_P}(u)^2 = \sum_{n=0}^{4g} a_n u^n$ we obtain

$$(4.3) \quad \begin{aligned} \sum_{n=0}^{4g} a_n u^n &= (qu^2)^g (qu^2)^g \sum_{m=0}^{4g} a_m q^{-m} u^{-m} = \sum_{m=0}^{4g} a_m q^{2g-m} u^{4g-m} \\ &= \sum_{k=0}^{4g} a_{4g-k} q^{k-2g} u^k. \end{aligned}$$

Equating coefficients we find that $a_n = a_{4g-n} q^{n-2g}$ and so we can write

$$(4.4) \quad L_{C_P}(u)^2 = \sum_{n=0}^{2g} a_n u^n + ((qu^2)^g)^2 \sum_{m=0}^{2g-1} a_m q^{-m} u^{-m}.$$

From $L(s, \chi_P)^2$ we see that the coefficients a_n are given by

$$(4.5) \quad a_n = \sum_{\substack{f \text{ monic} \\ \deg(f)=n}} \chi_P(f) d(f),$$

where

$$(4.6) \quad d(f) = \sum_{\substack{h_1 h_2 = f \\ h_1, h_2 \text{ monic}}} 1.$$

Therefore writing $s = 1/2$, i.e. $u = q^{-1/2}$, in (4.4) proves the lemma. ■

Our next lemma is quoted from Rosen [10, Proposition 2.5].

LEMMA 4.2.

$$(4.7) \quad \sum_{\substack{f \text{ monic} \\ \deg(f)=n}} d(f) \ll q^n n.$$

The next lemma is a minor modification of Theorem 17.4 in [10].

LEMMA 4.3. *Let $f : A^+ \rightarrow \mathbb{C}$ and let $\zeta_f(s)$ be the corresponding Dirichlet series. Suppose this series converges absolutely in the region $\Re(s) > 1$ and is holomorphic in the region $\{s \in B : \Re(s) = 1\}$ except for a single pole of order r at $s = 1$, where A^+ denotes the set of monic polynomials in $\mathbb{F}_q[T]$ and*

$$B = \left\{ s \in \mathbb{C} : -\frac{\pi i}{\log(q)} \leq \Im(s) \leq \frac{\pi i}{\log(q)} \right\}.$$

Let $\alpha = \lim_{s \rightarrow 1} (s - 1)^r \zeta_f(s)$. Then there is a $\delta < 1$ and constants c_{-i} with

$1 \leq i \leq r$ such that

$$(4.8) \quad \sum_{\deg(D)=n} f(D) = q^n \left(\sum_{i=1}^r c_{-i} \binom{n+i-1}{i-1} (-q)^i \right) + O(q^{\delta n}).$$

The sum in parenthesis is a polynomial in n of degree $r-1$ with leading term

$$\frac{\log(q)^r}{(r-1)!} \alpha n^{r-1}.$$

LEMMA 4.4. Let f be a monic polynomial in $A = \mathbb{F}_q[T]$. Then

$$(4.9) \quad \sum_{\substack{f \text{ monic} \\ \deg(f)=n}} d(f^2) = \frac{1}{2} \frac{1}{\zeta_A(2)} q^n n^2 + O(q^n n).$$

Proof. We consider the Dirichlet series associated to $d(f^2)$:

$$\begin{aligned} \zeta_f(s) &= \sum_{f \text{ monic}} \frac{d(f^2)}{|f|^s} = \prod_{\substack{P \text{ monic} \\ \text{irreducible}}} \left(1 + \frac{d(P^2)}{|P|^s} + \frac{d(P^4)}{|P|^{2s}} + \dots \right) \\ &= \prod_{\substack{P \text{ monic} \\ \text{irreducible}}} \left(1 + \left(\frac{-3}{|P|^s(|P|^s-1)^2} + \frac{1}{(|P|^s-1)^2} \right) \right) = \frac{\zeta_A(s)^3}{\zeta_A(2s)}. \end{aligned}$$

From (2.5) the sum converges absolutely for $\Re(s) > 1$, is holomorphic on the disc $\{u = q^{-s} \in \mathbb{C} : |u| \leq q^{-\delta}\}$ for some $\delta < 1$, and $\zeta_f(s)$ has a pole of order 3 at $s = 1$. We now apply Lemma 4.3 to obtain

$$(4.10) \quad \sum_{\substack{f \text{ monic} \\ \deg(f)=n}} d(f^2) = \frac{(\log q)^3}{2} \alpha q^n n^2 + O(q^n n),$$

where

$$(4.11) \quad \alpha = \lim_{s \rightarrow 1} (s-1)^3 \frac{\zeta_A(s)^3}{\zeta_A(2s)} = \frac{q-1}{q(\log q)^3}. \blacksquare$$

4.2. Preparation for the proof. From Lemma 4.1, $L(1/2, \chi_P)^2$ can be written as two similar sums. Our main aim is to average, over the prime polynomials, the first sum on the right-hand side of (4.1). We start by writing

$$(4.12) \quad \begin{aligned} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g}} \frac{\chi_P(f_1) d(f_1)}{|f_1|^{1/2}} \\ = \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 = \square}} \frac{\chi_P(f_1) d(f_1)}{|f_1|^{1/2}} + \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 \neq \square}} \frac{\chi_P(f_1) d(f_1)}{|f_1|^{1/2}}. \end{aligned}$$

4.3. The main term. The following proposition is established in this section.

PROPOSITION 4.5.

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 = \square}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} = \frac{1}{12} \frac{1}{\zeta_A(2)} \frac{|P|}{\log_q |P|} g(g+1)(2g+1) + O\left(\frac{|P|}{\log_q |P|} g^2\right).$$

Proof. We have

$$\begin{aligned} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 = \square}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} &= \sum_{n=0}^{2g} q^{-n/2} \sum_{\substack{f_1 = \square \\ \deg(f_1)=n}} d(f_1) \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \chi_P(f_1) \\ &= \sum_{m=0}^g q^{-m} \sum_{\substack{l \text{ monic} \\ \deg(l)=m}} d(l^2) \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} 1. \end{aligned}$$

We again make use of the Prime Polynomial Theorem 2.1 to obtain

$$\begin{aligned} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 = \square}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} &= \frac{|P|}{\log_q |P|} \sum_{m=0}^g q^{-m} \sum_{\substack{l \text{ monic} \\ \deg(l)=m}} d(l^2) + O\left(\frac{\sqrt{|P|}}{\log_q |P|} \sum_{m=0}^g q^{-m} \sum_{\substack{l \text{ monic} \\ \deg(l)=m}} d(l^2)\right). \end{aligned}$$

Invoking Lemma 4.3 we obtain the following equation:

$$\begin{aligned} \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 = \square}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} &= \frac{|P|}{\log_q |P|} \frac{1}{2} \frac{1}{\zeta_A(2)} \sum_{m=0}^g m^2 + O\left(\frac{|P|}{\log_q |P|} \sum_{m=0}^g m\right) + O\left(\frac{\sqrt{|P|}}{\log_q |P|} \sum_{m=0}^g m^2\right) \\ &= \frac{|P|}{\log_q |P|} \frac{1}{12} \frac{1}{\zeta_A(2)} g(g+1)(2g+1) + O\left(\frac{|P|}{\log_q |P|} g^2\right). \blacksquare \end{aligned}$$

In a similar way we can prove

PROPOSITION 4.6.

$$\begin{aligned} & \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2) \leq 2g-1 \\ f_2 \neq \square}} \frac{\chi_P(f_2)d(f_2)}{|f_2|^{1/2}} \\ &= \frac{1}{12} \frac{1}{\zeta_A(2)} \frac{|P|}{\log_q |P|} \left[\frac{2g-1}{2} \right] \left(1 + \left[\frac{2g-1}{2} \right] \right) \left(1 + 2 \left[\frac{2g-1}{2} \right] \right) \\ & \quad + O\left(\frac{|P|}{\log_q |P|} g^2 \right). \end{aligned}$$

4.4. Contributions of non-squares. The main result in this section is the following proposition.

PROPOSITION 4.7.

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 \neq \square}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} = O(|P|g).$$

Proof. We have

$$\begin{aligned} & \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 \neq \square}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} \ll \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g \\ f_1 \neq \square}} \frac{d(f_1)}{|f_1|^{1/2}} \left| \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \left(\frac{f_1}{P} \right) \right| \\ & \ll \frac{\sqrt{|P|}}{2g+1} \sum_{n=0}^{2g} \frac{n}{q^{n/2}} \sum_{\substack{f_1 \text{ monic} \\ \deg(f_1)=n}} d(f_1) \\ & \ll \frac{\sqrt{|P|}}{2g+1} \sum_{n=0}^{2g} n^2 q^{n/2} \ll |P|g, \end{aligned}$$

where we have used Proposition 2.3 in the first line and Lemma 4.2 in the second. ■

Similarly we have

PROPOSITION 4.8.

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2) \leq 2g-1 \\ f_2 \neq \square}} \frac{\chi_P(f_2)d(f_2)}{|f_2|^{1/2}} = O(|P|g).$$

4.5. Proof of the theorem for the second moment. We are finally in a position to prove Theorem 2.5.

Proof of Theorem 2.5. We can write

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} L(1/2, \chi_P)^2 = \sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} \left(\sum_{\substack{f_1 \text{ monic} \\ \deg(f_1) \leq 2g}} \frac{\chi_P(f_1)d(f_1)}{|f_1|^{1/2}} + \sum_{\substack{f_2 \text{ monic} \\ \deg(f_2) \leq 2g-1}} \frac{\chi_P(f_2)d(f_2)}{|f_2|^{1/2}} \right).$$

Making use of Propositions 4.5–4.8 we establish that

$$\sum_{\substack{P \text{ monic} \\ \text{irreducible} \\ \deg(P)=2g+1}} L(1/2, \chi_P)^2 = \frac{1}{12} \frac{1}{\zeta_A(2)} \frac{|P|}{\log_q |P|} \times \left[g(g+1)(2g+1) + \left[\frac{2g-1}{2} \right] \left(1 + \left[\frac{2g-1}{2} \right] \right) \left(1 + 2 \left[\frac{2g-1}{2} \right] \right) \right] + O(|P|g).$$

We use

$$\left[\frac{2g-1}{2} \right] \left(1 + \left[\frac{2g-1}{2} \right] \right) \left(1 + 2 \left[\frac{2g-1}{2} \right] \right) = (g-1)g(2g-1)$$

and

$$g(g+1)(2g+1) + (g-1)g(2g-1) = 4g^3 + O(g)$$

and after some simple arithmetical manipulations this gives the desired formula. ■

Acknowledgments. JCA is supported by a NSF Postdoctoral Grant and an ICERM-Brown University Postdoctoral Research Fellowship. JPK is sponsored by the Leverhulme Trust and the Air Force Office of Scientific Research, Air Force Material Command, USAF, under grant number FA8655-10-1-3088. The U.S. Government is authorized to reproduce and distribute reprints for Governmental purpose notwithstanding any copyright notation thereon.

We would like to thank Professor Zeév Rudnick for suggesting the problems tackled in this paper, and Professors Michael Rosen and Jeffrey Hoffstein for helpful and interesting discussions. We also wish to thank the referee for a careful reading of the paper and for the comments provided.

References

[1] J. C. Andrade, *Random matrix theory and L-functions in function fields*, Ph.D. thesis, Univ. of Bristol, Bristol, 2012.

- [2] J. C. Andrade, *Higher moments for the prime hyperelliptic ensemble*, in progress, 2013.
- [3] J. C. Andrade and J. P. Keating, *The mean value of $L(1/2, \chi)$ in the hyperelliptic ensemble*, *J. Number Theory* 132 (2012), 2793–2816.
- [4] E. Artin, *Quadratische Körper im Gebiete der höheren Kongruenzen I, II*, *Math. Z.* 19 (1924), 153–206, 207–246
- [5] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein and N. C. Snaith, *Integral moments of L -functions*, *Proc. London Math. Soc.* 91 (2005), 33–104.
- [6] A. Diaconu, D. Goldfeld and J. Hoffstein, *Multiple Dirichlet series and moments of zeta and L -functions*, *Compos. Math.* 139 (2003), 297–360.
- [7] D. Goldfeld and C. Viola, *Mean values of L -functions associated to elliptic, Fermat and other curves at the centre of the critical strip*, *J. Number Theory* 11 (1979), 305–320.
- [8] M. Jutila, *On the mean value of $L(1/2, \chi)$ for real characters*, *Analysis* 1 (1981), 149–161.
- [9] J. P. Keating and N. C. Snaith, *Random matrix theory and L -functions at $s = 1/2$* , *Comm. Math. Phys.* 214 (2000), 91–110.
- [10] M. Rosen, *Number Theory in Function Fields*, *Grad. Texts in Math.* 210, Springer, New York, 2002.
- [11] Z. Rudnick, *Traces of high powers of the Frobenius class in the hyperelliptic ensemble*, *Acta Arith.* 143 (2010), 81–99.
- [12] K. Soundararajan, *Nonvanishing of quadratic Dirichlet L -functions at $s = 1/2$* , *Ann. of Math.* 152 (2000), 447–488.
- [13] K. Soundararajan and M. P. Young, *The second moment of quadratic twists of modular L -functions*, *J. Eur. Math. Soc.* 12 (2010), 1097–1116.
- [14] A. Weil, *Sur les Courbes Algébriques et les Variétés qui s'en Déduisent*, Hermann, Paris, 1948.

Julio C. Andrade
Institute for Computational and Experimental
Research in Mathematics (ICERM)
Brown University
121 South Main Street
Providence, RI 02903, U.S.A.
E-mail: julio_andrade@brown.edu

Jonathan P. Keating
School of Mathematics
University of Bristol
Bristol BS8 1TW, UK
E-mail: j.p.keating@bristol.ac.uk

*Received on 12.3.2013
and in revised form on 13.6.2013*

(7369)

